

Bronchoscopy and Biopsy Pre-Procedure [1026]

General

Case Request (Single Response)

Case request operating room Scheduling/ADT, Scheduling/ADT

Nursing

Vital Signs

Vital signs - T/P/R/BP Routine, Every 15 min
Moderate sedation protocol monitoring, Pre-op

Consent

- Complete Consent For Bronchoscopy Routine, Once
Consent For: Bronchoscopy
Pre-op
- Complete Consent For Bronchoscopy with biopsy Routine, Once
Consent For: Bronchoscopy with biopsy
Pre-op
- Complete Consent For Bronchoscopy with lavage Routine, Once
Consent For: Bronchoscopy with lavage
Pre-op
- Complete Consent For Bronchoscopy with lavage, possible balloon dilation, stent revision, removal or placement Routine, Once
Consent For: Bronchoscopy with lavage, possible balloon dilation, stent revision, removal or placement
Pre-op
- Complete Consent For Bronchoscopy with biopsy, BAL studies, dilation or stent removal Routine, Once
Consent For: Bronchoscopy with biopsy, BAL studies, dilation or stent removal
Pre-op
- Complete consent for Fiducial marker placement Routine, Once
Procedure:
Diagnosis/Condition:
Physician:
Pre-op
- Complete consent for Bronchial Thermoplasty Routine, Once
Procedure:
Diagnosis/Condition:
Physician:
Pre-op
- Complete consent for Routine, Once
Procedure:
Diagnosis/Condition:
Physician:
Pre-op

Notify

- Notify Physician upon patient arrival to bronchoscopy suite Routine, Until discontinued, Starting S, Pre-op
- Notify Physician if O2 Sat less than 90% Routine, Until discontinued, Starting S, Pre-op

Diet

- NPO - except meds with sip of water after midnight Diet effective midnight, Starting S+1 at 12:01 AM
NPO: Except meds
Pre-Operative fasting options:
with sip of water after midnight, Pre-op

<input type="checkbox"/> NPO - except meds with sip of water after breakfast	Diet effective _____ NPO: Except meds Pre-Operative fasting options: with sip of water after breakfast, Pre-op
<input type="checkbox"/> NPO	Conditional Frequency NPO: Pre-Operative fasting options: 5 hours prior to procedure, Pre-op

IV Fluids

Peripheral IV Access

<input checked="" type="checkbox"/> Initiate and maintain IV	
<input checked="" type="checkbox"/> Insert peripheral IV	Routine, Once
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care

Medications

Pre-Bronchoscopy Medication

30 minutes prior to bronchoscopy.

<input type="checkbox"/> atropine injection	0.4 mg, intramuscular, once, For 1 Doses, Pre-op 30 minutes prior to bronchoscopy.
<input type="checkbox"/> promethazine (PHENERGAN) injection	12.5 mg, intramuscular, once PRN, nausea, vomiting, For 1 Doses, Pre-op 30 minutes prior to bronchoscopy. For Deep IM Administration.
<input type="checkbox"/> glycopyrrolate (ROBINUL) injection	0.2 mg, intramuscular, once, For 1 Doses, Pre-op
<input type="checkbox"/> lidocaine (XYLOCAINE) 2 % jelly	Topical, once, For 1 Doses, Pre-op
<input type="checkbox"/> ampicillin (OMNIPEN) IV	2 g, intravenous, for 30 Minutes, once, For 1 Doses, Pre-op Start 30 minutes prior to bronchoscopy Reason for Therapy:
<input type="checkbox"/> clindamycin (CLEOCIN) IV	300 mg, intravenous, for 30 Minutes, once, For 1 Doses, Pre-op Start 30 minutes prior to Bronchoscopy. Reason for Therapy:
<input type="checkbox"/> albuterol (PROVENTIL) nebulizer solution	2.5 mg, nebulization, once, For 1 Doses, Pre-op Administer during spirometry Aerosol Delivery Device:
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection	125 mg, intravenous, once, For 1 Doses, Pre-op Administer after spirometry

Intra Bronchoscopy Orders

<input checked="" type="checkbox"/> lidocaine (XYLOCAINE) 40 mg/mL (4 %) inhalation	6 mL, nebulization, once, For 1 Doses, Pre-op Administer immediately before procedure
<input type="checkbox"/> MIDAZolam (VERSED) injection	intravenous, once, For 1 Doses, Pre-op
<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	intravenous, once PRN, severe pain (score 7-10), For 1 Doses, Pre-op

Bronchoscopy Kit Medications

<input type="checkbox"/> sodium chloride (NS) 0.9 % irrigation solution	1,000 mL, Topical, once PRN, bronchoscopy, For 1 Doses, Pre-op
<input type="checkbox"/> lidocaine (XYLOCAINE) 4 % (40 mg/mL) external solution	Topical, once PRN, mild pain (score 1-3), bronchoscopy, For 1 Doses, Pre-op
<input type="checkbox"/> epINEPhrine (ADRENALIN) 1 mg/mL (1 mL) injection	1 mg, Topical, once PRN, anaphylaxis, bronchoscopy, For 1 Doses, Pre-op
<input type="checkbox"/> lidocaine HCl (XYLOCAINE) 2 % solution	15 mL, Topical, once PRN, bronchoscopy, For 1 Doses, Pre-op
<input type="checkbox"/> sodium chloride 0.9 % injection	20 mL, Topical, once PRN, line care, bronchoscopy, For 1 Doses, Pre-op

VTE

Labs

Labs STAT

<input type="checkbox"/>	CBC and differential	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Prothrombin time with INR	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Partial thromboplastin time	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/>	Basic metabolic panel	STAT For 1 Occurrences, Pre-op

Cardiology

Diagnostic Imaging

X-Ray

<input type="checkbox"/>	Chest 1 Vw Portable	Routine, 1 time imaging For 1 , Pre-op
--------------------------	---------------------	--

Other Diagnostic Studies

Respiratory

Respiratory

<input checked="" type="checkbox"/>	Oxygen therapy	Routine, Continuous Device 1: Nasal Cannula Rate in liters per minute: Rate in tenths of a liter per minute: O2 %: Titrate to keep O2 Sat Above: 92% Indications for O2 therapy: 4 to 6 liters per minute, Pre-op
<input type="checkbox"/>	Bedside spirometry	Routine, Once With bronchodilators, Pre-op

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders