

IV Contrast Allergy Prep**Contrast Allergy Prep**

<input type="checkbox"/> methylPREDNISolone (MEDROL) tablet	32 mg, oral, once, For 1 Doses Give 12 hours prior to the scheduled procedure
<input type="checkbox"/> methylPREDNISolone (MEDROL) tablet	32 mg, oral, once, For 1 Doses Give 2 hours prior to the scheduled procedure
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet	60 mg, oral, once, For 1 Doses Give 2 hours prior to the scheduled procedure
<input type="checkbox"/> famotidine (PEPCID) tablet	20 mg, oral, once, For 1 Doses Give 2 hours prior to the scheduled procedure

Contrast Allergy Prep

<input type="checkbox"/> methylPREDNISolone (MEDROL) tablet	32 mg, oral, once, For 1 Doses Give 12 hours prior to the scheduled procedure
<input type="checkbox"/> methylPREDNISolone (MEDROL) tablet	32 mg, oral, once, For 1 Doses Give 2 hours prior to the scheduled procedure
<input type="checkbox"/> famotidine (PEPCID) tablet	20 mg, oral, once, For 1 Doses Give 2 hours prior to the scheduled procedure

Contrast Allergy Prep

<input type="checkbox"/> methylPREDNISolone (MEDROL) tablet	32 mg, oral, once, For 1 Doses Give 12 hours prior to the scheduled procedure
<input type="checkbox"/> methylPREDNISolone (MEDROL) tablet	32 mg, oral, once, For 1 Doses Give 2 hours prior to the scheduled procedure
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet	60 mg, oral, once, For 1 Doses Give 2 hours prior to the scheduled procedure
<input type="checkbox"/> famotidine (PEPCID) tablet	20 mg, oral, once, For 1 Doses Give 2 hours prior to the scheduled procedure