

General

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Post-op
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Post-op
<input type="checkbox"/>	Acute Renal Failure	Post-op
<input type="checkbox"/>	Acute Respiratory Failure	Post-op
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
<input type="checkbox"/>	Anemia	Post-op
<input type="checkbox"/>	Bacteremia	Post-op
<input type="checkbox"/>	Bipolar disorder, unspecified	Post-op
<input type="checkbox"/>	Cardiac Arrest	Post-op
<input type="checkbox"/>	Cardiac Dysrhythmia	Post-op
<input type="checkbox"/>	Cardiogenic Shock	Post-op
<input type="checkbox"/>	Decubitus Ulcer	Post-op
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Post-op
<input type="checkbox"/>	Disorder of Liver	Post-op
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Post-op
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Post-op
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Post-op
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Post-op
<input type="checkbox"/>	Other Alteration of Consciousness	Post-op
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Post-op
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Post-op
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Post-op
<input type="checkbox"/>	Protein-calorie Malnutrition	Post-op
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Post-op
<input type="checkbox"/>	Schizophrenia Disorder	Post-op
<input type="checkbox"/>	Sepsis	Post-op
<input type="checkbox"/>	Septic Shock	Post-op
<input type="checkbox"/>	Septicemia	Post-op
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Post-op

Elective Outpatient, Observation, or Admission (Single Response)

<input type="checkbox"/>	Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

Admission or Observation (Single Response)

Patient has active outpatient status order on file

- | | |
|--|--|
| <input type="checkbox"/> Admit to Inpatient | Diagnosis:
Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Outpatient observation services under general supervision | Diagnosis:
Admitting Physician:
Patient Condition:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Outpatient in a bed - extended recovery | Diagnosis:
Admitting Physician:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Admission (Single Response)

Patient has active status order on file

- | | |
|---|--|
| <input type="checkbox"/> Admit to inpatient | Diagnosis:
Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Transfer (Single Response)

Patient has active inpatient status order on file

- | | |
|---|---|
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Code Status

- | | |
|---|--|
| <input type="checkbox"/> Full Code | Code Status decision reached by:
Post-op |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Does patient have decision-making capacity?
Post-op |

<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

Isolation

<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

Nursing

Vital Signs

<input type="checkbox"/> Vital signs-Per unit protocol	Routine, Per unit protocol, Post-op
<input type="checkbox"/> Vital signs-Q4H	Routine, Every 4 hours For Until specified, Post-op
<input type="checkbox"/> Pulse oximetry	Routine, Every 4 hours Current FIO2 or Room Air: Post-op

Activity

<input type="checkbox"/> Head of bed	Routine, Until discontinued, Starting S Head of bed: 30 degrees If not contraindicated, Post-op
<input type="checkbox"/> Bed rest with bedside commode	Routine, Until discontinued, Starting S Bathroom Privileges: with bedside commode Post-op
<input type="checkbox"/> Up in chair for meals	Routine, Until discontinued, Starting S Specify: Up in chair Additional modifier: for meals Post-op
<input type="checkbox"/> Ambulate with assistance	Routine, 3 times daily Specify: with assistance Post-op
<input type="checkbox"/> Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated Post-op

Nursing

<input checked="" type="checkbox"/> Height and weight	Routine, Once On arrival to unit, Post-op
<input type="checkbox"/> Daily weights	Routine, Daily, Post-op

<input type="checkbox"/>	Measure drainage	Routine, Every 8 hours Type of drain: Record output from drain every 8 hours , Post-op
<input checked="" type="checkbox"/>	Intake and Output	Routine, Every shift, Post-op
<input type="checkbox"/>	Oral care	Routine, Every 4 hours For intubated patients, Post-op
<input type="checkbox"/>	Oral care	Routine, Every shift For non intubated patients, Post-op
<input type="checkbox"/>	Assist with feeding patient	Routine, Until discontinued, Starting S Modifier: Post-op
<input type="checkbox"/>	Peripheral vascular assessment	Routine, Once, Post-op
Line/Drain Care		
<input type="checkbox"/>	Saline lock IV	Routine, Once For 1 Occurrences When patient tolerates oral intake , Post-op
<input type="checkbox"/>	Insert and Maintain Foley	
<input type="checkbox"/>	Insert Foley catheter	Routine, Once Type: Size: Urinometer needed: Post-op
<input type="checkbox"/>	Foley Catheter Care	Routine, Until discontinued, Starting S Orders: Maintain To Gravity 1) Remove Foley cath POD 1 or POD 2 or 2) Activate Nursing Protocol or 3) Document reason for not removing Foley_____. (must be documented on POD 1 or POD 2), Post-op
<input type="checkbox"/>	Chest tube to continuous suction	Routine, Until discontinued, Starting S Level of suction: 20 cm H2O Post-op
<input type="checkbox"/>	Chest tube to water seal	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/>	Drain care	Routine, Until discontinued, Starting S Drain 1: Drain 2: Drain 3: Drain 4: All Drains: Post-op
<input type="checkbox"/>	Nasogastric tube maintenance	Routine, Until discontinued, Starting S Tube Care Orders: Post-op
Wound/Incision Care		
<input type="checkbox"/>	Apply ice pack	Routine, Until discontinued, Starting S Affected area: Waking hours only? Nurse to schedule? Special Instructions: Post-op
<input type="checkbox"/>	Reinforce dressing	Routine, As needed Reinforce with: Post-op
<input type="checkbox"/>	Sitz bath	Routine, Once, Post-op
<input type="checkbox"/>	Surgical/incision site care	Routine, Once Location: Site: Apply: Dressing Type: Open to air? Post-op

<input type="checkbox"/> Wound care orders	Routine, Every 12 hours Wound care to be performed by: Location: Site: Irrigate wound? Apply: Dressing Type: Post-op
<input type="checkbox"/> Provide equipment / supplies at bedside	Routine, Once Supplies: Post-op
<input type="checkbox"/> Wound vac (Not Consult Order)	Routine, Every Mon, Wed, Fri Existing wound vac? Type of Wound: Wound Location: Pressure (mmHg): 125 Therapy Settings: Intensity: Foam Type: Post-op
<input type="checkbox"/> Consult to Wound Ostomy Care Nurse	Reason for consult: Reason for consult: Reason for consult: Reason for consult: Consult for NPWT: Reason for consult: Post-op

Diet

<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options: Post-op
<input type="checkbox"/> Diet- Clear liquid	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op

Notify

<input type="checkbox"/> Notify Physician for vitals:	Routine, Until discontinued, Starting S Temperature greater than: 100.5 Temperature less than: Systolic BP greater than: 160 Systolic BP less than: 90 Diastolic BP greater than: 100 Diastolic BP less than: 50 MAP less than: 60 Heart rate greater than (BPM): 100 Heart rate less than (BPM): 60 Respiratory rate greater than: 25 Respiratory rate less than: 8 SpO2 less than: 92
<input type="checkbox"/> Notify Physician (Specify)	Routine, Until discontinued, Starting S, Post-op

IV Fluids

IV Fluids (Single Response)

<input type="checkbox"/> lactated Ringer's infusion	intravenous, continuous, Post-op
<input type="checkbox"/> sodium chloride 0.9 % infusion	intravenous, continuous, Post-op

<input type="checkbox"/> sodium chloride 0.9 % with potassium chloride 20 mEq/L infusion	intravenous, continuous, Post-op
<input type="checkbox"/> dextrose 5 % and sodium chloride 0.45 % with potassium chloride 20 mEq/L infusion - for NPO Patients	intravenous, continuous, Post-op

Medications

Postoperative Antibiotics: For Patients LESS than or EQUAL to 120 kg (Single Response)

<input type="checkbox"/> cefazolin (ANCEF) IV - For Patients LESS than or EQUAL to 120 kg	2 g, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> cefoxitin (MEFOXIN) IV	2 g, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> ertapenem (INVanz) IV	1 g, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> ampicillin-sulbactam (UNASYN) IV	3 g, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis

Postoperative Antibiotics: For Patients GREATER than 120 kg (Single Response)

<input type="checkbox"/> cefazolin (ANCEF) IV - For Patients GREATER than 120 kg	3 g, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> cefoxitin (MEFOXIN) IV	2 g, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> ertapenem (INVanz) IV	1 g, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> ampicillin-sulbactam (UNASYN) IV	3 g, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis

Postoperative Antibiotics: If Beta-Lactam Allergy

<input type="checkbox"/> metronidazole (FLAGYL) IV	500 mg, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason of Therapy: Surgical Prophylaxis
<input type="checkbox"/> ciprofloxacin (CIPRO) IV	400 mg, intravenous, for 60 Minutes, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis

Postoperative Antibiotics: If MRSA Suspected

<input type="checkbox"/> vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
---	--

Beta-Blockers

<input type="checkbox"/> metoprolol (LOPRESSOR) injection	5 mg, intravenous, every 6 hours, Post-op hold if systolic blood pressure is LESS than 110 and heart rate is LESS than 60 bpm HOLD parameters for this order: Contact Physician if:
<input type="checkbox"/> metoprolol tartrate (LOPRESSOR) tablet	100 mg, oral, 2 times daily at 0600, 1800, Post-op hold if systolic blood pressure is LESS than 110 and heart rate is LESS than 60 bpm HOLD parameters for this order: Contact Physician if:

<input type="checkbox"/> labetalol (NORMODYNE) tablet	200 mg, oral, 2 times daily at 0600, 1800, Post-op HOLD parameters for this order: Contact Physician if:
<input type="checkbox"/> labetalol (NORMODYNE,TRANDATE) injection	intravenous, Post-op

Respiratory

<input type="checkbox"/> albuterol (PROVENTIL HFA;VENTOLIN HFA) inhaler	2 puff, inhalation, Respiratory Therapy - every 6 hours, Post-op
<input type="checkbox"/> tiotropium (SPIRIVA) 18 mcg per inhalation capsule	1 capsule, inhalation, Respiratory Therapy - Daily, Post-op

PRN Mild Pain (Pain Score 1-3) (Single Response)
(adjust dose for renal/liver function and age)

<input type="checkbox"/> acetaminophen (TYLENOL) tablet OR oral solution	"Or" Linked Panel Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op Maximum of 3 grams of acetaminophen per day from all sources. Give the tablet if the patient can tolerate oral medication. (Cirrhosis patients maximum: 2 grams per day from all sources)
<input type="checkbox"/> acetaminophen (TYLENOL)suspension	650 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Use if patient cannot tolerate oral tablet.

PRN Oral for Moderate Pain (Pain Score 4-6): For Patients LESS than 65 years old (Single Response)
(adjust dose for renal/liver function and age)

<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) tablet OR elixir	"Or" Linked Panel Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient is able to tolerate oral medication.
<input type="checkbox"/> acetaminophen-codeine 300 mg-30 mg /12.5 mL solution	12.5 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) Use if patient cannot swallow tablet.
<input type="checkbox"/> HYDROcodone-acetaminophen 5/325 (NORCO) tablet OR elixir	"Or" Linked Panel Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6)
<input type="checkbox"/> HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution	10 mL, oral, every 6 hours PRN, moderate pain (score 4-6)
<input type="checkbox"/> HYDROcodone-acetaminophen 7.5/325 (NORCO) tablet OR elixir	"Or" Linked Panel Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)

[] HYDROcodone-acetaminophen (NORCO) 7.5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient is able to tolerate oral medication.
[] HYDROcodone-acetaminophen (HYCET) 7.5-325 mg/15 mL solution	15 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) Use if patient cannot swallow tablet.
() HYDROcodone-acetaminophen 10/325 (NORCO) tablet OR elixir	"Or" Linked Panel
Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)	
[] HYDROcodone-acetaminophen (NORCO 10-325) 10-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient is able to tolerate oral medication.
[] HYDROcodone-acetaminophen (HYCET) 7.5-325 mg/15 mL solution	20 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) Use if patient can not swallow tablet.
() traMADol (ULTRAM) tablet - For eGFR LESS than 30 mL/min, change frequency to every 12 hours)	50 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op (Max Daily dose not to exceed 200 mg/day). Give if patient is able to tolerate oral medication

PRN Oral for Moderate Pain (Pain Score 4-6): For Patients GREATER than 65 years old (Single Response)
(adjust dose for renal/liver function and age)

() acetaminophen-codeine (TYLENOL #3) tablet OR elixir	"Or" Linked Panel
Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)	
[] acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient is able to tolerate oral medication.
[] acetaminophen-codeine 300 mg-30 mg /12.5 mL solution	12.5 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) Use if patient cannot swallow tablet.
() HYDROcodone-acetaminophen 5/325 (NORCO) tablet OR elixir	"Or" Linked Panel
Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)	
[] HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6)
[] HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution	10 mL, oral, every 6 hours PRN, moderate pain (score 4-6)

- | | |
|---|--|
| () traMADol (ULTRAM) tablet - For eGFR LESS than 30 mL/min, change frequency to every 12 hours | 25 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op
(Max Daily dose not to exceed 200 mg/day). Give if patient can tolerate oral medication. |
|---|--|

PRN IV for Moderate Pain (Pain Score 4-6): For Patients LESS than 65 years old (Single Response)

If you select a PCA option you will not be allowed to also order IV PRN pain medications from this section.
(adjust dose for renal/liver function and age)

- | | |
|---|--|
| () fentaNYL (SUBLIMAZE) injection | 25 mcg, intravenous, every 2 hour PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |
| () morphine 2 mg/mL injection | 2 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |
| () HYDROmorphine (DILAUDID) injection | 0.5 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |
| () ketorolac (TORADOL) IV (Single Response) | Do NOT use in patients with eGFR LESS than 30 mL/min AND/OR patients LESS than 17 years of age.
WARNING: Use is contraindicated for treatment of perioperative pain OR in the setting of coronary artery bypass graft (CABG) surgery. |
| () For patients ages GREATER than 64 OR weight LESS than 50 kg OR eGFR 30-59 mL/min - ketorolac (TORADOL) injection | 15 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6), For 5 Days |
| () For patients ages 17-64 AND weight GREATER than or EQUAL to 50 kg AND eGFR at least 60 mL/min - ketorolac (TORADOL) injection | 30 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6), For 5 Days |

PRN IV for Moderate Pain (Pain Score 4-6): For Patients GREATER than 65 years old (Single Response)

If you select a PCA option you will not be allowed to also order IV PRN pain medications from this section.
(adjust dose for renal/liver function and age)

- | | |
|---|---|
| () fentaNYL (SUBLIMAZE) injection | 12.5 mcg, intravenous, every 2 hour PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |
| () morphine 2 mg/mL injection | 1 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |
| () HYDROmorphine (DILAUDID) injection | 0.2 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |
| () ketorolac (TORADOL) injection - Do not use in patients with eGFR LESS than 30 mL/min. | 15 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6), For 5 Days, Post-op
Do not use in patients with eGFR LESS than 30 mL/min. Use if patient is unable to swallow or faster onset is needed. |

PRN Oral for Severe Pain (Pain Score 7-10): For Patients LESS than 65 years old (Single Response)

(adjust dose for renal/liver function and age)

- | | |
|---|--|
| () HYDROmorphine (DILAUDID) tablet | 2 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient is able to tolerate oral medication |
| () morphine (MSIR) tablet | 15 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient is able to tolerate oral medication |
| () oxyCODONE (ROXICODONE) immediate release tablet | 10 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient is able to tolerate oral medication |

PRN Oral for Severe Pain (Pain Score 7-10): For Patients GREATER than 65 years old (Single Response)

(adjust dose for renal/liver function and age)

<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 7.5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Give if patient is able to tolerate oral medication
<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO 10-325) 10-325 mg per tablet	1 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Give if patient is able to tolerate oral medication
<input type="checkbox"/> HYDROmorphine (DILAUDID) tablet	2 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Give if patient is able to tolerate oral medication
<input type="checkbox"/> morphine (MSIR) tablet	15 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Give if patient is able to tolerate oral medication
<input type="checkbox"/> oxyCODONE (ROXICODONE) immediate release tablet	5 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Give if patient is able to tolerate oral medication

PRN IV for Severe Pain (Pain Score 7-10): For Patients LESS than 65 years old (Single Response)

If you select a PCA option you will not be allowed to also order IV PRN pain medications from this section.
(adjust dose for renal/liver function and age)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	50 mcg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed
<input type="checkbox"/> morphine injection	4 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed
<input type="checkbox"/> HYDROmorphine (DILAUDID) injection	0.8 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed

PRN IV for Severe Pain (Pain Score 7-10): For Patients GREATER than 65 years old (Single Response)

If you select a PCA option you will not be allowed to also order IV PRN pain medications from this section.
(adjust dose for renal/liver function and age)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25 mcg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed
<input type="checkbox"/> morphine injection	2 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed
<input type="checkbox"/> HYDROmorphine (DILAUDID) injection	0.5 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed

Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.

<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.
--	---

Antiemetics - HMSL, HMWB Only

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9 % 0.9 % 20 mL for Alaris pump syringe option	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics - HMSTJ Only

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IVPB or Oral or Rectal	"Or" Linked Panel
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Insomnia: For Patients LESS than 70 years old (Single Response)

<input type="checkbox"/> zolpidem (AMBIEN) tablet	5 mg, oral, nightly PRN, sleep
<input type="checkbox"/> ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep

Insomnia: For Patients GREATER than or EQUAL to 70 years old (Single Response)

<input type="checkbox"/> ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep
---	--------------------------------

Itching: For Patients LESS than 70 years old (Single Response)

<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet	25 mg, oral, every 6 hours PRN, itching
<input type="checkbox"/> hydrOXYzine (ATARAX) tablet	10 mg, oral, every 6 hours PRN, itching
<input type="checkbox"/> cetirizine (ZyrTEC) tablet	5 mg, oral, daily PRN, itching

fexofenadine (ALLEGRA) tablet - For eGFR LESS than 80 mL/min, reduce frequency to once daily as needed 60 mg, oral, 2 times daily PRN, itching

Itching: For Patients between 70-76 years old (Single Response)

cetirizine (ZyrTEC) tablet 5 mg, oral, daily PRN, itching

Itching: For Patients GREATER than 77 years old (Single Response)

cetirizine (ZyrTEC) tablet 5 mg, oral, daily PRN, itching

GI Medications (Single Response)

	"Or" Linked Panel
<input type="checkbox"/> famotidine (PEPCID) Oral or IV	
<input type="checkbox"/> famotidine (PEPCID) tablet	20 mg, oral, 2 times daily, Post-op May crush and give per nasogastric tube if needed. Give the tablet if the patient can tolerate oral medication.
<input type="checkbox"/> famotidine (PEPCID) injection	20 mg, intravenous, 2 times daily, Post-op Use injection if patient cannot tolerate oral medication or requires a faster onset of action.

VTE

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.

Will encourage early ambulation

PACU & Post-op

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.

Therapy for the following:

PACU & Post-op

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	

<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee)	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op

<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1, PACU & Post-op Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1, PACU & Post-op To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.

Will encourage early ambulation

PACU & Post-op

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.

Therapy for the following:

PACU & Post-op

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCl LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCl LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Non-Surgical Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1, PACU & Post-op Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1, PACU & Post-op To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op

Labs Today

Hematology/Coagulation

<input type="checkbox"/> Hemoglobin and hematocrit	Once, Post-op
<input type="checkbox"/> CBC with platelet and differential	Once, Post-op
<input type="checkbox"/> Prothrombin time with INR	Once, Post-op
<input type="checkbox"/> Partial thromboplastin time	Once, Post-op
<input type="checkbox"/> Type and screen	Once, Post-op

Chemistry

<input type="checkbox"/> Basic metabolic panel	Once, Post-op
<input type="checkbox"/> Comprehensive metabolic panel	Once, Post-op
<input type="checkbox"/> Calcium	Once, Post-op
<input type="checkbox"/> Hepatic function panel	Once, Post-op
<input type="checkbox"/> Magnesium	Once, Post-op
<input type="checkbox"/> Phosphorus level	Once, Post-op

Labs Tomorrow

Hematology/Coagulation

<input type="checkbox"/>	CBC with platelet and differential	Once, Starting S+1, Post-op
<input type="checkbox"/>	Prothrombin time with INR	Once, Starting S+1, Post-op
<input type="checkbox"/>	Partial thromboplastin time	Once, Starting S+1, Post-op
<input type="checkbox"/>	Type and screen	Once, Starting S+1, Post-op

Chemistry

<input type="checkbox"/>	Amylase	Once, Starting S+1, Post-op
<input type="checkbox"/>	Basic metabolic panel	Once, Starting S+1, Post-op
<input type="checkbox"/>	Comprehensive metabolic panel	Once, Starting S+1, Post-op
<input type="checkbox"/>	Calcium	Once, Starting S+1, Post-op
<input type="checkbox"/>	Hepatic function panel	Once, Starting S+1, Post-op
<input type="checkbox"/>	Lipase level	Once, Starting S+1, Post-op
<input type="checkbox"/>	Magnesium	Once, Starting S+1, Post-op
<input type="checkbox"/>	Phosphorus level	Once, Starting S+1, Post-op

Cardiology

Imaging

X-Ray

<input type="checkbox"/>	Chest 1 Vw Portable	Routine, 1 time imaging For 1 , Post-op
<input type="checkbox"/>	Abdomen 1 Vw Portable	Routine, 1 time imaging For 1 , Post-op

Other Studies

Respiratory

Respiratory

<input checked="" type="checkbox"/>	Incentive spirometry	Routine, Every hour Every 1 hour while awake. Instruct on use: 10 repetitions, Post-op
<input checked="" type="checkbox"/>	Encourage deep breathing and coughing	Routine, Every hour Waking hours only , Post-op
<input checked="" type="checkbox"/>	Oxygen therapy	Routine, Continuous Device 1: Nasal Cannula Rate in liters per minute: 2 Lpm Rate in tenths of a liter per minute: O2 %: Device 2: Device 3: Titrate to keep O2 Sat Above: 92% Indications for O2 therapy: Post-op

Rehab

Consults

For Physician Consult orders use sidebar

Ancillary Consults

<input type="checkbox"/>	Consult to Case Management	Consult Reason: Post-op
<input type="checkbox"/>	Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/>	Consult PT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/>	Consult PT wound care	Special Instructions: Location of Wound? Post-op

<input type="checkbox"/> Consult OT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Purpose/Topic: Post-op
<input type="checkbox"/> Consult to Spiritual Care	Reason for consult? Post-op
<input type="checkbox"/> Consult to Speech Language Pathology	Routine, Once Reason for consult: Post-op
<input type="checkbox"/> Consult to Wound Ostomy Care nurse	Reason for consult: Reason for consult: Reason for consult: Reason for consult: Consult for NPWT: Reason for consult: Post-op
<input type="checkbox"/> Consult to Respiratory Therapy	Reason for Consult? Post-op

Additional Orders