

General

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Details
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Details
<input type="checkbox"/>	Acute Renal Failure	Details
<input type="checkbox"/>	Acute Respiratory Failure	Details
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Details
<input type="checkbox"/>	Anemia	Details
<input type="checkbox"/>	Bacteremia	Details
<input type="checkbox"/>	Bipolar disorder, unspecified	Details
<input type="checkbox"/>	Cardiac Arrest	Details
<input type="checkbox"/>	Cardiac Dysrhythmia	Details
<input type="checkbox"/>	Cardiogenic Shock	Details
<input type="checkbox"/>	Decubitus Ulcer	Details
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Details
<input type="checkbox"/>	Disorder of Liver	Details
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Details
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Details
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Details
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Details
<input type="checkbox"/>	Other Alteration of Consciousness	Details
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Details
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Details
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Details
<input type="checkbox"/>	Protein-calorie Malnutrition	Details
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Details
<input type="checkbox"/>	Schizophrenia Disorder	Details
<input type="checkbox"/>	Sepsis	Details
<input type="checkbox"/>	Septic Shock	Details
<input type="checkbox"/>	Septicemia	Details
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Details
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Details

Admission or Observation (Single Response)

<input type="checkbox"/> Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
<input type="checkbox"/> Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
<input type="checkbox"/> Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

Admission or Observation (Single Response)

Patient has active status order on file

<input type="checkbox"/> Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
---	--

<input type="checkbox"/> Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
--	---

<input type="checkbox"/> Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:
--	---

Admission (Single Response)

Patient has active status order on file.

<input type="checkbox"/> Admit to inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
---	--

Code Status

<input type="checkbox"/> Full code	Code Status decision reached by:
------------------------------------	----------------------------------

<input type="checkbox"/> DNR	Does patient have decision-making capacity?
------------------------------	---

<input type="checkbox"/> DNR (Do Not Resuscitate)	Priority:
---	-----------

<input type="checkbox"/> Consult to Palliative Care Service	Reason for Consult?
---	---------------------

	Order?
--	--------

	Name of referring provider:
--	-----------------------------

	Enter call back number:
--	-------------------------

<input type="checkbox"/> Consult to Social Work	Reason for Consult:
---	---------------------

<input type="checkbox"/> Modified Code	Does patient have decision-making capacity?
--	---

	Modified Code restrictions:
--	-----------------------------

<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by:
---	--

	Specify Treatment Restrictions:
--	---------------------------------

Isolation

<input type="checkbox"/> Airborne isolation status	Details
--	---------

<input type="checkbox"/> Contact isolation status	Details
---	---------

<input type="checkbox"/> Droplet isolation status	Details
---	---------

<input type="checkbox"/> Enteric isolation status	Details
---	---------

Precautions

<input type="checkbox"/> Aspiration precautions	Details
---	---------

<input type="checkbox"/> Fall precautions	Increased observation level needed:
---	-------------------------------------

<input type="checkbox"/> Latex precautions	Details
--	---------

<input type="checkbox"/> Seizure precautions	Increased observation level needed:
--	-------------------------------------

Nursing

Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP every 4 hours	Routine, Every 4 hours
---	------------------------

<input type="checkbox"/> Vital signs - T/P/R/BP per unit protocol	Routine, Per unit protocol
Activity	
<input type="checkbox"/> Ambulate	Routine, 3 times daily Specify:
<input type="checkbox"/> Ambulate as tolerated	Routine, 3 times daily Specify:
<input type="checkbox"/> Strict bed rest	Routine, Until discontinued, Starting S May elevate Head of Bed 30 degrees.
Nursing	
<input type="checkbox"/> Telemetry	"And" Linked Panel
<input type="checkbox"/> Telemetry monitoring	Routine, Continuous Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box) Reason for telemetry: Can be off of Telemetry for tests and baths? Yes
<input type="checkbox"/> Telemetry Additional Setup Information	Routine, Continuous High Heart Rate (BPM): 120 Low Heart Rate(BPM): 50 High PVC's (per minute): 10 High SBP(mmHg): 175 Low SBP(mmHg): 100 High DBP(mmHg): 95 Low DBP(mmHg): 40 Low Mean BP: 60 High Mean BP: 120 Low SPO2(%): 94
<input type="checkbox"/> Place TED hose	Routine, Once
<input type="checkbox"/> Collect initial labs before starting anticoagulation.	Routine, Once For 1 Occurrences
<input type="checkbox"/> Height and weight	Routine, Once For 1 Occurrences On Admission.
Notify	
<input type="checkbox"/> Notify Physician of following vitals	Routine, Until discontinued, Starting S, Systolic BP GREATER than 180 mmHg Systolic BP LESS than 80 mmHg Heart rate GREATER than 120 bpm Heart rate LESS than 55 bpm SpO2 LESS than 90
Diet	
<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options:
<input type="checkbox"/> Diet - Regular	Diet effective now, Starting S Diet(s): Regular,Other Potass/Phos Potassium/Phosphorus: 2 GM Potassium Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Low Vitamin K
<input type="checkbox"/> Diet - 2000 Kcal/225 gm Carb	Diet effective now, Starting S Diet(s): 2000 Kcal/225 gm Carbohydrate Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:

<input type="checkbox"/> Diet - Heart healthy	Diet effective now, Starting S Diet(s): Heart Healthy Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
<input type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:

IV Fluids

IV Fluids

<input type="checkbox"/> sodium chloride 0.45 % infusion	intravenous, continuous
<input type="checkbox"/> sodium chloride 0.9 % infusion	intravenous, continuous

Medications

Low Molecular Weight Heparins

<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> CrCl greater than or equal to 30 mL/min	1 mg/kg, subcutaneous, 2 times daily at 0600, 1800 (time critical)
<input type="checkbox"/> CrCl less than 30 mL/min	1 mg/kg, subcutaneous, daily at 1700 (time critical)

Vitamin K Antagonists (Single Response)

<input type="checkbox"/> warfarin (COUMADIN) with consult and labs	
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, once, S at 5:00 PM, For 1 Doses Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	Routine, Until discontinued, Starting S Indication:
<input type="checkbox"/> Prothrombin Time/INR STAT	STAT, Starting S For 1 Occurrences
<input type="checkbox"/> Prothrombin Time/INR AM Draw	AM draw, Starting S+1 For 1 Occurrences
<input type="checkbox"/> Prothrombin Time/INR every 72 hours	Every 72 hours, Starting S+1
<input type="checkbox"/> warfarin (COUMADIN) with consult and labs	
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, once, S at 5:00 PM, For 1 Doses Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	Routine, Until discontinued, Starting S Indication:
<input type="checkbox"/> Prothrombin Time/INR STAT	STAT, Starting S For 1 Occurrences
<input type="checkbox"/> Prothrombin Time/INR every 72 hours	Every 72 hours, Starting S+1
<input type="checkbox"/> warfarin (COUMADIN) with labs	
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Prothrombin Time/INR STAT	STAT, Starting S For 1 Occurrences
<input type="checkbox"/> Prothrombin Time/INR AM Draw	AM draw, Starting S+1 For 1 Occurrences
<input type="checkbox"/> Prothrombin Time/INR every 72 hours	Every 72 hours, Starting S+1
<input type="checkbox"/> warfarin (COUMADIN) with labs	
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Prothrombin Time/INR STAT	STAT, Starting S For 1 Occurrences
<input type="checkbox"/> Prothrombin Time/INR every 7 days	Weekly, Starting S+1

Heparin (Single Response)

<input type="checkbox"/> Pharmacy to dose heparin	Routine, Until discontinued, Starting S Heparin Indication: Specify: Monitoring: Anti-Xa
---	---

heparin infusion 50 units/mL in dextrose 5% intravenous, continuous
 Indication:
 Therapeutic Monitoring Target:

Direct Xa Inhibitors (Single Response)

<input type="checkbox"/> rivaroxaban (XARELTO) initial therapy and maintenance	"And" Linked Panel
<input type="checkbox"/> rivaroxaban (XARELTO) tablet	15 mg, oral, 2 times daily at 0900, 1700 (time critical), For 21 Days Indications: DVT/PE
<input type="checkbox"/> Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> rivaroxaban (XARELTO) tablet	20 mg, oral, daily at 1700 (time critical), Starting H+504 Hours Indications: DVT/PE
<input type="checkbox"/> apixaban (ELIQUIS) initial therapy and maintenance	"And" Linked Panel
<input type="checkbox"/> apixaban (ELIQUIS) tablet	10 mg, oral, 2 times daily, For 7 Days Indications: DVT/PE
<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S
<input type="checkbox"/> apixaban (ELIQUIS) tablet	5 mg, oral, 2 times daily, Starting S+7 Indications: DVT/PE
<input type="checkbox"/> dabigatran (PRADAXA) therapy - after 5 days of parenteral anticoagulation	"And" Linked Panel
<input type="checkbox"/> dabigatran etexilate (PRADAXA) capsule	150 mg, oral, 2 times daily Indications: DVT/PE
<input type="checkbox"/> Pharmacy consult to monitor dabigatran (PRADAXA) therapy	STAT, Until discontinued, Starting S

Analgesics

<input type="checkbox"/> acetaminophen (TYLENOL) oral/rectal	"Or" Linked Panel
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	325 mg, oral, every 4 hours PRN, fever Maximum dose of 3,000 mg per day (total from all sources).
<input type="checkbox"/> acetaminophen (TYLENOL) suppository	325 mg, rectal, every 4 hours PRN, fever Maximum dose of 3,000 mg per day (total from all sources). For rectal use only.

VTE

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition
 Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition
 Both pharmacologic AND mechanical prophylaxis must be addressed.
 Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:
 Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)
 Age 60 and above Severe fracture of hip, pelvis or leg
 Central line Acute spinal cord injury with paresis
 History of DVT or family history of VTE Multiple major traumas
 Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER
 Less than fully and independently ambulatory Acute ischemic stroke
 Estrogen therapy History of PE
 Moderate or major surgery (not for cancer)
 Major surgery within 3 months of admission

Low Risk of DVT
 Low Risk (Single Response)

<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
<input type="checkbox"/> Moderate Risk of DVT - Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	

<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee)	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

fondaparinux (ARIXTRA) injection

2.5 mg, subcutaneous, daily, Starting S+1

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

Moderate Risk of DVT - Non-Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE Routine, Once

Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)

<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

High Risk of DVT - Surgical

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

High Risk

High risk of VTE Routine, Once

High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

High Risk of DVT - Non-Surgical

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

High Risk

High risk of VTE Routine, Once

High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)

<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily, Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical) For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.

Will encourage early ambulation

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once

Labs

Labs STAT

<input type="checkbox"/> CBC and differential	STAT For 1 Occurrences
<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences
<input type="checkbox"/> POC occult blood stool and Notify	"And" Linked Panel
<input type="checkbox"/> POC occult blood stool	Once
<input type="checkbox"/> Notify Physician If occult stool is positive.	Routine, Until discontinued, Starting S
<input type="checkbox"/> Antithrombin III	STAT For 1 Occurrences
<input type="checkbox"/> Cardiolipin antibody	STAT For 1 Occurrences
<input type="checkbox"/> Factor V assay	STAT For 1 Occurrences
<input type="checkbox"/> Factor V leiden	STAT For 1 Occurrences
<input type="checkbox"/> Homocystine, plasma	STAT For 1 Occurrences
<input type="checkbox"/> Lupus anticoagulant	STAT For 1 Occurrences
<input type="checkbox"/> Protein C activity	STAT For 1 Occurrences
<input type="checkbox"/> Protein S activity	STAT For 1 Occurrences
<input type="checkbox"/> Prothrombin gene mutation	STAT For 1 Occurrences

Labs Tomorrow

<input type="checkbox"/> CBC and differential	AM draw For 1 Occurrences
<input type="checkbox"/> Partial thromboplastin time	AM draw For 1 Occurrences
<input type="checkbox"/> Comprehensive metabolic panel	AM draw For 1 Occurrences

Labs AM Repeat

<input type="checkbox"/> CBC and differential	AM draw repeats For 3 Occurrences
<input type="checkbox"/> Partial thromboplastin time	AM draw repeats For 3 Occurrences
<input type="checkbox"/> Prothrombin time with INR	AM draw repeats For 3 Occurrences

Cardiology

Imaging

CT

<input type="checkbox"/> CT Angiogram Pe Chest	Routine, 1 time imaging For 1 PE Protocol
--	--

X-Ray

<input type="checkbox"/> Chest 1 Vw Portable	Routine, 1 time imaging For 1 Occurrences
<input type="checkbox"/> Chest 2 Vw	Routine, 1 time imaging For 1 Occurrences

US for DVT Diagnosis

<input type="checkbox"/> PV duplex venous lower extremity bilat	Routine, 1 time imaging
<input type="checkbox"/> PV duplex venous lower extremity left	Routine, 1 time imaging
<input type="checkbox"/> PV duplex venous lower extremity right	Routine, 1 time imaging
<input type="checkbox"/> PV duplex venous upper extremity bilat	Routine, 1 time imaging
<input type="checkbox"/> PV duplex venous upper extremity left	Routine, 1 time imaging
<input type="checkbox"/> PV duplex venous upper extremity right	Routine, 1 time imaging

US

<input type="checkbox"/> USPV Venous Lower Extremity Bilateral	Routine, 1 time imaging For 1
<input type="checkbox"/> USPV Venous Lower Extremity Left	Routine, 1 time imaging For 1
<input type="checkbox"/> USPV Venous Lower Extremity Right	Routine, 1 time imaging For 1
<input type="checkbox"/> USPV Venous Upper Extremity Bilat	Routine, 1 time imaging For 1
<input type="checkbox"/> USPV Venous Upper Extremity Left	Routine, 1 time imaging For 1
<input type="checkbox"/> USPV Venous Upper Extremity Right	Routine, 1 time imaging For 1

Nuclear

<input type="checkbox"/> NM Lung Ventilation Perfusion	Routine, 1 time imaging For 1
--	-------------------------------

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Ancillary Consults

<input type="checkbox"/> Consult to Case Management	Consult Reason:
<input type="checkbox"/> Consult to Social Work	Reason for Consult:
<input type="checkbox"/> Consult to PT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult to OT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Other (Specify) Specify: Low vitamin K diet
<input type="checkbox"/> Consult to Spiritual Care	Reason for consult?
<input type="checkbox"/> Consult to Respiratory Therapy	Reason for Consult?

Additional Orders