

General

Elective Outpatient, Observation (Single Response)

() Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
() Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
() Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments: PACU & Post-op

Nursing

Vital Signs

[X] Vital signs (per unit protocol)	Routine, Per unit protocol, Post-op
-------------------------------------	-------------------------------------

Activity

[] Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated Post-op
[] Ambulate with assistance	Routine, 3 times daily Specify: with assistance Post-op
[] Bed rest with bathroom privileges	Routine, Until discontinued, Starting S Bathroom Privileges: Post-op
[] Strict bed rest	Routine, Until discontinued, Starting S May resume normal activity 24 hours after PEG placement., Post-op

Nursing

[X] Observe minimum of 1-2 hours post procedure	Routine, Until discontinued, Starting S, Post-op
[X] Return to floor/unit when stable per Endoscopy protocol	Routine, Until discontinued, Starting S, Post-op
[X] Resume active orders on return to floor/unit	Routine, Until discontinued, Starting S, Post-op
[] Positioning instruction	Routine, Until discontinued, Starting S Position: Left Lateral Recumbent Post-op
[] Have patient lie on side until fully awake	Routine, Until discontinued, Starting S, Post-op
[] Discontinue IV prior to discharge	Routine, Until discontinued, Starting S, Post-op
[] Discharge when meets discharge criteria	Routine, Until discontinued, Starting S, Post-op

PEG Orders

[] Positioning instruction	Routine, Until discontinued, Starting S For 24 Hours Position: Reverse Trendelenberg Post-op
[] Tube site care	Routine, Every morning Clean PEG site and change PEG dressings with hydrogen peroxide and povidone iodine., Post-op
[] Check residual	Routine, Per unit protocol Check gastric residual volume every 8 hours and hold tube feedings for 2 hours if more than 400 ml. When volume returns below 400 ml, resume tube feeding at previous rate., Post-op

[] Abdominal binder	Routine, Once Waking hours only? Nurse to schedule? Special Instructions: Post-op
[] OK to use PEG for meds now. Ok to start feeds via PEG in 3 hours.	Routine, Until discontinued, Starting S, Post-op
[] Keep patients hands away from PEG and prevent patient from pulling PEG out	Routine, Until discontinued, Starting S, Post-op

Notify

[] Notify Physician for signs of bleeding, abdominal or chest pain, or fever greater than 101.5	Routine, Until discontinued, Starting S, signs of bleeding, abdominal or chest pain, or fever greater than 101.5, Post-op
--	---

Diet

[] NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options: Post-op
[] NPO	Diet effective _____ NPO: Pre-Operative fasting options: Post-op
[] Diet	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op

IV Fluids

Medications

VTE

Labs

Cardiology

Imaging

Other Studies

Respiratory

Rehab

Consults

Additional Orders