

General

Common Present on Admission Diagnosis

<input type="checkbox"/> Acidosis	Details
<input type="checkbox"/> Acute Post-Hemorrhagic Anemia	Details
<input type="checkbox"/> Acute Renal Failure	Details
<input type="checkbox"/> Acute Respiratory Failure	Details
<input type="checkbox"/> Acute Thromboembolism of Deep Veins of Lower Extremities	Details
<input type="checkbox"/> Anemia	Details
<input type="checkbox"/> Bacteremia	Details
<input type="checkbox"/> Bipolar disorder, unspecified	Details
<input type="checkbox"/> Cardiac Arrest	Details
<input type="checkbox"/> Cardiac Dysrhythmia	Details
<input type="checkbox"/> Cardiogenic Shock	Details
<input type="checkbox"/> Decubitus Ulcer	Details
<input type="checkbox"/> Dementia in Conditions Classified Elsewhere	Details
<input type="checkbox"/> Disorder of Liver	Details
<input type="checkbox"/> Electrolyte and Fluid Disorder	Details
<input type="checkbox"/> Intestinal Infection due to Clostridium Difficile	Details
<input type="checkbox"/> Methicillin Resistant Staphylococcus Aureus Infection	Details
<input type="checkbox"/> Obstructive Chronic Bronchitis with Exacerbation	Details
<input type="checkbox"/> Other Alteration of Consciousness	Details
<input type="checkbox"/> Other and Unspecified Coagulation Defects	Details
<input type="checkbox"/> Other Pulmonary Embolism and Infarction	Details
<input type="checkbox"/> Phlebitis and Thrombophlebitis	Details
<input type="checkbox"/> Protein-calorie Malnutrition	Details
<input type="checkbox"/> Psychosis, unspecified psychosis type	Details
<input type="checkbox"/> Schizophrenia Disorder	Details
<input type="checkbox"/> Sepsis	Details
<input type="checkbox"/> Septic Shock	Details
<input type="checkbox"/> Septicemia	Details
<input type="checkbox"/> Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Details
<input type="checkbox"/> Urinary Tract Infection, Site Not Specified	Details

Admission or Observation (Single Response)

<input type="checkbox"/> Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
<input type="checkbox"/> Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
<input type="checkbox"/> Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

Admission or Observation (Single Response)

Patient has active status order on file

<input type="checkbox"/> Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
---	--

<input type="checkbox"/> Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
--	---

<input type="checkbox"/> Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:
--	---

Admission (Single Response)

Patient has active status order on file.

<input type="checkbox"/> Admit to inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
---	--

Code Status

<input type="checkbox"/> Full code	Code Status decision reached by:
------------------------------------	----------------------------------

<input type="checkbox"/> DNR	Does patient have decision-making capacity?
------------------------------	---

<input type="checkbox"/> DNR (Do Not Resuscitate)	Priority:
---	-----------

<input type="checkbox"/> Consult to Palliative Care Service	Reason for Consult?
---	---------------------

	Order?
--	--------

	Name of referring provider:
--	-----------------------------

	Enter call back number:
--	-------------------------

<input type="checkbox"/> Consult to Social Work	Reason for Consult:
---	---------------------

<input type="checkbox"/> Modified Code	Does patient have decision-making capacity?
--	---

	Modified Code restrictions:
--	-----------------------------

<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by:
---	--

	Specify Treatment Restrictions:
--	---------------------------------

Isolation

<input type="checkbox"/> Airborne isolation status	Details
--	---------

<input type="checkbox"/> Contact isolation status	Details
---	---------

<input type="checkbox"/> Droplet isolation status	Details
---	---------

<input type="checkbox"/> Enteric isolation status	Details
---	---------

Precautions

<input type="checkbox"/> Aspiration precautions	Details
---	---------

<input type="checkbox"/> Fall precautions	Increased observation level needed:
---	-------------------------------------

<input type="checkbox"/> Latex precautions	Details
--	---------

<input type="checkbox"/> Seizure precautions	Increased observation level needed:
--	-------------------------------------

Nursing

Vital Signs

<input type="checkbox"/> Vital Signs every 4 Hrs	Routine, Every 4 hours
--	------------------------

Vital signs every 8 Hrs Routine, Every 8 hours

Activity

Bed rest with bathroom privileges Routine, Until discontinued, Starting S
Bathroom Privileges: with bathroom privileges

Out of bed, Up with assistance Routine, Until discontinued, Starting S
Specify: Out of bed, Up with assistance

Nursing

Telemetry **"And" Linked Panel**

Telemetry monitoring Routine, Continuous
Order: Place in Centralized Telemetry Monitor: EKG
Monitoring Only (Telemetry Box)
Reason for telemetry:
Can be off of Telemetry for tests and baths? Yes

Telemetry Additional Setup Information Routine, Continuous
High Heart Rate (BPM): 120
Low Heart Rate(BPM): 50
High PVC's (per minute): 10
High SBP(mmHg): 175
Low SBP(mmHg): 100
High DBP(mmHg): 95
Low DBP(mmHg): 40
Low Mean BP: 60
High Mean BP: 120
Low SPO2(%): 94

Neurological assessment every 4 Hrs Routine, Every 4 hours
Assessment to Perform:

Toileting - bedside commode Routine, Until discontinued, Starting S
Specify: Bedside commode
Additional modifier:

Request for Central Supply Equipment - Walker Routine, Once
Equipment Requested: Regular Walker With Wheels
Special Instructions:

Height and weight on admission Routine, Once For 1 Occurrences
On admission.

OK to draw all ordered labs during plasma exchange Routine, Until discontinued, Starting S
If patient is receiving plasma exchange therapy, ok to draw all
ordered labs during plasma exchange process.

Notify

Notify Physician: If patient has: Shortness of Breath,
Difficulty with secretion, Forced Vital Capacity less than
15 milliliters per kilogram, Negative Inspiratory Force
less than -25 centimeters H2O Routine, Until discontinued, Starting S, If patient has:
Shortness of Breath, Difficulty with secretion, Forced Vital
Capacity less than 15 milliliters per kilogram, Negative
Inspiratory Force less than -25 centimeters H2O

Notify Physician for HR greater than 110 bpm Routine, Until discontinued, Starting S
Temperature greater than:
Temperature less than:
Systolic BP greater than:
Systolic BP less than:
Diastolic BP greater than:
Diastolic BP less than:
MAP less than:
Heart rate greater than (BPM): 110
Heart rate less than (BPM):
Respiratory rate greater than:
Respiratory rate less than:
SpO2 less than:

Diet

<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options:
<input type="checkbox"/> Diet - Regular	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Additional Instructions: Foods to Avoid:
<input type="checkbox"/> Diet - Heart healthy	Diet effective now, Starting S Diet(s): Heart Healthy Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:

IV Fluids

Medications

Medications

<input type="checkbox"/> pyridostigmine (MESTINON) CR tablet	180 mg, oral, nightly
<input type="checkbox"/> pyridostigmine (MESTINON) tablet	60 mg, oral, every 8 hours scheduled

Immunosuppressants

<input type="checkbox"/> azaTHIOprine (IMURAN) tablet	oral, every 12 hours scheduled
<input type="checkbox"/> cycloSPORINE modified (NEORAL) capsule	oral, 2 times daily at 0600, 1800 (time critical)
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) IV	intravenous
<input type="checkbox"/> mycophenolate (CELLCEPT) tablet	oral, 2 times daily at 0600, 1800 (time critical)
<input type="checkbox"/> predniSONE (DELTASONE) tablet	oral, daily

Insomnia (Single Response)

<input type="checkbox"/> ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep
<input type="checkbox"/> temazepam (RESTORIL) capsule	oral, nightly PRN, sleep
<input type="checkbox"/> traZODone (DESYREL) tablet	50 mg, oral, nightly PRN, sleep Indication:
<input type="checkbox"/> zolpidem (AMBIEN) tablet	5 mg, oral, nightly PRN, sleep

Medications - IVIG (Single Response)

<input type="checkbox"/> IVIG (Immune Globulin 10%) 400 mg/kg x5 doses with common predmedication orders	"And" Linked Panel
<input type="checkbox"/> diphenhydrAMINE (SOMINEX) tablet	25 mg, oral, every 24 hours, For 5 Doses Administer prior to IVIG administration.
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, every 24 hours, For 5 Doses Administer prior to IVIG administration.
<input type="checkbox"/> Immune Globulin (IGG) 10% IV	0.4 g/kg, intravenous, every 24 hours, Starting H+30 Minutes, For 5 Doses
<input type="checkbox"/> IVIG (Immune Globulin 10%) 500 mg/kg x4 doses with common predmedication orders	"And" Linked Panel
<input type="checkbox"/> diphenhydrAMINE (SOMINEX) tablet	25 mg, oral, every 24 hours, For 4 Doses Administer prior to IVIG administration.
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, every 24 hours, For 4 Doses Administer prior to IVIG administration.
<input type="checkbox"/> Immune Globulin (IGG) 10% IV	0.5 g/kg, intravenous, every 24 hours, Starting H+30 Minutes, For 4 Doses

Insomnia: For Patients GREATER than or EQUAL to 70 years old (Single Response)

<input type="checkbox"/> ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep
---	--------------------------------

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCl LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

<input type="checkbox"/> Low Risk of DVT	
<input type="checkbox"/> Low Risk (Single Response)	
<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
<input type="checkbox"/> Moderate Risk of DVT - Surgical	

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

Moderate Risk of DVT - Non-Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	

<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

High Risk of DVT - Surgical

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

High Risk

High risk of VTE

Routine, Once

High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
For Patients with CrCL LESS than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> High Risk of DVT - Non-Surgical Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily, Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical) For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee)	

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min

fondaparinux (ARIXTRA) injection

2.5 mg, subcutaneous, daily, Starting S+1

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:

<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Non-Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee)	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once

Labs

Labs Today

<input type="checkbox"/> CBC and differential	Once
<input type="checkbox"/> Basic metabolic panel	Once
<input type="checkbox"/> Comprehensive metabolic panel	Once
<input type="checkbox"/> Sedimentation rate	Once
<input type="checkbox"/> ANA	Once
<input type="checkbox"/> Hemoglobin A1c	Once
<input checked="" type="checkbox"/> Immunoglobulin A	Once
<input type="checkbox"/> T uptake	Once
<input type="checkbox"/> T3	Once
<input type="checkbox"/> T3, free	Once
<input type="checkbox"/> T4	Once
<input type="checkbox"/> T4, free	Once
<input type="checkbox"/> TSH	Once
<input type="checkbox"/> Acetylcholine receptor, binding	Once
<input type="checkbox"/> Thyroxine binding globulin	Once
<input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once
	Specimen Source: Urine
	Specimen Site:

Cardiology

Cardiology

<input type="checkbox"/> ECG 12 lead	Routine, Once Clinical Indications: Shortness of Breath Interpreting Physician:
--------------------------------------	---

Imaging

CT

<input type="checkbox"/> CT Chest W Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/> CT Chest Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/> CT Chest W Wo Contrast	Routine, 1 time imaging For 1

X-Ray

<input type="checkbox"/> Chest 2 Vw	Routine, 1 time imaging For 1 Occurrences
-------------------------------------	---

Other Studies

Respiratory

Respiratory Therapy

<input type="checkbox"/> Oxygen therapy - Nasal cannula	Routine, Continuous Device 1: Nasal Cannula Rate in liters per minute: Rate in tenths of a liter per minute: O2 %: Titrate to keep O2 Sat Above: 92% Indications for O2 therapy: Device 2: Device 3:
---	--

<input type="checkbox"/> Vital capacity	Routine, Every 4 hours PRN (RT) Forced Vital Capacity. Record in chart. If poor oral seal please use face mask.
<input type="checkbox"/> Negative inspiratory force	Routine, Every 4 hours PRN (RT) Record in chart.
<input type="checkbox"/> Mechanical ventilation - NPPV	Routine Mechanical Ventilation: Non-Invasive Inspiratory Pressure (cm H2O): PEEP (cm H2O): % O2 (%): Pressure Support (cm H2O): VT - Tidal Volume (mL): Vent Management Strategies: Non-Invasive Positive Pressure Ventilation Empiric Trial for {Condition:21144}.

Rehab

Consults

For Physician Consult orders use sidebar

Consults

<input type="checkbox"/> Consult to Case Management	Consult Reason:
<input type="checkbox"/> Consult to Social Work	Reason for Consult:
<input type="checkbox"/> Consult PT Eval and Treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult PT wound care	Special Instructions: Location of Wound?
<input type="checkbox"/> Consult OT Eval and Teat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Purpose/Topic:
<input type="checkbox"/> Consult to Spiritual Care	Reason for consult?
<input type="checkbox"/> Consult to Speech Language Pathology for bedside swallow exam	Routine, Once Reason for SLP? Bedside swallow exam
<input type="checkbox"/> Consult to Wound Ostomy Care nurse	Reason for consult: Reason for consult: Reason for consult: Reason for consult: Consult for NPWT: Reason for consult:
<input type="checkbox"/> Consult to Respiratory Therapy	Reason for Consult?

Additional Orders