

# BODYIR Pre Procedure [1376]

## General

## Nursing

### Consent

<input type="checkbox"/> Complete consent for	Routine, Once Procedure: Diagnosis/Condition: Physician: Pre-Procedure
---	--

### Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Once For 1 Occurrences Upon arrival to Interventional Radiology unit., Pre-Procedure
<input type="checkbox"/> Pulse oximetry	Routine, Once For 1 Occurrences Current FIO2 or Room Air: Upon arrival to Interventional Radiology unit., Pre-Procedure

### Nursing

<input type="checkbox"/> Hold AM meds	Routine, Once For 1 Occurrences, Pre-Procedure
<input type="checkbox"/> Insert peripheral IV	Routine, Once Use 20-22 gauge Jelco catheter, Pre-Procedure
<input type="checkbox"/> Insert and maintain Foley	
<input type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed: Pre-Procedure
<input type="checkbox"/> Foley Catheter Care	Routine, Until discontinued, Starting S Orders: Maintain Pre-Procedure
<input type="checkbox"/> Bedside glucose	Routine, Once If diabetic, upon arrival to Endovascular/AOD unit. Notify Radiologist if glucose level less than 60 or greater than 200., Pre-Procedure

### Diet

<input type="checkbox"/> NPO after midnight	Diet effective midnight, Starting S+1 at 12:01 AM NPO: Except meds Pre-Operative fasting options: Pre-Procedure
---	--

## Medications

### Buffered Lidocaine 1% Injection

<input type="checkbox"/> lidocaine 1% buffered with 8.4% sodium bicarbonate injection	10 mL, injection, once, For 1 Doses, Pre-Procedure
---	--

### Prophylaxis Recommendation for Patients with an Allergy to Iodine Contrast Media (Single Response)

Medications may be administered by oral or IV route. The equivalent of these medications may be administered as determined by the attending or referring physician.

In a true medical emergency situation the exam should proceed with an accelerated premedication schedule at the direction of the attending physician. (attachment to Radiology Policy R 711)

( ) Oral methylPREDNISolone (MEDROL) tablet	32 mg, oral, once, For 1 Doses, Pre-Procedure For iodine allergy prophylaxis. Give 6-12 hours prior to the examination.
( ) IV methylPREDNISolone succinate (Solu-MEDROL) 40 mg/mL injection	40 mg, intravenous, once, For 1 Doses, Pre-Procedure For iodine allergy prophylaxis. Give 2 hours prior to the examination.

#### For patients with allergic reaction

[ ] diphenhydrAMINE (BENADRYL) injection	50 mg, intravenous, once, For 1 Doses, Pre-Procedure Given in Interventional Radiology unit
[ ] methylPREDNISolone sodium succinate (Solu-MEDROL) injection	125 mg, intravenous, once, For 1 Doses, Pre-Procedure Given in Interventional Radiology unit

#### VTE

#### Labs

##### Labs

[ ] CBC and differential	Once, Starting S, Pre-Procedure
[ ] Comprehensive metabolic panel	Once, Pre-Procedure
[ ] Basic metabolic panel	Once, Starting S, Pre-Procedure
[ ] Protimes-INR	Once, Starting S, Pre-Procedure
[ ] Platelet count	Once, Pre-Procedure
[ ] Partial thromboplastin time	Once, Starting S, Pre-Procedure
[ ] hCG qualitative, serum	Once, Pre-Procedure

#### Cardiology

#### Imaging

#### Other Studies

#### Respiratory

#### Rehab

#### Consults

For Physician Consult orders use sidebar

#### Additional Orders