

General

Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 15 min Every 15 min times 4, then every 30 min times 4, then every 1 hour times 2, then every 4 hours., PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 30 min Every 30 min times 4, then every 1 hour times two, then every 4 hours., PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every hour Every 1 hour times 2, then every 4 hours., PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 4 hours, PACU & Post-op

Activity

<input type="checkbox"/> Strict bed rest	Routine, Until discontinued, Starting S For 4 Hours Keep right leg straight for 4 hours and advance activity as tolerated., PACU & Post-op
--	---

Nursing

<input type="checkbox"/> Assess cath site	Routine, With vitals As needed., PACU & Post-op
<input type="checkbox"/> Apply ice pack	Routine, As needed Affected area: Waking hours only? Nurse to schedule? Special Instructions: Apply to puncture site as needed., PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous At all times., PACU & Post-op
<input type="checkbox"/> Foley catheter - discontinue	Routine, Once For 1 Occurrences Once patient is ambulatory., PACU & Post-op
<input type="checkbox"/> Discharge Home	Routine, Once For 1 Occurrences At ***, if stable. Confirm medication reconciliation is complete., PACU

Notify

<input type="checkbox"/> Notify Endovascular Radiology if evidence of bleeding and/or hematoma around puncture site	Routine, Until discontinued, Starting S, Post-op
---	--

Diet

<input type="checkbox"/> NPO except meds	Diet effective now, Starting S For 1 Hours NPO: Except meds Pre-Operative fasting options: PACU & Post-op
<input type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op

Tube feeding

Diet effective now, Starting S
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Schedule:
Tube Feeding Schedule:
Dietitian to manage Tube Feed?
PACU & Post-op

Medications

Toradol IV

ketorolac (TORADOL) IV (Single Response)

Do NOT use in patients with eGFR LESS than 30 mL/min AND/OR patients LESS than 17 years of age.

WARNING: Use is contraindicated for treatment of perioperative pain OR in the setting of coronary artery bypass graft (CABG) surgery.

- | | |
|---|--|
| () For patients ages GREATER than 64 OR weight LESS than 50 kg OR eGFR 30-59 mL/min - ketorolac (TORADOL) injection | 15 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6), For 5 Days |
| () For patients ages 17-64 AND weight GREATER than or EQUAL to 50 kg AND eGFR at least 60 mL/min - ketorolac (TORADOL) injection | 30 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6), For 5 Days |

Medications

docusate sodium (COLACE) capsule

100 mg, oral, 2 times daily, PACU & Post-op

ondansetron (ZOFTRAN) injection

4 mg, intravenous, every 8 hours PRN, nausea, vomiting, PACU & Post-op

acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet

1 tablet, oral, every 4 hours PRN, moderate pain (score 4-6), PACU & Post-op
give if patient able to tolerate oral medications

acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet

2 tablet, oral, every 4 hours PRN, severe pain (score 7-10), PACU & Post-op
give if patient able to tolerate oral medications