

General

Nursing

Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 15 min Every 15 min times 4, then every 30 min times 4, then every 1 hour times 2, then every 4 hours., PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 30 min Every 30 min times 4, then every 1 hour times two, then every 4 hours., PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every hour Every 1 hour times 2, then every 4 hours., PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 4 hours, PACU & Post-op

Activity

<input type="checkbox"/> Bed rest	Routine, Until discontinued, Starting S For 2 Hours Bathroom Privileges: PACU & Post-op
-----------------------------------	---

Nursing

<input type="checkbox"/> Assess operative site	Routine, With vitals Assess puncture/surgical site with vital signs, PACU & Post-op
<input type="checkbox"/> Change dressing	Routine, Every 72 hours For central venous catheter. Change dressing and needleless connector every Tuesday and Friday or as needed., PACU & Post-op
<input type="checkbox"/> Head of bed 45 degrees	Routine, Until discontinued, Starting S Head of bed: 45 degrees PACU & Post-op
<input type="checkbox"/> Ok to use - portacath	Routine, Until discontinued, Starting S Device: Portacath Port has been flushed with Heparin: aspirate before use, PACU & Post-op
<input type="checkbox"/> Ok to use - central venous catheter	Routine, Until discontinued, Starting S Device: Other Other: Central venous catheter PACU & Post-op
<input type="checkbox"/> Ok to use - dialysis access	Routine, Until discontinued, Starting S Device: Dialysis Access PACU & Post-op
<input type="checkbox"/> Ok to use - tunneled central venous cath	Routine, Until discontinued, Starting S Device: Other Other: tunneled central venous cath PACU & Post-op
<input type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 8 hours, PACU & Post-op Flush each lumen with 10 milliliters normal saline after each use or every 8 hours.
<input type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care, PACU & Post-op Flush after every use, flush each lumen with 10 mL normal saline
<input type="checkbox"/> Nursing communication	Routine, Until discontinued, Starting S Instructions: ***, PACU & Post-op
<input type="checkbox"/> Discharge Home	Routine, Once For 1 Occurrences At ***, if stable. Confirm medication reconciliation is complete., PACU

Diet

<input type="checkbox"/> NPO except meds	Diet effective now, Starting S For 1 Hours NPO: Except meds Pre-Operative fasting options: PACU & Post-op
<input type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op
<input type="checkbox"/> Tube feeding	Diet effective now, Starting S Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Schedule: Tube Feeding Schedule: Dietitian to manage Tube Feed? PACU & Post-op

Notify

<input type="checkbox"/> Notify Endovascular Radiology of worsening abdominal pain or bleeding from cath site.	Routine, Until discontinued, Starting S, Please call 713-441-6540., PACU & Post-op
--	--

IV Fluids

Medications

Post Procedure Medications

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25 mcg, intravenous, every 15 min PRN, severe pain (score 7-10), persistent pain, For 2 Doses, PACU & Post-op Hold for respirations LESS than 12 breath/minute. For persistent pain, call interventional MD.
<input type="checkbox"/> ondansetron (ZOFTRAN) IV	4 mg, intravenous, once PRN, nausea, vomiting, For 1 Doses, PACU & Post-op

VTE

Labs

Cardiology

Diagnostic Imaging

Other Diagnostic Studies

Respiratory

Rehab

Consults

Additional Orders