

IR Contrasts [1467]

Nursing

IV Fluids

IV Fluids

[] sodium chloride 0.9 % bolus	50 mL, intravenous, for 1 Minutes, once in imaging, Flush, For 1 Doses
---------------------------------	--

Medications

IR Contrasts

[] iohexol (OMNIPAQUE) 180 mg iodine/mL injection	intrathecal, once in imaging, contrast, For 1 Doses
[] iohexol (OMNIPAQUE) 240 mg iodine/mL injection	intravenous, once in imaging, contrast, For 1 Doses
[] iohexol (OMNIPAQUE) 300 mg iodine/mL injection	intravenous, once in imaging, contrast, For 1 Doses
[] iohexol (OMNIPAQUE) 350 mg iodine/mL injection	intravenous, once in imaging, contrast, For 1 Doses
[] iodixanol (VISIPAQUE) 270 mg iodine/mL injection	50 mL, intravenous, once in imaging, contrast, For 1 Doses
[] iodixanol (VISIPAQUE) 320 mg iodine/mL injection	intravenous, once in imaging, contrast, For 1 Doses
[] carbon dioxide (CO2) gas	intra-arterial, once in imaging, contrast, For 1 Doses

XR Arthrograms - HMH, HMSJ, HMSTC Only

[] gadopentetate dimeglumine (MAGNEVIST) 5 mL vial solution	5 mL, intra-articular, once in imaging, contrast, xr arthrogram, For 1 Doses
[] iohexol (OMNIPAQUE 180)	20 mL, intra-arterial, once in imaging, contrast, For 1 Doses Intra-articular route
[] iohexol (OMNIPAQUE 300)	30 mL, Other, once in imaging, contrast, xr arthrogram, For 1 Doses Intra-articular route
[] lidocaine 1% buffered with 8.4% sodium bicarbonate injection	5 mL, Other, once in imaging, xr arthrogram, For 1 Doses Intra-articular route

XR Arthrograms - HMW Only

[] gadopentetate dimeglumine (MAGNEVIST) 5 mL vial solution	5 mL, intra-articular, once in imaging, contrast, xr arthrogram, For 1 Doses
[] gadobutrol (GADAVIST) 7.5 mL vial solution	7.5 mL, intravenous, once in imaging, contrast, For 1 Doses
[] iohexol (OMNIPAQUE 180)	20 mL, intra-arterial, once in imaging, contrast, For 1 Doses Intra-articular route
[] iohexol (OMNIPAQUE 300)	30 mL, Other, once in imaging, contrast, xr arthrogram, For 1 Doses Intra-articular route
[] lidocaine 1% buffered with 8.4% sodium bicarbonate injection	5 mL, Other, once in imaging, xr arthrogram, For 1 Doses Intra-articular route

XR Arthrograms - HMSL Only

[] gadopentetate dimeglumine (MAGNEVIST) 5 mL vial solution	5 mL, intra-articular, once in imaging, contrast, xr arthrogram, For 1 Doses
[] iohexol (OMNIPAQUE 180)	20 mL, intra-arterial, once in imaging, contrast, For 1 Doses Intra-articular route
[] iohexol (OMNIPAQUE 300)	30 mL, Other, once in imaging, contrast, xr arthrogram, For 1 Doses Intra-articular route
[] lidocaine 1% buffered with 8.4% sodium bicarbonate injection	5 mL, Other, once in imaging, xr arthrogram, For 1 Doses Intra-articular route
[] lidocaine PF (XYLOCAINE) 10 mg/mL (1 %) injection	5 mL, Other, once in imaging, xr arthrogram, For 1 Doses Intra-articular route
[] gadobutrol (GADAVIST) vial solution	7.5 mL, intravenous, once in imaging, contrast, xr arthrogram, For 1 Doses
[] methylPREDNISolone acetate (DEPO-MEDROL) injection	40 mg, intra-articular, once in imaging, xr arthrogram, For 1 Doses

[] bupivacaine (PF) (MARCAINE) 0.25 % (2.5 mg/mL) solution	injection, once in imaging, For 1 Doses
---	---

XR Arthrograms - HMWB Only

[] gadopentetate dimeglumine (MAGNEVIST) 5 mL vial solution	5 mL, intra-articular, once in imaging, contrast, xr arthrogram, For 1 Doses
[] iohexol (OMNIPAQUE 180)	20 mL, intra-arterial, once in imaging, contrast, For 1 Doses Intra-articular route
[] iohexol (OMNIPAQUE 300)	30 mL, Other, once in imaging, contrast, xr arthrogram, For 1 Doses Intra-articular route
[] lidocaine 2% buffered with 8.4% sodium bicarbonate injection	5 mL, Other, once in imaging, xr arthrogram, For 1 Doses Intra-articular route

XR Arthrograms - HMSTJ Only

[] gadopentetate dimeglumine (MAGNEVIST) 5 mL vial solution	5 mL, intra-articular, once in imaging, contrast, xr arthrogram, For 1 Doses
[] gadobutrol (GADAVIST) 10 mL vial solution	10 mL, intra-articular, once in imaging, contrast, xr arthrogram, For 1 Doses
[] iohexol (OMNIPAQUE 180)	20 mL, intra-arterial, once in imaging, contrast, For 1 Doses Intra-articular route
[] iohexol (OMNIPAQUE 300)	30 mL, Other, once in imaging, contrast, xr arthrogram, For 1 Doses Intra-articular route
[] lidocaine 1% buffered with 8.4% sodium bicarbonate injection	5 mL, Other, once in imaging, xr arthrogram, For 1 Doses

Additional Orders