

HM IP Labor Assessment, Admission and Delivery Nurse Protocol [1520]

General

Admission Order (Single Response)

(X) Admit to L&D

Admitting Physician:
Diagnosis: Pregnancy
Bed request comments:

Nursing

Activity

[X] Strict bed rest

Routine, Until discontinued, Starting S
If membranes ruptured and head not engaged., L&D
Pre-Delivery

[X] Bed rest with bathroom privileges

Routine, Until discontinued, Starting S
Bathroom Privileges: with bathroom privileges
L&D Pre-Delivery

Consent

[X] Complete consent form

Routine, Once
Procedure:
Diagnosis/Condition:
Physician:
Vaginal Delivery of fetus and placenta with possible
episiotomy and possible Cesarean Section with possible use
of blood products., L&D Pre-Delivery

Diet

[X] NPO (except ice chips)

Diet effective now, Starting S
NPO: Except Ice chips
Pre-Operative fasting options:
L&D Pre-Delivery

Nursing - General

[X] Monitor fetal heart tones (intermittent)

Routine, Once
Type: Intermittent
x 30 minutes, or as needed to obtain a Reactive NST, L&D
Pre-Delivery

[X] Monitor fetal heart tones (continuous)

Routine, Once
Type: Continuous
Internal scalp electrode, if needed to monitor fetus., L&D
Pre-Delivery

[X] Sterile vaginal exam

Routine, Once
if contracting, Contact MD prior to SVE if patient is bleeding,
L&D Pre-Delivery

[X] Sterile vaginal exam

Routine, Once
R/O suspected SROM, L&D Pre-Delivery

[X] Intake and output

Routine, Every 8 hours, L&D Pre-Delivery

[X] Straight cath

Routine, Per unit protocol
PRN if unable to void, L&D Pre-Delivery

[X] Insert and maintain Foley

[X] Insert Foley catheter

Routine, Once
Type:
Size:
Urinometer needed:
L&D Pre-Delivery

[X] Foley Catheter Care

Routine, Until discontinued, Starting S
Orders: Maintain
L&D Pre-Delivery

[X] Patient may have epidural	Routine, Once at 3cm dilation. , L&D Pre-Delivery
Notify	
[X] Notify MD (Specify)	Routine, Once For 1 Occurrences, when initial assessment is complete OR within one hour, L&D Pre-Delivery
[X] Notify Pediatrician	Routine, Once For 1 Occurrences, immediately for any newborn concerns, L&D Pre-Delivery

IV Fluids

IV Fluids

[X] lactated Ringer's bolus	1,000 mL, intravenous, for 30 Minutes, once PRN, Fluid Bolus, For 1 Doses, L&D Pre-Delivery
[X] lactated Ringer's infusion	125 mL/hr, intravenous, every 8 hours, L&D Pre-Delivery

Medications

Bleeding Medications

[X] oxytocin (PITOCIN) infusion	42 mL/hr, intravenous, continuous, For 8 Hours, L&D Pre-Delivery
[X] methylergonovine (METHERGINE) injection	200 mcg, intramuscular, once PRN, Bleeding that saturates a pad in LESS than 15 minutes, if BP LESS than 140/90 during labor, For 1 Doses, L&D Pre-Delivery Administer after placenta as need for excessive bleeding, if BP LESS than 140/90 during labor and notify physician.
[X] carboprost (HEMABATE) injection	250 mcg, intramuscular, once PRN, Bleeding that saturates a pad in less than 15 minutes for patients with B/P greater than 140/90, or if bleeding continues to saturate a pad in less than 15 minutes following administration of Methergine, For 1 Doses, L&D Pre-Delivery

Moderate Pain (Pain Score 4-6) (Single Response)

() butorphanol (STADOL) injection	1 mg, intravenous, once PRN, moderate pain (score 4-6), For 1 Doses, L&D Pre-Delivery
------------------------------------	---

Severe Pain (Pain Score 7-10) (Single Response)

() butorphanol (STADOL) injection	2 mg, intravenous, once PRN, severe pain (score 7-10), For 1 Doses, L&D Pre-Delivery
------------------------------------	--

Antiemetics - HMH, HMSJ, HMW, HMSTC, HMTW Only

[X] ondansetron (ZOFTRAN) IV or Oral	"Or" Linked Panel
[X] ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is able to tolerate oral medication.
[X] ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
[X] promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.

[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral medication.
--	---

Antiemetics - HML, HMWB Only

[X] ondansetron (ZOFRAN) IV or Oral	"Or" Linked Panel 4 mg, oral, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is able to tolerate oral medication.
[X] ondansetron ODT (ZOFRAN-ODT) disintegrating tablet	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] ondansetron (ZOFRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel 12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9 % 0.9 % 20 mL for Alaris pump syringe option	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics - HMSTJ Only

[X] ondansetron (ZOFRAN) IV or Oral	"Or" Linked Panel 4 mg, oral, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is able to tolerate oral medication.
[X] ondansetron ODT (ZOFRAN-ODT) disintegrating tablet	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] ondansetron (ZOFRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IVPB or Oral or Rectal	"Or" Linked Panel 12.5 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral medication.

VTE

Labs

Lab - Pt with Prenatal Record

[X] CBC with differential	Once For 1 Occurrences, L&D Pre-Delivery
[X] Syphilis treponemal IgG	Once For 1 Occurrences, L&D Pre-Delivery
[X] Type and Screen, Obstetrical Patient	Once For 1 Occurrences, L&D Pre-Delivery
[X] Rapid HIV 1 & 2	Once For 1 Occurrences, L&D Pre-Delivery

[X] Fern test	Once For 1 Occurrences, Vaginal fluid, L&D Pre-Delivery
[X] POC Amnisure	Once For 1 Occurrences R/O suspected SROM, L&D Pre-Delivery
[X] Hepatitis B surface antigen	Once For 1 Occurrences With patient consent for patients with prenatal records, L&D Pre-Delivery

Lab - Pt with Prenatal Record

[X] CBC with differential	Once For 1 Occurrences, L&D Pre-Delivery
[X] Syphilis treponemal IgG	Once For 1 Occurrences, L&D Pre-Delivery
[X] Type and Screen, Obstetrical Patient	Once For 1 Occurrences, L&D Pre-Delivery
[X] Rapid HIV 1 & 2	Once For 1 Occurrences, L&D Pre-Delivery
[X] Fern test	Once For 1 Occurrences, Vaginal fluid, L&D Pre-Delivery
[X] POC nitrazine	Once For 1 Occurrences R/O suspected SROM, L&D Pre-Delivery
[X] Hepatitis B surface antigen	Once For 1 Occurrences With patient consent for patients with prenatal records, L&D Pre-Delivery

Lab - Pt With No Prenatal Record

[X] Rubella antibody, IgG	Once For 1 Occurrences If no prenatal record, L&D Pre-Delivery
[X] Urine drugs of abuse screen	Once For 1 Occurrences If no prenatal record, L&D Pre-Delivery

Cardiology

Imaging

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders