

General

Nursing

Vital Signs

Vital signs - T/P/R/BP Routine, Per unit protocol

IV Fluids

Peripheral IV Access

<input checked="" type="checkbox"/> Initiate and maintain IV	
<input checked="" type="checkbox"/> Insert peripheral IV	Routine, Once
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care

IV Fluids

<input type="checkbox"/> sodium chloride 0.9 % bolus 1000 mL	1,000 mL, intravenous, for 30 Minutes, once, For 1 Doses
<input type="checkbox"/> sodium chloride 0.9 % bolus 500 mL	500 mL, intravenous, for 15 Minutes, once, For 1 Doses
<input type="checkbox"/> sodium chloride 0.9 % infusion - 100 mL/hr	100 mL/hr, intravenous, continuous

Medications

Antiemetics

<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet - 4 mg	4 mg, oral, once, For 1 Doses Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection - 4 mg	4 mg, intravenous, once, For 1 Doses Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IV - 12.5 mg	12.5 mg, intravenous, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet - 12.5 mg	12.5 mg, oral, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository - 12.5 mg	12.5 mg, rectal, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics

<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet - 4 mg	4 mg, oral, once PRN, nausea, vomiting, For 1 Doses Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) injection - 4 mg	4 mg, intravenous, once PRN, nausea, vomiting, For 1 Doses Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9 % 20 mL for Alaris pump syringe option - 12.5 mg	12.5 mg, intravenous, once PRN, nausea, vomiting, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet - 12.5 mg	12.5 mg, oral, once PRN, nausea, vomiting, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository - 12.5 mg	12.5 mg, rectal, once PRN, nausea, vomiting, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics

<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet - 4 mg	4 mg, oral, once, For 1 Doses Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection - 4 mg	4 mg, intravenous, once, For 1 Doses Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB - 12.5 mg	12.5 mg, intravenous, for 30 Minutes, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet - 12.5 mg	12.5 mg, oral, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository - 12.5 mg	12.5 mg, rectal, once, For 1 Doses Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

VTE

Labs

Laboratory STAT

<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences
<input type="checkbox"/> Reticulocyte count	STAT For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	STAT For 1 Occurrences
<input type="checkbox"/> Hepatic function panel	STAT For 1 Occurrences
<input type="checkbox"/> Troponin	STAT For 1 Occurrences
<input type="checkbox"/> Amylase	STAT For 1 Occurrences
<input type="checkbox"/> Lipase	STAT For 1 Occurrences
<input type="checkbox"/> Lactate dehydrogenase, LDH	STAT For 1 Occurrences
<input type="checkbox"/> Urine drugs of abuse screen	STAT For 1 Occurrences
<input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	STAT For 1 Occurrences Specimen Source: Urine Specimen Site:

Laboratory-STAT

<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences
<input type="checkbox"/> Reticulocyte count	STAT For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	STAT For 1 Occurrences
<input type="checkbox"/> Hepatic function panel	STAT For 1 Occurrences
<input type="checkbox"/> Troponin, I-Stat	STAT For 1 Occurrences
<input type="checkbox"/> Amylase	STAT For 1 Occurrences
<input type="checkbox"/> Lipase	STAT For 1 Occurrences
<input type="checkbox"/> Lactate dehydrogenase, LDH	STAT For 1 Occurrences
<input type="checkbox"/> Urine drugs of abuse screen	STAT For 1 Occurrences
<input type="checkbox"/> Urine Culture and Urinalysis	"And" Linked Panel
<input type="checkbox"/> Urine culture	Once For 1 Occurrences, Urine
<input type="checkbox"/> Urinalysis	STAT For 1 Occurrences

Laboratory - STAT

<input type="checkbox"/> CBC with differential	STAT For 1 Occurrences
<input type="checkbox"/> Reticulocyte count	STAT For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	STAT For 1 Occurrences
<input type="checkbox"/> Hepatic function panel	STAT For 1 Occurrences
<input type="checkbox"/> Troponin, I-Stat	STAT For 1 Occurrences
<input type="checkbox"/> Amylase	STAT For 1 Occurrences
<input type="checkbox"/> Lipase	STAT For 1 Occurrences
<input type="checkbox"/> Lactate dehydrogenase, LDH	STAT For 1 Occurrences
<input type="checkbox"/> Urinalysis	STAT For 1 Occurrences
<input type="checkbox"/> Urine culture	STAT For 1 Occurrences, Urine

Pregnancy Labs

<input type="checkbox"/> hCG QUALitative, serum	STAT For 1 Occurrences
<input type="checkbox"/> hCG QUALitative, urine	STAT For 1 Occurrences

Microbiology

<input type="checkbox"/> Blood culture x 2	"And" Linked Panel
<input type="checkbox"/> Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.
<input type="checkbox"/> Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.

Cardiology

EKG

<input type="checkbox"/> ED bedside monitoring	STAT, Continuous
<input type="checkbox"/> ECG 12 lead	STAT, Once Clinical Indications: Interpreting Physician: To be performed by ED Staff - Show immediately to MD

Imaging

Diagnostics X-Ray

<input type="checkbox"/> Chest 1 Vw	STAT, 1 time imaging For 1 Occurrences
<input type="checkbox"/> Chest 2 Vw	STAT, 1 time imaging For 1
<input type="checkbox"/> Abdomen Acute Inc Chest	STAT, 1 time imaging For 1

Other Studies

Respiratory

Respiratory (Single Response)

<input type="checkbox"/> Oxygen therapy	STAT, Continuous Device 1: Nasal Cannula Rate in liters per minute: 2 Lpm Rate in tenths of a liter per minute: O2 %: Device 2: Device 3: Titrate to keep O2 Sat Above: 92% Indications for O2 therapy:
---	---

Rehab

Consults

Additional Consults can be entered below in search field.

Ancillary Consults

Note: Due to the current resource constraints, consultation orders received after 10 AM Monday - Friday will be seen the following business day.

Day of discharge consults will not be accepted.
Reason for Consult?
Consult Tracking:

Additional Orders