

## General

## Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Post-op
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Post-op
<input type="checkbox"/>	Acute Renal Failure	Post-op
<input type="checkbox"/>	Acute Respiratory Failure	Post-op
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
<input type="checkbox"/>	Anemia	Post-op
<input type="checkbox"/>	Bacteremia	Post-op
<input type="checkbox"/>	Bipolar disorder, unspecified	Post-op
<input type="checkbox"/>	Cardiac Arrest	Post-op
<input type="checkbox"/>	Cardiac Dysrhythmia	Post-op
<input type="checkbox"/>	Cardiogenic Shock	Post-op
<input type="checkbox"/>	Decubitus Ulcer	Post-op
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Post-op
<input type="checkbox"/>	Disorder of Liver	Post-op
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Post-op
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Post-op
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Post-op
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Post-op
<input type="checkbox"/>	Other Alteration of Consciousness	Post-op
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Post-op
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Post-op
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Post-op
<input type="checkbox"/>	Protein-calorie Malnutrition	Post-op
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Post-op
<input type="checkbox"/>	Schizophrenia Disorder	Post-op
<input type="checkbox"/>	Sepsis	Post-op
<input type="checkbox"/>	Septic Shock	Post-op
<input type="checkbox"/>	Septicemia	Post-op
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Post-op

## Elective Outpatient, Observation, or Admission (Single Response)

<input type="checkbox"/>	Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

**Admission or Observation (Single Response)**

Patient has active outpatient status order on file

- |  |  |
|--|--|
| <input type="checkbox"/> Admit to Inpatient  | Diagnosis:<br>Admitting Physician:<br>Level of Care:<br>Patient Condition:<br>Bed request comments:<br>Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.<br>PACU & Post-op |
| <input type="checkbox"/> Outpatient observation services under general supervision | Diagnosis:<br>Admitting Physician:<br>Patient Condition:<br>Bed request comments:<br>PACU & Post-op  |
| <input type="checkbox"/> Outpatient in a bed - extended recovery                   | Diagnosis:<br>Admitting Physician:<br>Bed request comments:<br>PACU & Post-op  |
| <input type="checkbox"/> Transfer patient  | Level of Care:<br>Bed request comments:<br>Scheduling/ADT  |
| <input type="checkbox"/> Return to previous bed                                    | Routine, Until discontinued, Starting S, Scheduling/ADT  |

**Admission (Single Response)**

Patient has active status order on file

- |   |  |
|---|--|
| <input type="checkbox"/> Admit to inpatient     | Diagnosis:<br>Admitting Physician:<br>Level of Care:<br>Patient Condition:<br>Bed request comments:<br>Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.<br>PACU & Post-op |
| <input type="checkbox"/> Transfer patient       | Level of Care:<br>Bed request comments:<br>Scheduling/ADT  |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT  |

**Transfer (Single Response)**

Patient has active inpatient status order on file

- |   |   |
|---|---|
| <input type="checkbox"/> Transfer patient       | Level of Care:<br>Bed request comments:<br>Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT   |

**Code Status**

- |   |  |
|---|--|
| <input type="checkbox"/> Full Code                | Code Status decision reached by:<br>Post-op            |
| <input type="checkbox"/> DNR (Do Not Resuscitate) |  |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Does patient have decision-making capacity?<br>Post-op |

<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

### Isolation

<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

### Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

## Nursing

### Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP (Q 1 hour x 4)	Routine, Every hour For 4 Occurrences Then every 4 hours, Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP (Q 4 hours)	Routine, Every 4 hours, Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP (Per unit protocol)	Routine, Per unit protocol, Post-op

### Activity

<input type="checkbox"/> Bed rest	Routine, Until discontinued, Starting S Head of bed no more than 30 Degrees, may dangle at bedside PRN, Post-op
<input type="checkbox"/> Bed rest	Routine, Until discontinued, Starting S Elevate head of bed, may dangle at bedside PRN; advance as tolerated, Post-op
<input type="checkbox"/> Dangle at bedside	Routine, As needed, Post-op
<input type="checkbox"/> Activity (specify)	Routine, Until discontinued, Starting S Specify: Up in chair Additional modifier: Post-op
<input type="checkbox"/> Ambulate	Routine, 3 times daily Specify: with assistance Post-op
<input type="checkbox"/> Ambulate	Routine, 3 times daily Specify: Independently, Post-op

### Nursing

<input type="checkbox"/> Complete IV infusing from operating room, then change to ordered IV fluid	Routine, Until discontinued, Starting S, Post-op
--	--

<input type="checkbox"/> Saline lock IV	Routine, Continuous If there is a central line or when tolerating oral fluid intake., Post-op
<input type="checkbox"/> Normal Saline Flush	Routine, Continuous To maintain IV patency, Post-op
<input type="checkbox"/> Intake and output	Routine, Every 4 hours, Post-op
<input type="checkbox"/> Intake and output	Routine, Every shift, Post-op

### Tubes and Drains

<input type="checkbox"/> Chest tube insertion setup	Routine, Once, Post-op
<input type="checkbox"/> Foley catheter care	Routine, Until discontinued, Starting S Orders: Maintain To gravity, Post-op
<input type="checkbox"/> Nasogastric tube insert and maintain	
<input type="checkbox"/> Nasogastric tube insertion	Routine, Once Type: Post-op
<input type="checkbox"/> Nasogastric tube maintenance	Routine, Until discontinued, Starting S Tube Care Orders: Post-op

### Nursing Interventions

<input type="checkbox"/> Place antiembolic stockings	Routine, Once Bilateral, knee high, Post-op
<input type="checkbox"/> Reinforce dressing	Routine, As needed Reinforce with: Post-op
<input type="checkbox"/> Change dressing	Routine, As needed, Post-op
<input checked="" type="checkbox"/> Incentive spirometry	Routine, Every hour while awake 10 deep puffs every hour, Post-op

### Notify

<input type="checkbox"/> Notify Physician for temperature above 101 F; systolic blood pressure above 180; diastolic blood pressure above 100 or below 60; pulse is above 120 or below 60	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Notify Physician if patient unable to void within 6 hours of discontinuing foley catheter	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Notify Physician if patient complains of bladder discomfort after discontinuing foley	Routine, Until discontinued, Starting S, Post-op

### Diet

<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options: Sips of water; advance to goal diet of clear liquids, Post-op
<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options: Sips of water when alert, Post-op
<input type="checkbox"/> Diet - Clear Liquids	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op

<input type="checkbox"/> Diet - Full Liquids	Diet effective now, Starting S Diet(s): Full Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op
<input type="checkbox"/> Diet - Regular	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op

### Discharge Planning

<input type="checkbox"/> Assess discharge learning needs	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Patient education (specify)	Routine, Once Patient/Family: Patient Education for: Other (specify) Specify: Post-op urology instructions Post-op
<input type="checkbox"/> Discontinue epidural therapy	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Discontinue PCA	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Discontinue IV	Routine, Once, Post-op
<input type="checkbox"/> Discontinue NG tube	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Foley catheter - discontinue	Routine, Once, Post-op
<input type="checkbox"/> Foley catheter - discontinue	Routine, Once Only after epidural is discontinued, Post-op
<input type="checkbox"/> Discontinue central line	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Prepare for chest tube removal	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Remove dressing	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Discharge criteria	Routine, Until discontinued, Starting S Tolerates diet, ambulates independently, pain controlled with oral medication, passing flatus, patient afebrile, adequate urine output and urine is clear, able to verbalize discharge instructions., Post-op

## IV Fluids

### IV Fluid

<input type="checkbox"/> dextrose 5 % and sodium chloride 0.45 % with potassium chloride 20 mEq/L infusion	intravenous, continuous, Post-op
<input type="checkbox"/> Custom IV Infusion	intravenous, continuous, Post-op

## Medications

### Restricted Medications

<input type="checkbox"/> No ketorolac (TORADOL)	Routine, Until discontinued, Starting S Reason for "No" order: Post-op
---	--

### Mild Pain (Single Response)

(adjust dose for renal/liver function and age)

<input type="checkbox"/> acetaminophen (TYLENOL) tablet OR oral solution	<b>"Or" Linked Panel</b>
Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)	

[ ] acetaminophen (TYLENOL) tablet	650 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op Maximum of 3 grams of acetaminophen per day from all sources. Give the tablet if the patient can tolerate oral medication. (Cirrhosis patients maximum: 2 grams per day from all sources)
[ ] acetaminophen (TYLENOL)suspension	650 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Use if patient cannot tolerate oral tablet.
() ibuprofen (MOTRIN) tablet OR oral solution	<b>"Or" Linked Panel</b> Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury.
[ ] ibuprofen (ADVIL,MOTRIN) tablet	600 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury. Give if patient is able to tolerate oral medication.
[ ] ibuprofen (ADVIL,MOTRIN) 100 mg/5 mL suspension	600 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op Not recommended for patients with eGFR LESS than 30 mL/min or acute kidney injury. Use if patient cannot swallow tablet.
() naproxen (NAPROSYN) tablet - Not recommended for patients with eGFR LESS than 30 mL/min.	250 mg, oral, every 8 hours PRN, mild pain (score 1-3) Not recommended for patients with eGFR LESS than 30 mL/min.

**PRN Oral Medications for Moderate Pain (Pain Score 4-6): For Patients LESS than 65 years old (Single Response)**  
(adjust dose for renal/liver function and age)

() acetaminophen-codeine (TYLENOL #3) tablet OR elixir	<b>"Or" Linked Panel</b> Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
[ ] acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6) Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient is able to tolerate oral medication.
[ ] acetaminophen-codeine 300 mg-30 mg /12.5 mL solution	12.5 mL, oral, every 6 hours PRN, moderate pain (score 4-6) Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) Use if patient cannot swallow tablet.
() HYDROcodone-acetaminophen 5/325 (NORCO) tablet OR elixir	<b>"Or" Linked Panel</b> Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
[ ] HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6)
[ ] HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution	10 mL, oral, every 6 hours PRN, moderate pain (score 4-6)
() HYDROcodone-acetaminophen 7.5/325 (NORCO) tablet OR elixir	<b>"Or" Linked Panel</b> Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)

<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 7.5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6) Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient is able to tolerate oral medication.
<input type="checkbox"/> HYDROcodone-acetaminophen (HYCET) 7.5-325 mg/15 mL solution	15 mL, oral, every 6 hours PRN, moderate pain (score 4-6) Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) Use if patient cannot swallow tablet.
<input type="checkbox"/> HYDROcodone-acetaminophen 10/325 (NORCO) tablet OR elixir	<b>"Or" Linked Panel</b> Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO 10-325) 10-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6) Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient is able to tolerate oral medication.
<input type="checkbox"/> HYDROcodone-acetaminophen (HYCET) 7.5-325 mg/15 mL solution	20 mL, oral, every 6 hours PRN, moderate pain (score 4-6) Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) Use if patient can not swallow tablet.
<input type="checkbox"/> traMADol (ULTRAM) tablet - For eGFR LESS than 30 mL/min, change frequency to every 12 hours)	50 mg, oral, every 6 hours PRN, moderate pain (score 4-6) (Max Daily dose not to exceed 200 mg/day)

**PRN Oral Medications for Moderate Pain (Pain Score 4-6): For Patients GREATER than 65 years old (Single Response)**  
(adjust dose for renal/liver function and age)

<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) tablet OR elixir	<b>"Or" Linked Panel</b> Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6) Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources). Give if patient is able to tolerate oral medication.
<input type="checkbox"/> acetaminophen-codeine 300 mg-30 mg /12.5 mL solution	12.5 mL, oral, every 6 hours PRN, moderate pain (score 4-6) Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources) Use if patient cannot swallow tablet.
<input type="checkbox"/> HYDROcodone-acetaminophen 5/325 (NORCO) tablet OR elixir	<b>"Or" Linked Panel</b> Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6)
<input type="checkbox"/> HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution	10 mL, oral, every 6 hours PRN, moderate pain (score 4-6)
<input type="checkbox"/> traMADol (ULTRAM) tablet - For eGFR LESS than 30 mL/min, change frequency to every 12 hours)	25 mg, oral, every 6 hours PRN, moderate pain (score 4-6) (Max Daily dose not to exceed 200 mg/day)

**PRN IV Medications for Moderate Pain (Pain Score 4-6): For Patients LESS than 65 years old (Single Response)**  
If you select a PCA option above you will not be allowed to also order IV PRN pain medications from this section.  
(adjust dose for renal/liver function and age)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25 mcg, intravenous, every 2 hour PRN, moderate pain (score 4-6)
<input type="checkbox"/> morphine 2 mg/mL injection	2 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6)

<input type="checkbox"/> HYDROmorphone (DILAUDID) injection	0.5 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6)
<input type="checkbox"/> ketorolac (TORADOL) IV (Single Response)	
Do NOT use in patients with eGFR LESS than 30 mL/min AND/OR patients LESS than 17 years of age. WARNING: Use is contraindicated for treatment of perioperative pain OR in the setting of coronary artery bypass graft (CABG) surgery.	
<input type="checkbox"/> For patients ages GREATER than 64 OR weight LESS than 50 kg OR eGFR 30-59 mL/min - ketorolac (TORADOL) injection	15 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6), For 5 Days
<input type="checkbox"/> For patients ages 17-64 AND weight GREATER than or EQUAL to 50 kg AND eGFR at least 60 mL/min - ketorolac (TORADOL) injection	30 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6), For 5 Days

**PRN IV Medications for Moderate Pain (Pain Score 4-6) For Patients GREATER than 65 years old (Single Response)**

If you select a PCA option above you will not be allowed to also order IV PRN pain medications from this section.  
(adjust dose for renal/liver function and age)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	12.5 mcg, intravenous, every 2 hour PRN, moderate pain (score 4-6)
<input type="checkbox"/> morphine 2 mg/mL injection	1 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6)
<input type="checkbox"/> HYDROmorphone (DILAUDID) injection	0.2 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6)
<input type="checkbox"/> ketorolac (TORADOL) injection - Do not use in patients with eGFR LESS than 30 mL/min.	15 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6), For 5 Days Do not use in patients with eGFR LESS than 30 mL/min.

**PRN Oral Medications for Severe Pain (Pain Score 7-10): For Patients LESS than 65 years old (Single Response)**

(adjust dose for renal/liver function and age)

<input type="checkbox"/> HYDROmorphone (DILAUDID) tablet	2 mg, oral, every 6 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> morphine (MSIR) tablet	15 mg, oral, every 6 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> oxyCODONE (ROXICODONE) immediate release tablet	10 mg, oral, every 6 hours PRN, severe pain (score 7-10)

**PRN Oral Medications for Severe Pain (Pain Score 7-10): For Patients GREATER than 65 years old (Single Response)**

(adjust dose for renal/liver function and age)

<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 7.5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO 10-325) 10-325 mg per tablet	1 tablet, oral, every 6 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> HYDROmorphone (DILAUDID) tablet	2 mg, oral, every 6 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> morphine (MSIR) tablet	15 mg, oral, every 6 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> oxyCODONE (ROXICODONE) immediate release tablet	5 mg, oral, every 6 hours PRN, severe pain (score 7-10)

**PRN IV Medications for Severe Pain (Pain Score 7-10): For Patients LESS than 65 years old (Single Response)**

If you select a PCA option above you will not be allowed to also order IV PRN pain medications from this section.  
(adjust dose for renal/liver function and age)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	50 mcg, intravenous, every 3 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> morphine injection	4 mg, intravenous, every 3 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> HYDROmorphone (DILAUDID) injection	0.8 mg, intravenous, every 3 hours PRN, severe pain (score 7-10)

**PRN IV Medications for Severe Pain (Pain Score 7-10): For Patients GREATER than 65 years old (Single Response)**

If you select a PCA option above you will not be allowed to also order IV PRN pain medications from this section.  
(adjust dose for renal/liver function and age)



<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25 mcg, intravenous, every 3 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> morphine injection	2 mg, intravenous, every 3 hours PRN, severe pain (score 7-10)
<input type="checkbox"/> HYDROmorphine (DILAUDID) injection	0.5 mg, intravenous, every 3 hours PRN, severe pain (score 7-10)

**Scheduled Pain Medications - Mild Pain (Pain Score 1-3): For Patients GREATER than 65 years old (Single Response)**  
 (Consider scheduled options if pain source is present and patient unable to reliably communicate needs. Monitor closely for response)

Max Acetaminophen: 3 grams/day; cirrhosis patients: max Acetaminophen 2 grams/day  
 Adjust dose for renal/liver function and age.

<input type="checkbox"/> acetaminophen (TYLENOL) tablet	500 mg, oral, every 6 hours scheduled
---	---------------------------------------

**Scheduled Pain Medications - Moderate Pain (Pain Score 4-6): For Patients GREATER than 65 years old (Single Response)**

(Consider scheduled options if pain source is present and patient unable to reliably communicate needs. Monitor closely for response)

Max Acetaminophen: 3 grams/day; cirrhosis patients: max Acetaminophen 2 grams/day  
 Adjust dose for renal/liver function and age.

<input type="checkbox"/> HYDROcodone-acetaminophen (LORTAB) 2.5-167 mg/5 mL solution	10 mL, oral, every 6 hours scheduled
<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 6 hours scheduled
<input type="checkbox"/> traMADol (ULTRAM) tablet - If eGFR is LESS than 30 mL/min: change frequency to every 12 hours and max daily dose not to exceed 200 mg/day.	25 mg, oral, every 6 hours scheduled If eGFR is LESS than 30 mL/min: change frequency to every 12 hours and max daily dose not to exceed 200 mg/day.

**Scheduled Pain Medications - Severe Pain (Pain Score 7-10): For Patients GREATER than 65 years old (Single Response)**

(Consider scheduled options if pain source is present and patient unable to reliably communicate needs. Monitor closely for response)

Max Acetaminophen: 3 grams/day; cirrhosis patients: max Acetaminophen 2 grams/day  
 Adjust dose for renal/liver function and age.

<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 7.5-325 mg per tablet	1 tablet, oral, every 6 hours scheduled
<input type="checkbox"/> oxyCODONE (ROXICODONE) immediate release tablet	5 mg, oral, every 6 hours scheduled

**Antiemetics**

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.

<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.
--	--

#### Antiemetics - HMSTJ Only

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IVPB or Oral or Rectal	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

#### Antiemetics - HMSL, HMWB Only

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9 % 0.9 % 20 mL for Alaris pump syringe option	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

#### Antispasmodics

<input type="checkbox"/> belladonna alkaloids-opium (B&O SUPPRETTES) 16.2-60 MG suppository	60 mg, rectal, every 6 hours PRN, bladder spasms, Post-op
<input type="checkbox"/> oxybutynin (DITROPAN) tablet	5 mg, oral, 3 times daily PRN, bladder spasms, Post-op

#### Gastrointestinal

<input type="checkbox"/> simethicone (MYLICON) chewable tablet	160 mg, oral, 3 times daily PRN, flatulence, Post-op
--	--

#### Laxatives/Bowel Care

<input type="checkbox"/> bisacodyl (DULCOLAX) suppository	10 mg, rectal, every 8 hours PRN, constipation, Post-op
<input type="checkbox"/> docusate sodium (COLACE) capsule	100 mg, oral, daily PRN, constipation, Post-op

Miscellaneous

acetaminophen (TYLENOL) tablet 650 mg, oral, every 6 hours PRN, fever, temp greater than 101 F, Post-op

VTE

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition  
 Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition  
 Both pharmacologic AND mechanical prophylaxis must be addressed.  
 Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:  
 Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)  
 Age 60 and above Severe fracture of hip, pelvis or leg  
 Central line Acute spinal cord injury with paresis  
 History of DVT or family history of VTE Multiple major traumas  
 Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER  
 Less than fully and independently ambulatory Acute ischemic stroke  
 Estrogen therapy History of PE  
 Moderate or major surgery (not for cancer)  
 Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)  
 Low risk of VTE Routine, Once  
 Low risk: Due to low risk, no VTE prophylaxis is needed.  
 Will encourage early ambulation  
 PACU & Post-op

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk  
 Moderate risk of VTE Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation Routine, Once  
 No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.  
 Therapy for the following:  
 PACU & Post-op

Contraindications exist for pharmacologic prophylaxis Routine, Once  
 No pharmacologic VTE prophylaxis due to the following contraindication(s):  
 PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe 40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min 30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1  
 For Patients with CrCL LESS than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min 30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1  
 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1, PACU & Post-op Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1, PACU & Post-op To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op

#### DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT



<input type="checkbox"/> Low Risk (Single Response)	
<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation PACU & Post-op
<input type="checkbox"/> Moderate Risk of DVT - Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op

<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>

<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>

<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Place antiembolic stockings	Routine, Once, PACU & Post-op
<b>( ) High Risk of DVT - Non-Surgical</b>		
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.		
<b>[ ] High Risk</b>		
<input type="checkbox"/>	High risk of VTE	Routine, Once, PACU & Post-op
<b>[ ] High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)</b>		
<input type="checkbox"/>	Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/>	Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<b>( ) enoxaparin (LOVENOX) injection (Single Response)</b>		
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/>	enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/>	fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/>	heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/>	heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/>	warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/>	Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<b>[ ] Mechanical Prophylaxis (Single Response)</b>		
<input type="checkbox"/>	Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/>	Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Place antiembolic stockings	Routine, Once, PACU & Post-op
<b>( ) High Risk of DVT - Surgical (Hip/Knee)</b>		

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1, PACU & Post-op Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1, PACU & Post-op To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op

## Labs

### Lab Today

<input type="checkbox"/> Hemoglobin	Once, Starting S, Post-op
<input type="checkbox"/> Hematocrit	Once, Starting S, Post-op
<input type="checkbox"/> Basic metabolic panel	Once, Starting S, Post-op

### Lab - Tomorrow in AM

<input type="checkbox"/> Hemoglobin	AM draw For 1 Occurrences, Post-op
<input type="checkbox"/> Hematocrit	AM draw For 1 Occurrences, Post-op
<input type="checkbox"/> CBC hemogram	AM draw For 1 Occurrences, Post-op
<input type="checkbox"/> Electrolyte panel	AM draw For 1 Occurrences, Post-op
<input type="checkbox"/> Creatinine level	AM draw For 1 Occurrences, Post-op
<input type="checkbox"/> Basic metabolic panel	AM draw For 1 Occurrences, Post-op

## Cardiology

## Imaging

### X-Ray

<input type="checkbox"/> Chest 2 Vw W Apical Lordotic	Routine, 1 time imaging For 1 In PACU, Post-op
<input type="checkbox"/> Chest 1 Vw	Routine, 1 time imaging For 1 Upright after chest tube discontinued, Post-op

## Other Studies

## Respiratory

### Respiratory Therapy

<input type="checkbox"/> Turn cough deep breathe	Routine, Every hour While awake, Post-op
<input type="checkbox"/> Turn cough deep breathe	Routine, Every 2 hours While awake, Post-op
<input type="checkbox"/> Incentive spirometry	Routine, Every hour While awake, Post-op
<input type="checkbox"/> Incentive spirometry	Routine, Every 2 hours while awake, Post-op

## Rehab

## Consults

For Physician Consult orders use sidebar

### Ancillary Consults

<input type="checkbox"/> Consult to Case Management	Consult Reason: Post-op
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op

<input type="checkbox"/> Consult PT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult PT wound care	Special Instructions: Location of Wound? Post-op
<input type="checkbox"/> Consult OT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Purpose/Topic: Post-op
<input type="checkbox"/> Consult to Spiritual Care	Reason for consult? Post-op
<input type="checkbox"/> Consult to Speech Language Pathology	Routine, Once Reason for consult: Post-op
<input type="checkbox"/> Consult to Wound Ostomy Care nurse	Reason for consult: Reason for consult: Reason for consult: Reason for consult: Consult for NPWT: Reason for consult: Post-op
<input type="checkbox"/> Consult to Respiratory Therapy	Reason for Consult? Post-op

**Additional Orders**