

# Psychiatry Admission [1769]

## General

### Common Psychiatry Present on Admission Diagnosis

<input type="checkbox"/> Bipolar disorder, unspecified	Details
<input type="checkbox"/> Dementia in Conditions Classified Elsewhere	Details
<input type="checkbox"/> Major Depression Disorder	Details
<input type="checkbox"/> Opioid Withdrawal	Details
<input type="checkbox"/> Other	Details
<input type="checkbox"/> Psychosis, unspecified psychosis type	Details
<input type="checkbox"/> Schizophrenia Disorder	Details
<input type="checkbox"/> Schizoaffective Disorder	Details
<input type="checkbox"/> Substance Induced Mood Disorder	Details

### Admission (Single Response)

<input checked="" type="checkbox"/> Admit to psychiatry	Diagnosis: Admitting Physician: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
---	--

### Code Status

<input type="checkbox"/> Full code	Code Status decision reached by:
<input type="checkbox"/> DNR	
<input type="checkbox"/> DNR (Do Not Resuscitate)	Does patient have decision-making capacity?
<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult:
<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions:
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions:

### Precautions

<input checked="" type="checkbox"/> Suicide precautions	Increased observation level needed:
<input type="checkbox"/> Seizure precautions	Increased observation level needed:
<input type="checkbox"/> Elopement risk	Increased observation level needed:
<input type="checkbox"/> Fall precautions	Increased observation level needed:
<input type="checkbox"/> Aggressive precautions	Increased observation level needed:
<input type="checkbox"/> Bizarre precautions	Increased observation level needed:

## Nursing

### Vital Signs

<input checked="" type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol, Starting S
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 4 hours, Starting S

### Nursing

<input type="checkbox"/> 1:1 patient	Routine, Continuous
<input type="checkbox"/> Clinical institute withdrawal assessment	Routine, Every 4 hours
<input type="checkbox"/> Clinical opiate withdrawal scale (COWS) screening	Routine, Every 4 hours
<input checked="" type="checkbox"/> Search all clothes and belongings	Routine, Once For 1 Occurrences

<input checked="" type="checkbox"/> Safety search	Routine, Per unit protocol Perform a safety search and skin assessment per procedure
---	---

### Restrictions

<input checked="" type="checkbox"/> Restrict to unit	Routine, Continuous
--	---------------------

### Diet

<input type="checkbox"/> Diet - Regular	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
---	--

## IV Fluids

### Medications

#### PRN Anxiety/Agitation (Single Response)

<input type="checkbox"/> diphenhydramine (BENADRYL) tablet	25 mg, oral, every 6 hours PRN, agitation
<input type="checkbox"/> diphenhydramine (BENADRYL) injection	25 mg, intramuscular, every 6 hours PRN, agitation, Severe agitation and Refusal of PO
<input type="checkbox"/> haloperidol (HALDOL) tablet	5 mg, oral, every 6 hours PRN, agitation Indication:
<input type="checkbox"/> haloperidol (HALDOL) injection	5 mg, intramuscular, every 6 hours PRN, agitation, and Refusal of PO. Indication:
<input type="checkbox"/> hydroxyzine (ATARAX) tablet	25 mg, oral, every 4 hours PRN, anxiety
<input type="checkbox"/> lorazepam (ATIVAN) tablet	2 mg, oral, every 6 hours PRN, anxiety, withdrawal, agitation Indication:
<input type="checkbox"/> lorazepam (ATIVAN) injection	2 mg, intramuscular, every 6 hours PRN, agitation, withdrawal, and Refusal of PO
<input type="checkbox"/> chlordiazepoxide (LIBRIUM) capsule	25 mg, oral, every 6 hours PRN, anxiety, withdrawal Indication:
<input type="checkbox"/> olanzapine (ZYPREXA) tablet	oral, daily PRN Indication:
<input type="checkbox"/> olanzapine (ZYPREXA) injection - DO NOT give within 4 hours of IM lorazepam (ATIVAN) injection	intramuscular, daily PRN, agitation NOT to give within 4 hours of IM lorazepam (ATIVAN) injection Indication:
<input type="checkbox"/> ziprasidone (GEODON) injection	intramuscular, daily PRN, agitation Indication:

### Bowel Care

<input type="checkbox"/> loperamide (IMODIUM) capsule	2 mg, oral, every 6 hours PRN, diarrhea
<input type="checkbox"/> alum-mag hydroxide-simeth (MAALOX) 200-200-20 mg/5 mL suspension	30 mL, oral, every 4 hours PRN, indigestion
<input type="checkbox"/> magnesium hydroxide suspension	30 mL, oral, every 12 hours PRN, constipation

#### Sleeping Aid: For Patients LESS than 70 years old (Single Response)

<input type="checkbox"/> ramelteon (ROZEREM) tablet	8 mg, oral, nightly
<input type="checkbox"/> trazodone (DESYREL) tablet	100 mg, oral, nightly Indication:

#### Sleeping Aid: For Patients GREATER than 70 years old

<input type="checkbox"/> ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep
---	--------------------------------

### Nicotine Patch w/ Remove Patch

Dosing suggestions for nicotine patch:

Number of cigarettes per day Dosing (mg/day)

Less than 10, patients with heart disease or weighing less than 100 pounds (45 kg) 7-14

10-20 14-21

21-40 21-42

Greater than 40 42+

NICOTINE PATCH (Single Response)

( ) nicotine (NICODERM CQ) 7 mg/24 hr	1 patch, transdermal, for 24 Hours, daily Time to remove patch:
( ) nicotine (NICODERM CQ) 14 mg/24 hr	1 patch, transdermal, for 24 Hours, daily Time to remove patch:
( ) nicotine (NICODERM CQ) 21 mg/24 hr	1 patch, transdermal, for 24 Hours, daily Time to remove patch:
( ) nicotine (NICODERM CQ) 42 mg/24 hr	2 patch, transdermal, for 24 Hours, daily Time to remove patch:

## VTE

**DVT Risk and Prophylaxis Tool (Single Response)**

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

( ) Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.  
Will encourage early ambulation

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

( ) Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.  
Therapy for the following:

( ) Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

( ) enoxaparin (LOVENOX) injection (Single Response)

( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<b>[ ] Mechanical Prophylaxis (Single Response)</b>	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once
( ) Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<b>[ ] Moderate Risk</b>	
[ ] Moderate risk of VTE	Routine, Once
<b>[ ] Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)</b>	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<b>( ) enoxaparin (LOVENOX) injection (Single Response)</b>	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S

( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input checked="" type="checkbox"/> Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Surgical	

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCl LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.
( ) heparin (porcine) injection	This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 8 hours 5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Surgical (Hip/Knee)	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once

#### DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

( ) Low Risk of DVT	
[ ] Low Risk (Single Response)	
( ) Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
( ) Moderate Risk of DVT - Surgical	

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
( ) Moderate Risk of DVT - Non-Surgical	

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	

( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

( ) High Risk of DVT - Surgical

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
( ) High Risk of DVT - Non-Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily, Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily, Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical) For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
( ) High Risk of DVT - Surgical (Hip/Knee)	

<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

## DVT Risk and Prophylaxis Tool (Single Response)

## Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

### ( ) Low Risk of DVT

#### [ ] Low Risk (Single Response)

( ) Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.  
Will encourage early ambulation

### ( ) Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

#### [ ] Moderate Risk

[ ] Moderate risk of VTE

Routine, Once

#### [ ] Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

( ) Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.  
Therapy for the following:

( ) Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

( ) enoxaparin (LOVENOX) injection (Single Response)

( ) enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min

( ) fondaparinux (ARIXTRA) injection

2.5 mg, subcutaneous, daily, Starting S+1

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
( ) Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCl LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:

( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
[ ] High Risk	
[ ] High risk of VTE	Routine, Once
[ ] High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response)	

( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Non-Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
[ ] High Risk	
[ ] High risk of VTE	Routine, Once
[ ] High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous

<input type="checkbox"/> Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Surgical (Hip/Knee)	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<b>[ ] High Risk</b>	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<b>[ ] Mechanical Prophylaxis (Single Response)</b>	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous

( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once

## Labs

### Laboratory

[X] Hemoglobin A1c	AM draw For 1 Occurrences
[ ] Basic metabolic panel	AM draw For 1 Occurrences
[ ] Comprehensive metabolic panel	AM draw For 1 Occurrences
[X] Lipid panel	AM draw For 1 Occurrences
[ ] Hepatic function panel	AM draw For 1 Occurrences
[ ] CBC with platelet and differential	AM draw For 1 Occurrences
[ ] TSH	AM draw For 1 Occurrences
[ ] T4, free	AM draw For 1 Occurrences
[ ] Ammonia level	AM draw For 1 Occurrences
[ ] Magnesium level	AM draw For 1 Occurrences
[ ] Lithium level	AM draw For 1 Occurrences
[ ] Valproic acid level	AM draw For 1 Occurrences
[ ] Carbamazepine level	AM draw For 1 Occurrences
[ ] Vitamin B12 level	AM draw For 1 Occurrences
[ ] Vitamin D 25 hydroxy level	AM draw For 1 Occurrences
[ ] Urinalysis, automated with microscopy	Once
[ ] hCG qualitative, urine	Once
[ ] Urine drugs of abuse screen	Once
[X] HIV Ag/Ab combination	AM draw For 1 Occurrences
[X] Syphilis treponemal IgG	AM draw For 1 Occurrences
[X] hCG qualitative, serum screen	AM draw For 1 Occurrences

## Cardiology

### Cardiology

[ ] ECG 12 lead	STAT, Once Clinical Indications: Interpreting Physician: Evaluate QT interval
-----------------	--

## Rehab

### Consults

For Physician Consult orders use sidebar

### Ancillary Consults

[X] Consult to Psychiatric OT eval and treat	Special Instructions:
[X] Consult to Social Work	Reason for Consult: Other Specify Specify: Patient may begin to participate in Group and or Psychological education Patient may begin to participate in Group and or Psychological education
[ ] Consult to Case Management	Consult Reason:
[ ] Consult PT eval and treat	Special Instructions: Weight Bearing Status:
[ ] Consult PT wound care	Special Instructions: Location of Wound?
[ ] Consult OT eval and treat	Special Instructions: Weight Bearing Status:
[ ] Consult to Nutrition Services	Reason For Consult? Purpose/Topic:

<input type="checkbox"/> Consult to Spiritual Care	Reason for consult?
<input type="checkbox"/> Consult to Speech Language Pathology	Routine, Once Reason for consult:
<input type="checkbox"/> Consult to Wound Ostomy Care nurse	Reason for consult: Reason for consult: Reason for consult: Reason for consult: Consult for NPWT: Reason for consult:
<input type="checkbox"/> Consult to Respiratory Therapy	Reason for Consult?

## Additional Orders