

General

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Post-op
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Post-op
<input type="checkbox"/>	Acute Renal Failure	Post-op
<input type="checkbox"/>	Acute Respiratory Failure	Post-op
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
<input type="checkbox"/>	Anemia	Post-op
<input type="checkbox"/>	Bacteremia	Post-op
<input type="checkbox"/>	Bipolar disorder, unspecified	Post-op
<input type="checkbox"/>	Cardiac Arrest	Post-op
<input type="checkbox"/>	Cardiac Dysrhythmia	Post-op
<input type="checkbox"/>	Cardiogenic Shock	Post-op
<input type="checkbox"/>	Decubitus Ulcer	Post-op
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Post-op
<input type="checkbox"/>	Disorder of Liver	Post-op
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Post-op
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Post-op
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Post-op
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Post-op
<input type="checkbox"/>	Other Alteration of Consciousness	Post-op
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Post-op
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Post-op
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Post-op
<input type="checkbox"/>	Protein-calorie Malnutrition	Post-op
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Post-op
<input type="checkbox"/>	Schizophrenia Disorder	Post-op
<input type="checkbox"/>	Sepsis	Post-op
<input type="checkbox"/>	Septic Shock	Post-op
<input type="checkbox"/>	Septicemia	Post-op
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Post-op

Elective Outpatient, Observation, or Admission (Single Response)

<input type="checkbox"/>	Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="checkbox"/>	Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments: PACU & Post-op
<input type="checkbox"/>	Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

Admission or Observation (Single Response)

Patient has active outpatient status order on file

- | | |
|--|--|
| <input type="checkbox"/> Admit to Inpatient | Diagnosis:
Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Outpatient observation services under general supervision | Diagnosis:
Admitting Physician:
Patient Condition:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Outpatient in a bed - extended recovery | Diagnosis:
Admitting Physician:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Admission (Single Response)

Patient has active status order on file

- | | |
|---|--|
| <input type="checkbox"/> Admit to inpatient | Diagnosis:
Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Transfer (Single Response)

Patient has active inpatient status order on file

- | | |
|---|---|
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Code Status

- | | |
|---|--|
| <input type="checkbox"/> Full Code | Code Status decision reached by:
Post-op |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Does patient have decision-making capacity?
Post-op |

<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

Isolation

<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Hip precautions	Precaution: Pillow between legs at all times. No bedpan., Post-op
<input type="checkbox"/> Hip precautions - Posterior	Precaution: Pillow between legs at all times. No bedpan., Post-op
<input type="checkbox"/> Hip precautions - Anterior	Precaution: Pillow between legs at all times. No bedpan., Post-op

Nursing

Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP (Q4 hours)	Routine, Every 4 hours For 24 Hours Then every 8 hours., Post-op
--	---

Activity

<input type="checkbox"/> CPM	STAT, Until discontinued, Starting S Location to start: Floor Apply to: CPM range from (degrees): 0 CPM range to (degrees): 90 Daily duration: Other Specify: TID wear as tolerated Advance or Progress: As tolerated Night time use: Advance to 90 degrees as tolerated, Post-op
<input checked="" type="checkbox"/> Up in chair	Routine, As needed Specify: Up in chair Additional modifier: Day of Surgery, Post-op
<input type="checkbox"/> Dangle at bedside	Routine, Once On day of surgery., PACU & Post-op

Nursing Assessments

<input type="checkbox"/> Telemetry	"And" Linked Panel
------------------------------------	--------------------

<input type="checkbox"/> Telemetry monitoring	Routine, Continuous Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box) Reason for telemetry: Can be off of Telemetry for tests and baths? Yes PACU & Post-op
<input type="checkbox"/> Telemetry Additional Setup Information	Routine, Continuous High Heart Rate (BPM): 120 Low Heart Rate(BPM): 50 High PVC's (per minute): 10 High SBP(mmHg): 175 Low SBP(mmHg): 100 High DBP(mmHg): 95 Low DBP(mmHg): 40 Low Mean BP: 60 High Mean BP: 120 Low SPO2(%): 94 PACU & Post-op
<input type="checkbox"/> Neurovascular checks	Routine, Every 4 hours With vital signs.
Nursing Interventions	
<input type="checkbox"/> Turn cough deep breathe	Routine, Every hour While awake, Post-op
<input type="checkbox"/> Cryocuff	Routine, Until discontinued, Starting S For 3 Days Affected area: Cold therapy x 3 days
<input type="checkbox"/> Patient position: Float heels	Routine, Until discontinued, Starting S Position: Additional instructions: Float Heels
<input type="checkbox"/> Patient position: elevate heels and all osseous prominence	Routine, Until discontinued, Starting S Position: Additional instructions: Elevate heels and all osseous prominence to prevent decubitus formation.
<input type="checkbox"/> Insert and Maintain Foley	
<input type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed: Insert as needed for urinary retention leave in. Discontinue Post-Op day 2 in A.M., Post-op
<input type="checkbox"/> Foley Catheter Care	Routine, Until discontinued, Starting S For 48 Hours Orders: to gravity Discontinue post-op day 2 in A.M., Post-op
<input type="checkbox"/> Foley catheter - discontinue	Routine, Once, Starting S+2 at 6:00 AM Post-op day 2 in AM., Post-op
<input type="checkbox"/> Wound care orders - dressing change	Routine, Daily, Starting S+2 Wound care to be performed by: Floor Nurse Location: Site: Irrigate wound? Apply: Dressing Type: Dressing Change Post Operative Day 2 and then daily until dry., Post-op
<input type="checkbox"/> Intake and output	Routine, Every shift For 48 Hours, Post-op
<input type="checkbox"/> Intake and output	Routine, Every shift JP drain to be emptied and recorded every shift., Post-op
<input type="checkbox"/> Saline Lock and Flush	

<input type="checkbox"/> Saline lock	Routine, Continuous If taking PO
<input type="checkbox"/> sodium chloride 0.9 % flush	3 mL, intravenous, PRN, line care Flush every shift and as needed POD#1
<input type="checkbox"/> Nursing communication	Routine, Until discontinued, Starting S, Post-op

Equipment

<input type="checkbox"/> Obtain supply / device:	Routine, Once Obtain: Post-op
--	-------------------------------------

Diet

<input type="checkbox"/> Diet - Clear liquids, advance as tolerated to Regular	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Yes Target Diet: Regular Advance target diet criteria: Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op
<input type="checkbox"/> Diet - Low Sodium	Diet effective now, Starting S Diet(s): Regular, Other Chol/Fat/Sodium Chol/Fat/Sodium: 2 GM Sodium Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Low sodium, PACU & Post-op
<input type="checkbox"/> Diet - Regular	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Please assess bowel sounds between progressions., PACU & Post-op
<input type="checkbox"/> Diet - 1800 Carb Control Diabetic	Diet effective now, Starting S Diet(s): Other Diabetic/Cal Diabetic/Calorie: 1800 Kcal/202 gm Carbohydrate Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op

Notify

<input type="checkbox"/> Notify NP or IM for pain score greater than 7 uncontrolled with PRN medications.	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Notify MD regarding med rec	Routine, Until discontinued, Starting S, To start home meds on Post-Op day 1, Post-op
<input type="checkbox"/> Notify Medicine/NP after Epidural has been discontinued	Routine, Until discontinued, Starting S, For pain score greater than 7, uncontrolled with PRN pain medications., Post-op
<input type="checkbox"/> Notify Physician if Hgb less than or equal to 7 OR if Hgb less than or equal to 8	Routine, Until discontinued, Starting S, And symptomatic of acute anemia

Education

<input type="checkbox"/> Patient education - for self admin of medication	Routine, Once Patient/Family: Patient Education for: Other (specify) Specify: Reinforce home teaching for Lovenox self-injection. Post-op
---	---

IV Fluids

Peripheral IV Access

<input checked="" type="checkbox"/> Initiate and maintain IV	
<input checked="" type="checkbox"/> Insert peripheral IV	Routine, Once
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care

IV Fluids

<input type="checkbox"/> dextrose 5%-0.45% sodium chloride infusion	80 mL/hr, intravenous, for 12.5 Hours, continuous, Post-op Until PO, then Saline Lock
<input type="checkbox"/> lactated ringer's infusion	80 mL/hr, intravenous, for 12.5 Hours, continuous, Post-op Until PO, then Saline Lock

IV Fluids with Potassium

<input type="checkbox"/> dextrose 0.5% in sodium chloride 0.45% with potassium chloride 20 mEq	80 mL/hr, intravenous, for 12.5 Hours, continuous, Post-op Until PO, then Saline Lock
--	---

Medications

Consult Pharmacy

<input type="checkbox"/> Pharmacy consult to manage dose adjustments for renal function	STAT, Until discontinued, Starting S Adjust dose for:
---	--

Vitamin Supplementation

<input type="checkbox"/> ferrous sulfate tablet	325 mg, oral, 3 times daily with meals, Post-op
<input type="checkbox"/> multivitamin (THERAGRAN) per tablet	1 tablet, oral, daily, Post-op

IV Antibiotics: For Patients LESS than or EQUAL to 120 kg

<input type="checkbox"/> cefazolin (ANCEF) IV - For Patients LESS than or EQUAL to 120 kg	2 g, intravenous, once, For 1 Doses, Post-op For patients LESS than or EQUAL to 120 kg. Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> clindamycin (CLEOCIN) IV - For Penicillin Allergic Patients	900 mg, intravenous, for 30 Minutes, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> vancomycin (VANCOGIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis

IV Antibiotics: For Patients GREATER than 120 kg

<input type="checkbox"/> cefazolin (ANCEF) IV - For Patients GREATER than 120 kg	3 g, intravenous, once, For 1 Doses, Post-op For patients GREATER than 120 kg; Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> clindamycin (CLEOCIN) IV - For Penicillin Allergic Patients	900 mg, intravenous, for 30 Minutes, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> vancomycin (VANCOGIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis

Multimodal Pain Control (Single Response)

<input type="checkbox"/> gabapentin (NEURONTIN)	oral, 3 times daily, Post-op
<input type="checkbox"/> celecoxib (CeleBREX) capsule	100 mg, oral, 2 times daily, Post-op

Moderate Pain (Pain Score 4-6) (Single Response)

<input type="checkbox"/> celecoxib (CeleBREX) capsule	200 mg, oral, 2 times daily PRN, moderate pain (score 4-6), Post-op
<input type="checkbox"/> If Sulfa Drug Allergic - naproxen (NAPROSYN) tablet	250 mg, oral, every 8 hours PRN, mild pain (score 1-3), moderate pain (pain score 4-6), Post-op
<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op
<input type="checkbox"/> oxyCODone-acetaminophen (PERCOCET) 5-325 mg per tablet	1 tablet, oral, every 4 hours PRN, moderate pain (score 4-6), Post-op

Severe Pain (Pain Score 7-10)

<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 7.5-325 mg per tablet	2 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op If Anesthesia on case, begin POD #2.
<input type="checkbox"/> oxyCODone-acetaminophen (PERCOCET) 10-325 mg per tablet	1 tablet, oral, every 4 hours PRN, severe pain (score 7-10), Post-op
<input type="checkbox"/> morPHINE injection	1 mg, intravenous, every 4 hours PRN, severe pain (score 7-10), Post-op

Respiratory Depression and Somnolence

<input checked="" type="checkbox"/> naloxone (NARCAN) injection	0.2 mg, intravenous, once PRN, respiratory depression, as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3), For 1 Doses, Post-op Repeat Naloxone 0.2 mg once in 2 minutes if necessary (MAXIMUM 0.4 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times.
---	--

Fever

<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, every 6 hours PRN, headaches, fever, Temperature greater than 101, Post-op
---	--

Antiemetics - HHM, HMSJ, HMW, HMSTC, HMTW Only (Single Response)

<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics - HMSL, HMWB Only

<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is able to tolerate oral medication.
--	---

<input type="checkbox"/> ondansetron (ZOFTRAN) injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9 % 20 mL for Alaris pump syringe option	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics - HMSTJ Only

<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, PACU & Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, PACU & Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Stool Softener

<input checked="" type="checkbox"/> docusate sodium (COLACE) capsule	100 mg, oral, 2 times daily PRN, constipation, for stool softener, Post-op
--	--

Laxatives (Single Response)

<input type="checkbox"/> sennosides-docusate sodium (SENOKOT-S) 8.6-50 mg per tablet	2 tablet, oral, nightly PRN, constipation, Post-op
<input type="checkbox"/> magnesium hydroxide suspension - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR WORSE	30 mL, oral, every 6 hours PRN, constipation, Post-op Do not give if patient is on hemodialysis or is in chronic renal failure.
<input type="checkbox"/> bisacodyl (DULCOLAX) EC tablet	10 mg, oral, daily PRN, constipation, Post-op
<input type="checkbox"/> bisacodyl (DULCOLAX) suppository	10 mg, rectal, daily PRN, constipation, Post-op
<input type="checkbox"/> polyethylene glycol (MIRALAX) packet	17 g, oral, daily PRN, constipation, Post-op

Itching: For Patients LESS than 70 years old (Single Response)

<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet	25 mg, oral, every 6 hours PRN, itching, PACU & Post-op
<input type="checkbox"/> hydrOXYzine (ATARAX) tablet	10 mg, oral, every 6 hours PRN, itching, PACU & Post-op
<input type="checkbox"/> cetirizine (ZyrTEC) tablet	5 mg, oral, daily PRN, itching, PACU & Post-op
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet - For eGFR LESS than 80 mL/min, reduce frequency to once daily as needed	60 mg, oral, 2 times daily PRN, itching, PACU & Post-op

Itching: For Patients BETWEEN 70 to 76 years old (Single Response)

cetirizine (ZyrTEC) tablet 5 mg, oral, daily PRN, itching, PACU & Post-op

Itching: For Patients GREATER than 77 years old (Single Response)

cetirizine (ZyrTEC) tablet 5 mg, oral, daily PRN, itching, PACU & Post-op

Insomnia: For Patients GREATER than or EQUAL to 70 years old (Single Response)

ramelteon (ROZEREM) tablet 8 mg, oral, nightly PRN, sleep, PACU & Post-op

Insomnia: For Patients LESS than 70 years old (Single Response)

zolpidem (AMBIEN) tablet 5 mg, oral, nightly PRN, sleep, PACU & Post-op

ramelteon (ROZEREM) tablet 8 mg, oral, nightly PRN, sleep, PACU & Post-op

VTE**DVT Risk and Prophylaxis Tool (Single Response)**

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.

Will encourage early ambulation

PACU & Post-op

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

PACU & Post-op

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response)

<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op

<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1, PACU & Post-op Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1, PACU & Post-op To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.

Will encourage early ambulation

PACU & Post-op

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.

Therapy for the following:

PACU & Post-op

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCl LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Non-Surgical Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCl LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1, PACU & Post-op Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1, PACU & Post-op To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op

Labs

Labs Today

<input type="checkbox"/> Hemoglobin and hematocrit	STAT For 1 Occurrences, PACU
<input type="checkbox"/> Sodium level	STAT For 1 Occurrences, PACU
<input type="checkbox"/> Potassium level	STAT For 1 Occurrences, PACU

Labs POD 1, 2, 3

<input type="checkbox"/> Hemoglobin and hematocrit	AM draw, Starting S+1 For 1 Occurrences POD 1, Post-op
<input type="checkbox"/> Hemoglobin & hematocrit	AM draw, Starting S+2 For 1 Occurrences POD 2, Post-op
<input type="checkbox"/> CBC hemogram	AM draw, Starting S+2 For 1 Occurrences POD 2, Post-op

Labs Repeat

<input type="checkbox"/> CBC hemogram	AM draw repeats For 3 Occurrences, PACU & Post-op
<input type="checkbox"/> Hemoglobin & hematocrit	AM draw repeats For 3 Occurrences, PACU & Post-op

Cardiology

Imaging

Diagnostic X-Ray

<input type="checkbox"/>	XR Hip 2-3 View Left	Routine, 1 time imaging For 1 , PACU
<input type="checkbox"/>	XR Hip 2-3 View Right	Routine, 1 time imaging For 1 , PACU
<input type="checkbox"/>	XR Knee 3 Vw Left	Routine, 1 time imaging For 1 , PACU
<input type="checkbox"/>	XR Knee 3 Vw Right	Routine, 1 time imaging For 1 , PACU
<input type="checkbox"/>	Pelvis 1 Or 2 Vw	Routine, 1 time imaging For 1 , PACU
<input type="checkbox"/>	Femur 2 Vw Left	Routine, 1 time imaging For 1 , PACU
<input type="checkbox"/>	Femur 2 Vw Right	Routine, 1 time imaging For 1 , PACU

Other Studies

Respiratory

Respiratory

<input type="checkbox"/>	Oxygen therapy	Routine, Continuous Device 1: Nasal Cannula Rate in liters per minute: Rate in tenths of a liter per minute: O2 %: Titrate to keep O2 Sat Above: 90% Indications for O2 therapy: Post-op
<input type="checkbox"/>	Pulse oximetry, overnight	Routine, At bedtime, Post-op
<input checked="" type="checkbox"/>	Incentive spirometry	Routine, Every hour while awake Waking hours only., Post-op

Rehab

Consults

For Physician Consult orders use sidebar

Consults

<input type="checkbox"/>	IR Consult To Interventional Radiology	Routine, 1 time imaging For 1 Occurrences
<input type="checkbox"/>	Consult Pain Management	Reason for Consult? pain management Patient/Clinical information communicated? Patient/clinical information communicated? Post-op
<input type="checkbox"/>	Consult to PT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/>	Consult to PT eval and treat - Weight bearing as needed	Special Instructions: Weight Bearing Status: Weight bearing as tolerated
<input type="checkbox"/>	Consult to PT eval and treat - Non-weight bearing	Special Instructions: Weight Bearing Status: Non-weight bearing
<input type="checkbox"/>	Consult to PT eval and treat - Touch down weight bearing	Special Instructions: Weight Bearing Status: Touch down weight bearing
<input type="checkbox"/>	Consult to OT eval and treat - for ADLs	Special Instructions: for ADLs Weight Bearing Status:
<input type="checkbox"/>	Consult to Case Management - DME: bedside commode and rolling walker	Consult Reason: DME Post-op, bedside commode and rolling walker
<input type="checkbox"/>	Consult to Case Management - DME: Wheelchair	Consult Reason: DME Post-op, Wheelchair
<input type="checkbox"/>	Consult to Case Management - DME: Crutches	Consult Reason: DME Post-op, Crutches
<input type="checkbox"/>	Consult to Case Management - DME: Rolling Walker	Consult Reason: DME Post-op, Rolling Walker
<input type="checkbox"/>	Consult to Case Management - Discharge planning	Consult Reason: Discharge Planning Post-op

Consult to Case Management - Home Health Safety Eval

Consult Reason: Home Health
Post-op, Home Health Safety Eval

Consult to Case Management - Home Health: Physical
Therapy Eval & Treat

Consult Reason: Home Health
Post-op, Home Health: Physical Therapy Eval & Treat

Additional Orders