

General

Common Present on Admission Diagnosis

[] Acidosis	Post-op
[] Acute Post-Hemorrhagic Anemia	Post-op
[] Acute Renal Failure	Post-op
[] Acute Respiratory Failure	Post-op
[] Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
[] Anemia	Post-op
[] Bacteremia	Post-op
[] Bipolar disorder, unspecified	Post-op
[] Cardiac Arrest	Post-op
[] Cardiac Dysrhythmia	Post-op
[] Cardiogenic Shock	Post-op
[] Decubitus Ulcer	Post-op
[] Dementia in Conditions Classified Elsewhere	Post-op
[] Disorder of Liver	Post-op
[] Electrolyte and Fluid Disorder	Post-op
[] Intestinal Infection due to Clostridium Difficile	Post-op
[] Methicillin Resistant Staphylococcus Aureus Infection	Post-op
[] Obstructive Chronic Bronchitis with Exacerbation	Post-op
[] Other Alteration of Consciousness	Post-op
[] Other and Unspecified Coagulation Defects	Post-op
[] Other Pulmonary Embolism and Infarction	Post-op
[] Phlebitis and Thrombophlebitis	Post-op
[] Protein-calorie Malnutrition	Post-op
[] Psychosis, unspecified psychosis type	Post-op
[] Schizophrenia Disorder	Post-op
[] Sepsis	Post-op
[] Septic Shock	Post-op
[] Septicemia	Post-op
[] Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
[] Urinary Tract Infection, Site Not Specified	Post-op

Elective Outpatient, Observation, or Admission (Single Response)

() Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
() Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
() Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments: PACU & Post-op
() Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

Admission or Observation (Single Response)

Patient has active outpatient status order on file

() Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op
() Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
() Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments: PACU & Post-op
() Transfer patient	Level of Care: Bed request comments: Scheduling/ADT
() Return to previous bed	Routine, Until discontinued, Starting S, Scheduling/ADT

Admission (Single Response)

Patient has active status order on file

() Admit to inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op
() Transfer patient	Level of Care: Bed request comments: Scheduling/ADT
() Return to previous bed	Routine, Until discontinued, Starting S, Scheduling/ADT

Transfer (Single Response)

Patient has active inpatient status order on file

() Transfer patient	Level of Care: Bed request comments: Scheduling/ADT
() Return to previous bed	Routine, Until discontinued, Starting S, Scheduling/ADT

Code Status

[] Full Code	Code Status decision reached by: Post-op
[] DNR (Do Not Resuscitate)	Does patient have decision-making capacity? Post-op
[] DNR (Do Not Resuscitate)	Does patient have decision-making capacity? Post-op

<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

Isolation

<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

Nursing

Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP - Per Unit Protocol - PACU	Routine, Per unit protocol, PACU
<input type="checkbox"/> Vital signs - T/P/R/BP - Per Unit Protocol - Post-Op Floor	Routine, Per unit protocol, Post-op

Activity

<input type="checkbox"/> Activity (specify)	Routine, Until discontinued, Starting S Specify: Activity as tolerated PACU
---	---

Notify

<input type="checkbox"/> Notify Physician for vitals:	Routine, Until discontinued, Starting S Temperature greater than: 101.5 Temperature less than: 96 Systolic BP greater than: 180 Systolic BP less than: 90 Diastolic BP greater than: 100 Diastolic BP less than: 40 MAP less than: Heart rate greater than (BPM): 120 Heart rate less than (BPM): 50 Respiratory rate greater than: 22 Respiratory rate less than: 10 SpO2 less than: 90
---	--

<input type="checkbox"/> Notify Resident for vitals:	Routine, Until discontinued, Starting S Temperature greater than: 101.5 Temperature less than: 96 Systolic BP greater than: 180 Systolic BP less than: 90 Diastolic BP greater than: 100 Diastolic BP less than: 40 MAP less than: Heart rate greater than (BPM): 120 Heart rate less than (BPM): 50 Respiratory rate greater than: 22 Respiratory rate less than: 10 SpO2 less than: 90
--	--

Nursing

<input type="checkbox"/> Intake and output	Routine, Every shift, PACU & Post-op
<input type="checkbox"/> Discontinue IV fluids when tolerating by mouth	Routine, Once For 1 Occurrences Flush with NS 5mL every 8 hours and PRN, Post-op
<input type="checkbox"/> Saline lock IV	Routine, Continuous, Post-op
<input type="checkbox"/> Change IV site dressing	Routine, Per unit protocol, Post-op
<input type="checkbox"/> Foley catheter care - to gravity	Routine, Until discontinued, Starting S Orders: to gravity Post-op
<input type="checkbox"/> Meatal care	Routine, Until discontinued, Starting S Every shift with soap and water , Post-op
<input type="checkbox"/> Apply ice pack	Routine, Until discontinued, Starting S Affected area: Waking hours only? Nurse to schedule? Special Instructions: To scrotum once until bedtime, Post-op
<input type="checkbox"/> Incentive spirometry	Routine, Every 2 hours while awake, PACU & Post-op
<input type="checkbox"/> Instruct patient on how to create and use a penile sling	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Supplies at bedside once patient arrives to floor	Routine, Until discontinued, Starting S 20 mL syringe 2 inch silk tape, Post-op
<input type="checkbox"/> Insert and maintain Foley	
<input type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed: Post-op
<input type="checkbox"/> Foley Catheter Care	Routine, Until discontinued, Starting S Orders: Maintain Post-op
<input type="checkbox"/> Perineal care	Routine, 2 times daily, Post-op
<input type="checkbox"/> Foley catheter - discontinue POD #1	Routine, Once, Post-op

Diet

<input type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op
-------------------------------	---

IV Fluids

IV Fluids

<input type="checkbox"/> sodium chloride 0.9% infusion	intravenous, at 125 mL/hr, continuous, Post-op
--	--

<input type="checkbox"/> lactated ringers infusion	intravenous, at 125 mL/hr, continuous, Post-op
<input type="checkbox"/> dextrose 5%-lactated ringers infusion	intravenous, at 125 mL/hr, continuous, Post-op
<input type="checkbox"/> dextrose 5 % and sodium chloride 0.45 % with potassium chloride 20 mEq/L infusion	intravenous, at 125 mL/hr, continuous, Post-op

Medications

Combination Antibiotics (Single Response)

HOP-Surgery approved antibiotic options for penile prosthesis surgery. If patient allergy prevents use of options below, consult pharmacy or infectious disease consultant for alternative options.

<input type="checkbox"/> ampicillin-sulbactam (UNASYN) 3 g injection	3 g, intravenous, once, Starting S, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> gentamicin (GARAMYCIN) IVPB plus ceFAZolin (ANCEF) IVPB - For patients GREATER THAN 120 kg	"And" Linked Panel
<input type="checkbox"/> gentamicin (GARAMICIN) IVPB	5 mg/kg, intravenous, for 60 Minutes, once, For 1 Doses, Pre-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> cefazolin (ANCEF) IVPB	3 g, intravenous, once, Starting S, For 1 Doses, Pre-op Give within 1 hour of procedure. Infuse over 30 minutes. Repeat 8 hours after initial dose if still intra-op. Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> gentamicin (GARAMYCIN) IVPB plus ceFAZolin (ANCEF) IVPB - For patients LESS than or EQUAL to 120 kg	"And" Linked Panel
<input type="checkbox"/> gentamicin (GARAMICIN) IVPB	5 mg/kg, intravenous, for 60 Minutes, once, For 1 Doses, Pre-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> cefazolin (ANCEF) IVPB	2 g, intravenous, once, Starting S, For 1 Doses, Pre-op Give within 1 hour of procedure. Infuse over 30 minutes. Repeat 8 hours after initial dose if still intra-op. Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> gentamicin (GARAMYCIN) IVPB plus clindamycin (CLEOCIN) IVPB - For Penicillin or Vancomycin Allergic patients	"And" Linked Panel
<input type="checkbox"/> gentamicin (GARAMICIN) IVPB	5 mg/kg, intravenous, for 60 Minutes, once, For 1 Doses, Pre-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> clindamycin (CLEOCIN) IVPB	900 mg, intravenous, for 30 Minutes, once, For 1 Doses, Pre-op Give within 1 hour of penile prosthesis procedure. Infuse over 30 minutes. Repeat 3 hours after initial dose if still intra-op. Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> gentamicin (GARAMYCIN) IVPB plus vancomycin (VANCOCIN) IVPB	"And" Linked Panel
<input type="checkbox"/> gentamicin (GARAMICIN) IVPB	5 mg/kg, intravenous, for 60 Minutes, once, For 1 Doses, Pre-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis

[] vancomycin (VANCOCIN) IVPB

15 mg/kg, intravenous, once, Starting S, For 1 Doses, Pre-op
Give within 2 hours of penile prosthesis procedure. Infuse
over 60 minutes. Repeat 8 hours after initial dose if still
intra-op.

Type of Therapy: New Anti-Infective Order
Reason for Therapy: Surgical Prophylaxis

Other Antibiotics: For Patients GREATER than 120 kg (Single Response)

() ceFAZolin (ANCEF) IV - For Patients GREATER than 120 kg	3 g, intravenous, once, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() cefoxitin (MEFOXIN) IV	2 g, intravenous, once, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() metronidazole (FLAGYL) IV	500 mg, intravenous, once, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() ampicillin-sulbactam (UNASYN) IV	3 g, intravenous, once, Starting S, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() levofloxacin (LEVAQUIN) IV	500 mg, intravenous, once, Starting S, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, PACU & Post-op Dose to be given 12 hours after pre-op dose. Use of vancomycin is indicated due to high prevalence rates for MRSA, for all areas within the hospital. Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() gentamicin (GARAMICIN) IVPB - For Penicillin Allergic Patients	5 mg/kg, intravenous, for 60 Minutes, once, Starting S, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis

Other Antibiotics: For Patients LESS than or EQUAL to 120 kg (Single Response)

() ceFAZolin (ANCEF) IV - For Patients LESS than or EQUAL to 120 kg	2 g, intravenous, once, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() cefoxitin (MEFOXIN) IV	2 g, intravenous, once, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() metronidazole (FLAGYL) IV	500 mg, intravenous, once, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() ampicillin-sulbactam (UNASYN) IV	3 g, intravenous, once, Starting S, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() levofloxacin (LEVAQUIN) IV	500 mg, intravenous, once, Starting S, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, PACU & Post-op Dose to be given 12 hours after pre-op dose. Use of vancomycin is indicated due to high prevalence rates for MRSA, for all areas within the hospital. Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis

() gentamicin (GARAMICIN) IVPB - For Penicillin Allergic Patients	5 mg/kg, intravenous, for 60 Minutes, once, Starting S, For 1 Doses, PACU & Post-op Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
--	--

Beta Blockers - Day of Surgery (Single Response)

If patient was on beta blocker therapy prior to the procedure:

1. The patient must have a dose of beta blocker on day of surgery (pre-op or post-op) OR a contraindication to beta blocker should be documented on the day of surgery (pre or post op)
2. A beta blocker should be resumed (via order or medication reconciliation) pos-op Day 1 OR a contraindication to beta blocker should be documented on POD 1 or POD 2.

() Reason for no beta-blocker in periop	Routine, Once For 1 Occurrences Reasons for not administering Beta-Blockers-Perioperatively: PACU
() metoprolol tartrate (LOPRESSOR) tablet	12.5 mg, oral, once, For 1 Doses, Post-op Hold for SBP less than 100 mmHg, DBP less than 60, HR less than 50 or patient is on a vasopressor or inotrope. HOLD parameters for this order: Contact Physician if:
() metoprolol (LOPRESSOR) injection	2.5 mg, intravenous, once, For 1 Doses, Post-op Hold for SBP less than 100 mmHg, DBP less than 60, HR less than 50 or patient is on a vasopressor or inotrope. HOLD parameters for this order: Contact Physician if:
() carvedilol (COREG) tablet	3.125 mg, oral, once, For 1 Doses, Post-op Hold for SBP less than 100 mmHg, DBP less than 60, HR less than 50 or patient is on a vasopressor or inotrope. HOLD parameters for this order: Contact Physician if:

Beta Blockers - Day After Surgery (Single Response)

If patient was on beta-blocker therapy prior to the procedure:

1. The patient must have a dose of beta blocker on day or surgery (Pre-Op or Post-Op) OR a contraindication to beta blocker should be documented on the day of surgery (Pre or Post Op).
2. A beta blocker should be resumed (via order or medication reconciliation) Post-Op Day 1 OR a contraindication to beta blocker should be documented on POD 1 or POD 2.

() metoprolol tartrate (LOPRESSOR) tablet	12.5 mg, oral, 2 times daily at 0600, 1800, Starting S+1, Post-op Hold for SBP less than 100 mmHg, DBP less than 60, HR less than 50 or patient is on a vasopressor or inotrope. HOLD parameters for this order: Contact Physician if:
() carvedilol (COREG) tablet	3.125 mg, oral, 2 times daily at 0600, 1800, Starting S+1, Post-op Hold for SBP less than 100 mmHg, DBP less than 60, HR less than 50 or patient is on a vasopressor or inotrope. HOLD parameters for this order: Contact Physician if:
() metoprolol (LOPRESSOR) injection	2.5 mg, intravenous, 2 times daily, Starting S+1, Post-op Hold for SBP less than 100 mmHg, DBP less than 60, HR less than 50 or patient is on a vasopressor or inotrope. HOLD parameters for this order: Contact Physician if:

Mild Pain (Pain Score 1-3)

[] acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	1 tablet, oral, every 4 hours PRN, mild pain (score 1-3), Post-op May administer when patient is passing flatus.
---	---

<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 4 hours PRN, mild pain (score 1-3), Post-op May administer when patient is passing flatus
<input type="checkbox"/> traMADol (ULTRAM) tablet	50 mg, oral, every 6 hours PRN, mild pain (score 1-3), Post-op

VTE

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

() Low Risk of DVT

Low Risk (Single Response)

() Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.

Will encourage early ambulation

PACU & Post-op

() Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical

Patient (Single Response)

() Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.

Therapy for the following:

PACU & Post-op

() Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

PACU & Post-op

() enoxaparin (LOVENOX) injection (Single Response)

() enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

() enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

() enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() Place sequential compression device and antiembolic stockings	"And" Linked Panel
[] Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
[] Place antiembolic stockings	Routine, Once, PACU & Post-op
() Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
[] Moderate Risk	
[] Moderate risk of VTE	Routine, Once, PACU & Post-op
[] Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
() Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
() Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() enoxaparin (LOVENOX) injection (Single Response)	
() enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
() enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCl LESS than 30 mL/min
() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

() enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):	
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op	
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.	
() warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:	
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:	
[] Mechanical Prophylaxis (Single Response)		
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op	
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op	
() Place sequential compression device and antiembolic stockings	"And" Linked Panel	
[] Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op	
[] Place antiembolic stockings	Routine, Once, PACU & Post-op	
() High Risk of DVT - Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
[] High Risk		
[] High risk of VTE	Routine, Once, PACU & Post-op	
[] High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	[] Patient is currently receiving therapeutic anticoagulation	
	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op	
() Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op	
() enoxaparin (LOVENOX) injection (Single Response)	[] enoxaparin (LOVENOX) syringe	
	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1	
() enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCl LESS than 30 mL/min	
() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	
() enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	

() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
() High Risk of DVT - Non-Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
() Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
() Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() enoxaparin (LOVENOX) injection (Single Response)	
() enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
() enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
() enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input checked="" type="checkbox"/> Mechanical Prophylaxis (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input checked="" type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input checked="" type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
() High Risk of DVT - Surgical (Hip/Knee)	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<input checked="" type="checkbox"/> High Risk	
<input checked="" type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input checked="" type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
() Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
() Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1, PACU & Post-op Indications:
() aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
() aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
() enoxaparin (LOVENOX) injection (Single Response)	
() enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
() enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
() enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCl LESS than 30 mL/min.
() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.

() enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1, PACU & Post-op To be Given on Post Op Day 1. Indications:
() warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() Place sequential compression device and antiembolic stockings	"And" Linked Panel
[] Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
[] Place antiembolic stockings	Routine, Once, PACU & Post-op

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

[] Low Risk (Single Response)

() Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.

Will encourage early ambulation

PACU & Post-op

() Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

[] Moderate Risk

[] Moderate risk of VTE

Routine, Once, PACU & Post-op

[] Moderate Risk Pharmacological Prophylaxis - Surgical

Patient (Single Response)

() Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.

Therapy for the following:

PACU & Post-op

() Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

PACU & Post-op

() enoxaparin (LOVENOX) injection (Single Response)

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

() enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCl LESS than 30 mL/min

() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

() enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min

() fondaparinux (ARIXTRA) injection

2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

() heparin (porcine) injection

5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op

() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)

5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op

Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

() warfarin (COUMADIN) tablet

oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op

Indication:

() Pharmacy consult to manage warfarin (COUMADIN)

STAT, Until discontinued, Starting S

Indication:

[] Mechanical Prophylaxis (Single Response)

() Contraindications exist for mechanical prophylaxis

Routine, Once

No mechanical VTE prophylaxis due to the following contraindication(s):

PACU & Post-op

() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() Place sequential compression device and antiembolic stockings	"And" Linked Panel
[] Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
[] Place antiembolic stockings	Routine, Once, PACU & Post-op
() Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
[] Moderate Risk	
[] Moderate risk of VTE	Routine, Once, PACU & Post-op
[] Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
() Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
() Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() enoxaparin (LOVENOX) injection (Single Response)	
() enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
() enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
() enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() Place sequential compression device and antiembolic stockings	"And" Linked Panel

<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
() High Risk of DVT - Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
() Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
() Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() enoxaparin (LOVENOX) injection (Single Response)	
() enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
() enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
() enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() Place sequential compression device and antiembolic stockings	"And" Linked Panel

<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
() High Risk of DVT - Non-Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
() Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
() Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() enoxaparin (LOVENOX) injection (Single Response)	
() enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
() enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
() enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Place antiembolic stockings	Routine, Once, PACU & Post-op
() High Risk of DVT - Surgical (Hip/Knee)	

<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
() Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
() Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1, PACU & Post-op Indications:
() aspirin chewable tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
() aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1, PACU & Post-op
() enoxaparin (LOVENOX) injection (Single Response)	
() enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
() enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
() enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
() enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
() enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
() fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
() heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
() heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
() rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1, PACU & Post-op To be Given on Post Op Day 1. Indications:
() warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1, PACU & Post-op Indication:
() Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	

() Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
() Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
() Place sequential compression device and antiembolic stockings	"And" Linked Panel
[] Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
[] Place antiembolic stockings	Routine, Once, PACU & Post-op

Labs

Laboratory

[] Potassium level - PACU	STAT For 1 Occurrences, PACU
[] Bedside glucose - PACU	STAT, Once, PACU
[] Hemoglobin	Once, PACU
[] Basic metabolic panel	Once, PACU

Cardiology

Imaging

Imaging

[] Upper GI	Routine, 1 time imaging For 1 , Post-op
--------------	---

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders