

General

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Details
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Details
<input type="checkbox"/>	Acute Renal Failure	Details
<input type="checkbox"/>	Acute Respiratory Failure	Details
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Details
<input type="checkbox"/>	Anemia	Details
<input type="checkbox"/>	Bacteremia	Details
<input type="checkbox"/>	Bipolar disorder, unspecified	Details
<input type="checkbox"/>	Cardiac Arrest	Details
<input type="checkbox"/>	Cardiac Dysrhythmia	Details
<input type="checkbox"/>	Cardiogenic Shock	Details
<input type="checkbox"/>	Decubitus Ulcer	Details
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Details
<input type="checkbox"/>	Disorder of Liver	Details
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Details
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Details
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Details
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Details
<input type="checkbox"/>	Other Alteration of Consciousness	Details
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Details
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Details
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Details
<input type="checkbox"/>	Protein-calorie Malnutrition	Details
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Details
<input type="checkbox"/>	Schizophrenia Disorder	Details
<input type="checkbox"/>	Sepsis	Details
<input type="checkbox"/>	Septic Shock	Details
<input type="checkbox"/>	Septicemia	Details
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Details
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Details

Admission or Observation (Single Response)

<input type="checkbox"/> Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
<input type="checkbox"/> Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
<input type="checkbox"/> Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

Admission or Observation (Single Response)

Patient has active status order on file

Admit to Inpatient
Diagnosis:
Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.

Outpatient observation services under general supervision
Diagnosis:
Admitting Physician:
Patient Condition:
Bed request comments:

Outpatient in a bed - extended recovery
Diagnosis:
Admitting Physician:
Bed request comments:

Admission (Single Response)

Patient has active status order on file.

Admit to inpatient
Diagnosis:
Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.

Code Status

Full code
Code Status decision reached by:

DNR

DNR (Do Not Resuscitate) Does patient have decision-making capacity?

Consult to Palliative Care Service
Priority:

Reason for Consult?

Order?

Name of referring provider:

Enter call back number:

Consult to Social Work
Reason for Consult:

Modified Code
Does patient have decision-making capacity?

Modified Code restrictions:

Treatment Restrictions
Treatment Restriction decision reached by:

Specify Treatment Restrictions:

Isolation

Airborne isolation status
Details

Contact isolation status
Details

Droplet isolation status
Details

Enteric isolation status
Details

Precautions

Aspiration precautions
Details

Fall precautions
Increased observation level needed:

Latex precautions
Details

Seizure precautions
Increased observation level needed:

Nursing

Vitals

Vital signs - T/P/R/BP
Routine, Every 6 hours

Activity

<input type="checkbox"/> Ambulate	Routine, 3 times daily Specify:
<input type="checkbox"/> Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated
<input type="checkbox"/> Strict bed rest	Routine, Until discontinued, Starting S

Nursing

<input checked="" type="checkbox"/> Height and weight	Routine, Daily With same scale and manner and record in EPIC
<input checked="" type="checkbox"/> Intake and output	Routine, Every shift
<input type="checkbox"/> Telemetry	"And" Linked Panel
<input type="checkbox"/> Telemetry monitoring	Routine, Continuous Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box) Reason for telemetry: Can be off of Telemetry for tests and baths? Yes Upon arrival to the unit
<input type="checkbox"/> Telemetry Additional Setup Information	Routine, Continuous High Heart Rate (BPM): 120 Low Heart Rate(BPM): 50 High PVC's (per minute): 10 High SBP(mmHg): 175 Low SBP(mmHg): 100 High DBP(mmHg): 95 Low DBP(mmHg): 40 Low Mean BP: 60 High Mean BP: 120 Low SPO2(%): 94
<input type="checkbox"/> Nursing communication	Ask patient if enrolled in research study, if so notify study coordinator
<input type="checkbox"/> Limb precautions- No Venipuncture or Blood Pressure to Hemodialysis Arm	Location: Precaution: No venipuncture, No blood pressure To Hemodialysis arm. Place sign at bedside.

Diet

<input type="checkbox"/> Diet - Post Transplant	Diet effective now, Starting S Diet(s): Post Transplant Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
<input type="checkbox"/> NPO Except meds	Diet effective now, Starting S NPO: Except meds Pre-Operative fasting options:

Notify

<input checked="" type="checkbox"/> Notify Transplant Surgeon of patient arrival	Routine, Until discontinued, Starting S
<input checked="" type="checkbox"/> Notify Nephrologist of patient arrival	Routine, Until discontinued, Starting S
<input checked="" type="checkbox"/> Notify Study Coordinator	Routine, Once Ask patient if enrolled in research study, if so notify study coordinator
<input checked="" type="checkbox"/> Notify Physician for blood glucose greater than 30 milligrams/deciliter	Routine, Until discontinued, Starting S, for blood glucose greater than 30 milligrams/deciliter
<input checked="" type="checkbox"/> Notify Physician: For urine output less than 200 milliliters in 8 hours	Routine, Until discontinued, Starting S, For urine output less than 200 milliliters in 8 hours

Notify Physician for vitals:

Routine, Until discontinued, Starting S
Temperature greater than: 100
Temperature less than:
Systolic BP greater than: 180
Systolic BP less than: 100
Diastolic BP greater than: 110
Diastolic BP less than: 60
MAP less than:
Heart rate greater than (BPM): 110
Heart rate less than (BPM): 60
Respiratory rate greater than: 25
Respiratory rate less than: 10
SpO2 less than:

IV Fluids

Peripheral IV Access

Initiate and maintain IV

Insert peripheral IV

Routine, Once

sodium chloride 0.9 % flush

10 mL, intravenous, every 12 hours scheduled

sodium chloride 0.9 % flush

10 mL, intravenous, PRN, line care

IV Fluids

sodium chloride 0.9 % bolus

1,000 mL, intravenous, for 60 Minutes, once, For 1 Doses

sodium chloride 0.9 % infusion

75 mL/hr, intravenous, continuous

sodium chloride 0.45 % 1,000 mL with sodium bicarbonate 75 mEq/L infusion

75 mL/hr, intravenous, continuous

Medications

Alternate Route Immunosuppression

IV - methylPREDNISolone sodium succinate (Solu-MEDROL) injection

intravenous, daily

If given by IV Push, administer over no less than 3 minutes.

IV - mycophenolate (CELLCEPT) in dextrose 5% IVPB

intravenous, for 2 Hours, 2 times daily at 0600, 1800

Sublingual - tacrolimus (PROGRAF) capsule

sublingual, 2 times daily at 0600, 1800 (time critical)

Sublingual - tacrolimus (PROGRAF) capsule - Select this option if patients doses are different for Morning and Evening doses

"And" Linked Panel

tacrolimus (PROGRAF) capsule

sublingual, daily at 0600 (time critical)

tacrolimus (PROGRAF) capsule

sublingual, daily at 1800

Stress Ulcer Prophylaxis

pantoprazole (PROTONIX) EC tablet

40 mg, oral, daily at 0600

Indication(s) for Proton Pump Inhibitor (PPI) Therapy:

famotidine (PEPCID) tablet

20 mg, oral, daily

Antiemetics

ondansetron (ZOFTRAN) IV or Oral

"Or" Linked Panel

ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet

4 mg, oral, every 8 hours PRN, nausea, vomiting
Give if patient is able to tolerate oral medication.

ondansetron (ZOFTRAN) 4 mg/2 mL injection

4 mg, intravenous, every 8 hours PRN, nausea, vomiting
Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.

promethazine (PHENERGAN) IV or Oral or Rectal

"Or" Linked Panel

promethazine (PHENERGAN) 12.5 mg IV

12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting
Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.

promethazine (PHENERGAN) tablet

12.5 mg, oral, every 6 hours PRN, nausea, vomiting
Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.

<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.
--	---

Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
<input type="checkbox"/> promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9 % 0.9 % 20 mL for Alaris pump syringe option	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IVPB or Oral or Rectal	"Or" Linked Panel
<input type="checkbox"/> promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Restricted Medications

<input checked="" type="checkbox"/> No NSAIDs EXcluding aspirin	STAT, Until discontinued, Starting S
---	--------------------------------------

Insomnia: Ramelteon For Patients GREATER than 70 years old

<input checked="" type="checkbox"/> ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep
--	--------------------------------

Insomnia: Zolpidem For Patients LESS than 70 years old (Single Response)

<input checked="" type="checkbox"/> zolpidem (AMBIEN) tablet	5 mg, oral, nightly PRN, sleep
--	--------------------------------

VTE

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min

fondaparinux (ARIXTRA) injection

2.5 mg, subcutaneous, daily, Starting S+1

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:

<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Non-Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee)	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

<input type="checkbox"/> Low Risk of DVT	
<input type="checkbox"/> Low Risk (Single Response)	
<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

<input type="checkbox"/> Moderate Risk of DVT - Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	

<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S+1 For Patients with CrCl LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:

<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> High Risk of DVT - Surgical Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> High Risk of DVT - Non-Surgical Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):

<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily, Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical) For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee)	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.
Will encourage early ambulation

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	

<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
<input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place sequential compression device and antiembolic stockings	"And" Linked Panel
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once

Labs

Admission Labs

<input type="checkbox"/> CBC with platelet and differential	STAT For 1 Occurrences
<input type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences
<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	STAT For 1 Occurrences
<input type="checkbox"/> Hepatic function panel	STAT For 1 Occurrences
<input type="checkbox"/> HLA antibody testing - pre transplant	STAT For 1 Occurrences
<input type="checkbox"/> Magnesium level	STAT For 1 Occurrences
<input type="checkbox"/> BK virus by PCR	STAT For 1 Occurrences Specimen Source: Plasma
<input type="checkbox"/> Cytomegalovirus by PCR	STAT For 1 Occurrences Specimen Source: Plasma
<input type="checkbox"/> Blood culture x 2	
<input type="checkbox"/> Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.
<input type="checkbox"/> Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.
<input type="checkbox"/> Blood culture, fungus	STAT For 1 Occurrences, Blood
<input type="checkbox"/> Urinalysis, automated with microscopy	STAT For 1 Occurrences

<input type="checkbox"/>	Urinalysis screen and microscopy, with reflex to culture	STAT For 1 Occurrences Specimen Source: Urine Specimen Site:
<input type="checkbox"/>	Sputum culture	STAT For 1 Occurrences, Sputum
<input type="checkbox"/>	Aerobic culture	STAT For 1 Occurrences, Wound
<input type="checkbox"/>	Respiratory pathogen panel	STAT For 1 Occurrences, Nasopharyngeal

Laboratory AM 5:00

<input type="checkbox"/>	CBC with platelet and differential	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Prothrombin time with INR	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Partial thromboplastin time	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Basic metabolic panel	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Magnesium level	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Phosphorus level	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Amylase	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Lipase	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	FK506 Tacrolimus level, random	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Cyclosporine level, random	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Sirolimus level, random	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Everolimus level, random	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Prealbumin	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Ferritin level	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Total iron binding capacity and % saturation	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Iron level	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Phosphorus level	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Insulin, fasting	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	C-peptide	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Creatinine level, urine, random	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM
<input type="checkbox"/>	Protein, urine, random	AM draw, Starting S+1 For 1 Occurrences Draw at 5 AM

Every morning times 3 days

<input checked="" type="checkbox"/>	CBC with platelet and differential	AM draw repeats For 3 Days
<input type="checkbox"/>	Basic metabolic panel	AM draw repeats For 3 Days
<input checked="" type="checkbox"/>	Magnesium level	AM draw repeats For 3 Days
<input checked="" type="checkbox"/>	Phosphorus level	AM draw repeats For 3 Days
<input type="checkbox"/>	Amylase	AM draw repeats For 3 Days
<input type="checkbox"/>	Lipase	AM draw repeats For 3 Days
<input type="checkbox"/>	Prothrombin time with INR	AM draw repeats For 3 Days
<input type="checkbox"/>	Partial thromboplastin time	AM draw repeats For 3 Days

<input type="checkbox"/>	FK506 Tacrolimus level, random	AM draw repeats For 3 Days
<input type="checkbox"/>	Cyclosporine level, random	AM draw repeats For 3 Days
<input type="checkbox"/>	Sirolimus level, random	AM draw repeats For 3 Days
<input type="checkbox"/>	Everolimus level, random	AM draw repeats For 3 Days

Cardiology

Imaging

Diagnostic CT

<input type="checkbox"/>	CT Abdomen Pelvis WO Contrast (Omnipaque)	"And" Linked Panel
	For those with iodine allergies, please order the panel with Readi-Cat (barium sulfate).	
<input type="checkbox"/>	CT Abdomen Pelvis Wo Contrast	STAT, 1 time imaging For 1 Occurrences Oral contrast only. With iohexol Oral
<input type="checkbox"/>	iohexol (OMNIPAQUE) 300 mg iodine/mL oral solution	30 mL, oral, once
<input type="checkbox"/>	CT Abdomen Pelvis WO Contrast (Readi-Cat)	"And" Linked Panel
	Ordered as secondary option for those with iodine allergies.	
<input type="checkbox"/>	CT Abdomen Pelvis Wo Contrast	STAT, 1 time imaging For 1 Occurrences Oral contrast only. With iohexol Oral
<input type="checkbox"/>	barium (READI-CAT 2) 2.1 % (w/v), 2.0 % (w/w) suspension	450 mL, oral, once in imaging, contrast, For 1 Doses

Diagnostic X-Ray

<input type="checkbox"/>	Chest 2 Vw W Apical Lordotic	STAT, 1 time imaging For 1 Occurrences
<input type="checkbox"/>	Abdomen 1 Vw	STAT, 1 time imaging For 1 Occurrences

Diagnostic US

<input type="checkbox"/>	US Renal Doppler	STAT, 1 time imaging For 1 Occurrences Upon arrival to Unit
<input type="checkbox"/>	US Pancreas	STAT, 1 time imaging For 1 Occurrences Pancreas with Flow - Upon arrival
<input type="checkbox"/>	US Needle Biopsy	STAT, 1 time imaging For 1 Occurrences Schedule for ultrasound-guided biopsy of kidney allograft

Other Studies

Respiratory

Respiratory Therapy

<input type="checkbox"/>	Oxygen therapy	Routine, Continuous Device 1: Nasal Cannula Rate in liters per minute: Rate in tenths of a liter per minute: O2 %: Titrate to keep O2 Sat Above: 90% Indications for O2 therapy: Titrate to maintain oxygen saturation greater than 90%; wean to room air
<input type="checkbox"/>	Encourage deep breathing and coughing	Routine, Every 2 hours while awake
<input type="checkbox"/>	Incentive spirometry	Routine, Every 2 hours while awake

Rehab

Consults

For Physician Consult orders use sidebar

Consults

<input type="checkbox"/> Consult to PT eval and treat	Special Instructions: Evaluate and treat for muscle strengthening and activity Weight Bearing Status:
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Other (Specify), Positive Nutrition Screen Specify: Nutrition assessment
<input type="checkbox"/> Consult Transplant Social Work	Reason for Consult? Transplant Psychosocial Evaluation Organ Transplant: Kidney 713-441-5451

Additional Orders