

General

Common Present on Admission Diagnosis

[] Acidosis	Details
[] Acute Post-Hemorrhagic Anemia	Details
[] Acute Renal Failure	Details
[] Acute Respiratory Failure	Details
[] Acute Thromboembolism of Deep Veins of Lower Extremities	Details
[] Anemia	Details
[] Bacteremia	Details
[] Bipolar disorder, unspecified	Details
[] Cardiac Arrest	Details
[] Cardiac Dysrhythmia	Details
[] Cardiogenic Shock	Details
[] Decubitus Ulcer	Details
[] Dementia in Conditions Classified Elsewhere	Details
[] Disorder of Liver	Details
[] Electrolyte and Fluid Disorder	Details
[] Intestinal Infection due to Clostridium Difficile	Details
[] Methicillin Resistant Staphylococcus Aureus Infection	Details
[] Obstructive Chronic Bronchitis with Exacerbation	Details
[] Other Alteration of Consciousness	Details
[] Other and Unspecified Coagulation Defects	Details
[] Other Pulmonary Embolism and Infarction	Details
[] Phlebitis and Thrombophlebitis	Details
[] Protein-calorie Malnutrition	Details
[] Psychosis, unspecified psychosis type	Details
[] Schizophrenia Disorder	Details
[] Sepsis	Details
[] Septic Shock	Details
[] Septicemia	Details
[] Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Details
[] Urinary Tract Infection, Site Not Specified	Details

Case Request

[] Case request operating room	Scheduling/ADT, Scheduling/ADT
---------------------------------	--------------------------------

Planned ICU Admission Post-Operatively (Admit to Inpatient Order) (Single Response)

Patients who are having an Inpatient Only Procedure as determined by CMS and patients with prior authorization for Inpatient Care may have an Admit to Inpatient order written pre-operatively.

() Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. Pre-op
------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Nursing

Vitals

[] Vital signs - T/P/R/BP	Routine, Per unit protocol On arrival and then routine, Pre-op
Activity	
[] Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated Pre-op
Nursing	
[] Height and weight	Routine, Once On Arrival and document, Pre-op
Consent	
[] Complete consent for	Routine, Once For 1 Occurrences Procedure: Orthotopic Cardiac Transplant Diagnosis/Condition: Physician: Pre-op
Notify	
[X] Notify Transplant Cardiology Service	Routine, Until discontinued, Starting S, Transplant Cardiology Service at 713-441-1100 upon patient arrival to unit, Pre-op
[X] Notify Transplant Coordinator On-Call	Routine, Until discontinued, Starting S, Transplant Coordinator On-Call at 713-441-5451 upon patient arrival to unit, Pre-op
[X] Notify - Consultants of patient's location	Routine, Until discontinued, Starting S, Consultants of patient's location, Pre-op
[X] Notify team if patient has taken Coumadin dose	Routine, Until discontinued, Starting S, Notify team if patient has taken Coumadin dose
Diet	
[X] NPO	Diet effective now, Starting S NPO: Except meds Pre-Operative fasting options: Give only specifically ordered medications, Pre-op
IV Fluids	
Medications	
Restricted Medications	
[X] No anti-platelet agents EXcluding aspirin	STAT, Until discontinued, Starting S Reason for "No" order: Post-op
[X] No ketorolac (Toradol)	STAT, Until discontinued, Starting S Reason for "No" order: Post-op
[X] No NSAIDs EXcluding aspirin	STAT, Until discontinued, Starting S, Post-op
Preoperative Antibiotics (Single Response)	
(X) Ceftriaxone + Vancomycin	
[X] cefTRIAxone (ROCEPHIN) IV	2 g, intravenous, for 30 Minutes, once, For 1 Doses, Pre-op To be dispensed in FBOR and administered by Anesthesia. To be given 1 hour PRIOR to skin incision. Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis

<input checked="" type="checkbox"/> vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Pre-op To be dispensed in FBOR and administered by Anesthesia. To be given 1 hour PRIOR to skin incision. Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis Indication: Increased MRSA rate (operation specific)
() Aztreonam + Vancomycin - For severe beta-lactam allergy	
<input type="checkbox"/> aztreonam (AZACTAM) IV	2 g, intravenous, once, For 1 Doses, Pre-op To be dispensed in FBOR and administered by Anesthesia. To be given 1 hour PRIOR to skin incision Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> vancomycin (VANCOCIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Pre-op To be dispensed in FBOR and administered by Anesthesia. To be given 1 hour PRIOR to skin incision Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis Indication: Increased MRSA rate (operation specific)
() For Vancomycin Allergy - ceFAZolin (ANCEF) IV - For Patients LESS than or EQUAL to 120 kg	
<input type="checkbox"/> ceFAZolin (ANCEF) IV	2 g, intravenous, once, For 1 Doses, Pre-op To be dispensed in FBOR and administered by Anesthesia. To be given 1 hour PRIOR to skin incision Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis
() For Vancomycin Allergy - ceFAZolin (ANCEF) IV - For Patients GREATER than 120 kg	
<input type="checkbox"/> ceFAZolin (ANCEF) IV	3 g, intravenous, once, For 1 Doses, Pre-op To be dispensed in FBOR and administered by Anesthesia. To be given 1 hour PRIOR to skin incision Type of Therapy: New Anti-Infective Order Reason for Therapy: Surgical Prophylaxis

Intraoperative Medications

<input checked="" type="checkbox"/> methylPREDNISolone (Solu-MEDROL) IVPB in 100 mL NS	1,000 mg, intravenous, for 15 Minutes Administer in OR by Anesthesiologist 15 minutes PRIOR to opening of aortic clamp.
<input type="checkbox"/> basiliximab (SIMULECT) injection	20 mg, intravenous, once, For 1 Doses, Pre-op

Labs

Laboratory STAT

<input checked="" type="checkbox"/> CBC with platelet and differential	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Comprehensive metabolic panel	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Hepatic function panel	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Magnesium level	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Phosphorus level	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> LDH	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Uric acid level	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Cholesterol	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Triglycerides	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Fibrinogen	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> Platelet function analysis	STAT For 1 Occurrences, Pre-op
<input checked="" type="checkbox"/> West Nile virus antibody IgM, serum	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> HLA antibody testing - pre transplant	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> HLA deceased donor	STAT For 1 Occurrences, Pre-op
<input type="checkbox"/> HLA stored sample - crossmatch	STAT For 1 Occurrences, Pre-op

[X] Urinalysis screen and microscopy, with reflex to culture	STAT For 1 Occurrences Specimen Source: Urine Specimen Site: Pre-op
[] hCG qualitative, urine screen	STAT For 1 Occurrences, Pre-op

Transmission Donor/Baseline Serology - HMH

[] HIV Ag/Ab combination	Once Perform on HBV negative recipient, Pre-op
[] HIV quantitative, PCR	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B surface antigen	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B core antibody, total	Once Perform on HBV negative recipient, Pre-op
[] HBV, quantitative PCR	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis C antibody	Once Perform regardless recipient HCV status, Pre-op
[] Hepatitis C quantitative, PCR	Once Perform regardless recipient HCV status, Pre-op

Transmission Donor/Baseline Serology - HMH

[] HIV Ag/Ab combination	Once Perform on HBV negative recipient, Pre-op
[] HIV-1 RNA, qualitative TMA	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B surface antigen	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B core antibody, total	Once Perform on HBV negative recipient, Pre-op
[] HBV, quantitative PCR	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis C antibody	Once Perform regardless recipient HCV status, Pre-op
[] Hepatitis C quantitative, PCR	Once Perform regardless recipient HCV status, Pre-op

Transmission Donor/Baseline Serology - HMSL/HMW

[] Rapid HIV 1 & 2	Once
[] HIV quantitative, PCR	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B surface antigen	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B core antibody, total	Once Perform on HBV negative recipient, Pre-op
[] HBV, quantitative PCR	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis C antibody	Once Perform regardless recipient HCV status, Pre-op
[] Hepatitis C quantitative, PCR	Once Perform regardless recipient HCV status, Pre-op

Transmission Donor/Baseline Serology - HMSL/HMW

[] Rapid HIV 1 & 2	Once
[] HIV-1 RNA, qualitative TMA	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B surface antigen	Once Perform on HBV negative recipient, Pre-op

[] Hepatitis B core antibody, total	Once Perform on HBV negative recipient, Pre-op
[] HBV, quantitative PCR	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis C antibody	Once Perform regardless recipient HCV status, Pre-op
[] Hepatitis C quantitative, PCR	Once Perform regardless recipient HCV status, Pre-op

Transmission Donor/Baseline Serology - HMSTJ, HMTW, HMSJ, HMWB

[] HIV 1, 2 antibody	Once Perform on HBV negative recipient, Pre-op
[] HIV quantitative, PCR	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B surface antigen	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B core antibody, total	Once Perform on HBV negative recipient, Pre-op
[] HBV, quantitative PCR	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis C antibody	Once Perform regardless recipient HCV status, Pre-op
[] Hepatitis C quantitative, PCR	Once Perform regardless recipient HCV status, Pre-op

Transmission Donor/Baseline Serology - HMSTJ, HMTW, HMSJ, HMWB

[] HIV 1, 2 antibody	Once Perform on HBV negative recipient, Pre-op
[] HIV-1 RNA, qualitative TMA	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B surface antigen	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis B core antibody, total	Once Perform on HBV negative recipient, Pre-op
[] HBV, quantitative PCR	Once Perform on HBV negative recipient, Pre-op
[] Hepatitis C antibody	Once Perform regardless recipient HCV status, Pre-op
[] Hepatitis C quantitative, PCR	Once Perform regardless recipient HCV status, Pre-op

Blood Products

[] Type and screen	Once, Pre-op
[] Blood Administration - Red Blood Cells	
[] Prepare RBC	Routine Transfusion Indications: Perioperative Special Requirements: Irradiated Transfusion date: This is a transplant patient. All blood products to be irradiated and leukocyte reduced If recipient is CMV Negative, then give only CMV Negative blood. Pre-op
[] Transfuse RBC	Routine Transfusion duration per unit (hrs): Pre-op
[] Adult Blood Administration - Platelets	

[] Prepare plateletpheresis

Routine
Transfusion Indications: Perioperative
Special Requirements: Irradiated
Transfusion date:
This is a transplant patient. All blood products to be
irradiated and leukocyte reduced. If recipient are CMV
Negative, then give only CMV Negative blood.
Pre-op

[] Transfuse platelets

Routine
Transfusion duration per unit (hrs):
Pre-op

Cardiology

Cardiology

[X] ECG 12 lead

STAT, Once
Clinical Indications: Other:
Other: Pre-operative
Interpreting Physician:
Pre-op

Imaging

Diagnostic X-Ray

[X] Chest 1 Vw Portable

STAT, 1 time imaging For 1
on arrival to unit, Pre-op

Other Studies

Respiratory

Respiratory

[] Oxygen therapy

Routine, Continuous
Device 1: Nasal Cannula
Rate in liters per minute:
Rate in tenths of a liter per minute:
O2 %:
Titrate to keep O2 Sat Above: 90%
Indications for O2 therapy:
Pre-op

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders