

General

Nursing

Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 15 min Every 15 min times 4, then every 30 min times 4, then every 1 hour times 2, then every 4 hours., PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 30 min Every 30 min times 4, then every 1 hour times two, then every 4 hours., PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every hour Every 1 hour times 2, then every 4 hours., PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 4 hours, PACU & Post-op

Activity

<input type="checkbox"/> Strict bed rest	Routine, Until discontinued, Starting S For 2 Hours Then resume pre-procedure activity, PACU & Post-op
--	---

Nursing

<input type="checkbox"/> Assess cath site	Routine, As needed Check catheter exit site for bleeding/hematoma with vital signs., PACU & Post-op
<input type="checkbox"/> Ok to use - dialysis access	Routine, Until discontinued, Starting S Device: Dialysis Access Enter site and laterality: _**_, PACU & Post-op
<input type="checkbox"/> Head of bed 45 degrees	Routine, Until discontinued, Starting S Head of bed: 45 degrees PACU & Post-op
<input type="checkbox"/> Bedside glucose	Routine, Once For 1 Occurrences Upon arrival to AOD/PACU. Notify Radiologist if glucose below 60 or above 200., PACU & Post-op
<input type="checkbox"/> Nursing communication	Routine, Until discontinued, Starting S Instruction: **, PACU & Post-op
<input type="checkbox"/> Discharge Home	Routine, Once For 1 Occurrences At **, if stable. Confirm medication reconciliation is complete., PACU

Diet

<input type="checkbox"/> NPO except meds	Diet effective now, Starting S For 1 Hours NPO: Except meds Pre-Operative fasting options: PACU & Post-op
<input type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op

Tube feeding

Diet effective now, Starting S
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Formula:
Tube Feeding Schedule:
Tube Feeding Schedule:
Dietitian to manage Tube Feed?
PACU & Post-op

Notify

Notify ordering physician if evidence of bleeding and/or hematoma around catheter exit site

Routine, Until discontinued, Starting S, Post-op

IV Fluids

Medications

Post Procedure Medications

fentaNYL (SUBLIMAZE) injection

25 mcg, intravenous, every 15 min PRN, severe pain (score 7-10), persistent pain, For 2 Doses, PACU & Post-op
Hold for respirations LESS than 12 breath/minute. For persistent pain, call interventional MD.

ondansetron (ZOFTRAN) IV

4 mg, intravenous, once PRN, nausea, vomiting, For 1 Doses, PACU & Post-op

VTE

Labs

Cardiology

Diagnostic Imaging

Other Diagnostic Studies

Respiratory

Rehab

Consults

Additional Orders