

## General

### Common Present on Admission Diagnosis

[ ] Acidosis	Details
[ ] Acute Post-Hemorrhagic Anemia	Details
[ ] Acute Renal Failure	Details
[ ] Acute Respiratory Failure	Details
[ ] Acute Thromboembolism of Deep Veins of Lower Extremities	Details
[ ] Anemia	Details
[ ] Bacteremia	Details
[ ] Bipolar disorder, unspecified	Details
[ ] Cardiac Arrest	Details
[ ] Cardiac Dysrhythmia	Details
[ ] Cardiogenic Shock	Details
[ ] Decubitus Ulcer	Details
[ ] Dementia in Conditions Classified Elsewhere	Details
[ ] Disorder of Liver	Details
[ ] Electrolyte and Fluid Disorder	Details
[ ] Intestinal Infection due to Clostridium Difficile	Details
[ ] Methicillin Resistant Staphylococcus Aureus Infection	Details
[ ] Obstructive Chronic Bronchitis with Exacerbation	Details
[ ] Other Alteration of Consciousness	Details
[ ] Other and Unspecified Coagulation Defects	Details
[ ] Other Pulmonary Embolism and Infarction	Details
[ ] Phlebitis and Thrombophlebitis	Details
[ ] Protein-calorie Malnutrition	Details
[ ] Psychosis, unspecified psychosis type	Details
[ ] Schizophrenia Disorder	Details
[ ] Sepsis	Details
[ ] Septic Shock	Details
[ ] Septicemia	Details
[ ] Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Details
[ ] Urinary Tract Infection, Site Not Specified	Details

### Admission or Observation (Single Response)

( ) Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
( ) Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
( ) Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

### Admission or Observation (Single Response)

Patient has active status order on file

( ) Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
( ) Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
( ) Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

#### Admission (Single Response)

Patient has active status order on file.

( ) Admit to inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
------------------------	--

#### Code Status

[ ] Full code	Code Status decision reached by:
[ ] DNR	
[ ] DNR (Do Not Resuscitate)	Does patient have decision-making capacity?
[ ] Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
[ ] Consult to Social Work	Reason for Consult:
[ ] Modified Code	Does patient have decision-making capacity? Modified Code restrictions:
[ ] Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions:

#### Isolation

[ ] Airborne isolation status	Details
[ ] Contact isolation status	Details
[ ] Droplet isolation status	Details
[ ] Enteric isolation status	Details

#### Precautions

[ ] Aspiration precautions	Details
[ ] Fall precautions	Increased observation level needed:
[ ] Latex precautions	Details
[ ] Seizure precautions	Increased observation level needed:

## Nursing

#### Vital Signs

[ ] Vital signs - Per Unit Protocol	Routine, Per unit protocol
Printed on 4/18/2019 at 2:06 PM from SUP	Page 2 of 21

<input type="checkbox"/> Vital signs - Every 4 hrs	Routine, Every 4 hours For Until specified
<input type="checkbox"/> Vital signs - Every shift	Routine, Every shift
<b>Activity</b>	
<input type="checkbox"/> Strict bed rest	Routine, Until discontinued, Starting S
<input type="checkbox"/> Bed rest with bathroom privileges	Routine, Until discontinued, Starting S Bathroom Privileges: with bathroom privileges
<input type="checkbox"/> Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated
<input type="checkbox"/> Ambulate with assistance	Routine, 3 times daily Specify: with assistance

<b>Nursing</b>	
<input type="checkbox"/> Intake and output	Routine, Every shift
<input type="checkbox"/> Nasogastric tube insert and maintain	
<input type="checkbox"/> Nasogastric tube insertion	Routine, Once Type:
<input type="checkbox"/> Nasogastric tube maintenance	Routine, Until discontinued, Starting S Tube Care Orders:
<input type="checkbox"/> Insert and maintain Foley	
<input type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed:
<input type="checkbox"/> Foley Catheter Care	Routine, Until discontinued, Starting S Orders: Maintain

<b>Notify Physician</b>	
<input type="checkbox"/> Notify Physician if patient reports alcohol use	Routine, Until discontinued, Starting S, If patient reports alcohol use
<input type="checkbox"/> Notify Physician for vitals:	Routine, Until discontinued, Starting S Temperature greater than: 100.5 Temperature less than: Systolic BP greater than: 160 Systolic BP less than: 90 Diastolic BP greater than: 100 Diastolic BP less than: 50 MAP less than: 60 Heart rate greater than (BPM): 100 Heart rate less than (BPM): 60 Respiratory rate greater than: 25 Respiratory rate less than: 8 SpO2 less than: 92

<b>Diet</b>	
<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options:
<input type="checkbox"/> Diet-Clear liquids	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Yes Target Diet: regular low fat Advance target diet criteria: Liquid Consistency: Fluid Restriction: Foods to Avoid:

## IV Fluids

<b>Peripheral IV Access</b>	
<input checked="" type="checkbox"/> Initiate and maintain IV	

<input checked="" type="checkbox"/> Insert peripheral IV	Routine, Once
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care

#### Bolus IV (Single Response)

( ) sodium chloride 0.9 % bolus	500 mL, intravenous, at 1000 mL/hr, once, For 1 Doses
( ) sodium chloride 0.9 % bolus	1,000 mL, intravenous, at 1000 mL/hr, once, For 1 Doses
( ) lactated Ringer's bolus	500 mL, intravenous, at 1000 mL/hr, once, For 1 Doses
( ) lactated Ringer's bolus	1,000 mL, intravenous, at 1000 mL/hr, once, For 1 Doses

#### Continuous IV (Single Response)

( ) dextrose 5 % and sodium chloride 0.45 % with potassium chloride 20 mEq/L infusion	100 mL/hr, intravenous, continuous
( ) dextrose 5%-0.45% sodium chloride 1,000 mL infusion	100 mL/hr, intravenous, continuous
( ) dextrose 5%-0.9% sodium chloride infusion	100 mL/hr, intravenous, continuous
( ) sodium chloride 0.9 % infusion	100 mL/hr, intravenous, continuous
( ) lactated Ringer's infusion	100 mL/hr, intravenous, continuous

## Medications

#### Pharmacy Consults

<input type="checkbox"/> Pharmacy to dose Tramadol for renal and hepatic dysfunction	Routine, Until discontinued, Starting S Which drug do you need help dosing? tramadol Adjust dose for: renal and hepatic dysfunction
--	---

## VTE

#### DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

( ) Low Risk of DVT

Low Risk (Single Response)

( ) Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.  
Will encourage early ambulation

( ) Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical

Patient (Single Response)

( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once
( ) Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
[ ] Moderate Risk	
[ ] Moderate risk of VTE	Routine, Once
[ ] Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
[ ] High Risk	
[ ] High risk of VTE	Routine, Once
[ ] High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min

( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<b>[ ] Mechanical Prophylaxis (Single Response)</b>	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Non-Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<b>[ ] High Risk</b>	
[ ] High risk of VTE	Routine, Once
[ ] High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<b>( ) enoxaparin (LOVENOX) injection (Single Response)</b>	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Surgical (Hip/Knee)	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCl LESS than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.

( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once

#### DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

( ) Low Risk of DVT

[ ] Low Risk (Single Response)

( ) Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
( ) Moderate Risk of DVT - Surgical	Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCl LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
( ) Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

( ) High Risk of DVT - Surgical

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

[ ] High Risk	
[ ] High risk of VTE	Routine, Once
[ ] High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
( ) High Risk of DVT - Non-Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily, Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily, Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical) For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
( ) High Risk of DVT - Surgical (Hip/Knee)	

<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

## DVT Risk and Prophylaxis Tool (Single Response)

## Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

### ( ) Low Risk of DVT

#### [ ] Low Risk (Single Response)

( ) Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.  
Will encourage early ambulation

### ( ) Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

#### [ ] Moderate Risk

[ ] Moderate risk of VTE

Routine, Once

#### [ ] Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

( ) Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.  
Therapy for the following:

( ) Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

( ) enoxaparin (LOVENOX) injection (Single Response)

( ) enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min

( ) fondaparinux (ARIXTRA) injection

2.5 mg, subcutaneous, daily, Starting S+1

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Place antiembolic stockings	Routine, Once
( ) Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCl LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:

( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
[ ] High Risk	
[ ] High risk of VTE	Routine, Once
[ ] High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response)	

( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Non-Surgical	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
[ ] High Risk	
[ ] High risk of VTE	Routine, Once
[ ] High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous

<input type="checkbox"/> Place antiembolic stockings	Routine, Once
( ) High Risk of DVT - Surgical (Hip/Knee)	Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.
<b>[ ] High Risk</b>	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
( ) Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
( ) enoxaparin (LOVENOX) injection (Single Response)	
( ) enoxaparin (LOVENOX) syringe - hip arthroplasty	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - knee arthroplasty	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min - knee/hip arthroplasty	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<b>[ ] Mechanical Prophylaxis (Single Response)</b>	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous

( ) Place sequential compression device and antiembolic stockings	<b>"And" Linked Panel</b>
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
[ ] Place antiembolic stockings	Routine, Once

## Labs

### Lab Today

[X] CBC and differential	Once
[ ] Basic metabolic panel	Once
[X] Comprehensive metabolic panel	Once
[ ] Hepatic function panel	Once
[ ] APTT	Once
[ ] Protime-INR	Once
[ ] Calcium	Once
[ ] Cholesterol, total	Once
[ ] Triglycerides	Once
[ ] Lactate dehydrogenase	Once
[X] Amylase	Once
[X] Lipase	Once
[ ] C-reactive protein	Once
[ ] Alcohol	Once
[ ] MWS - Urinalysis, auto. with microscopy	Once
[ ] Toxicology screen, urine	Once

### Labs Tomorrow

[ ] CBC and differential	AM draw For 1 Occurrences
[ ] Protime-INR	AM draw For 1 Occurrences
[ ] APTT	AM draw For 1 Occurrences
[ ] Basic metabolic panel	AM draw For 1 Occurrences
[ ] Comprehensive metabolic panel	AM draw For 1 Occurrences
[ ] Amylase	AM draw For 1 Occurrences
[ ] Lipase	AM draw For 1 Occurrences
[ ] Lactate dehydrogenase	AM draw For 1 Occurrences

### Microbiology

[ ] Blood culture x 2	<b>"And" Linked Panel</b>
[ ] Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.
[ ] Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.

## Cardiology

### Diagnostic Imaging

#### X-RAY

[ ] Abdomen Ap Lat Both Obliques	Routine, 1 time imaging For 1
[ ] Chest 2 Vw	Routine, 1 time imaging For 1
[ ] Chest 1 Vw Portable	Routine, 1 time imaging For 1

## US

<input type="checkbox"/> US Abdomen Complete	Routine, 1 time imaging For 1
--	-------------------------------

## CT

<input type="checkbox"/> CT Abdomen W Contrast (Omnipaque)	<b>"And" Linked Panel</b>
--	---------------------------

For those with iodine allergies, please order the panel with Readi-Cat (barium sulfate).

<input type="checkbox"/> CT Abdomen W Contrast	Routine, 1 time imaging For 1
--	-------------------------------

<input type="checkbox"/> iohexol (OMNIPACQUE) 300 mg iodine/mL oral solution	30 mL, oral, once
--	-------------------

<input type="checkbox"/> CT Abdomen WO Contrast (Omnipaque)	<b>"And" Linked Panel</b>
---	---------------------------

For those with iodine allergies, please order the panel with Readi-Cat (barium sulfate).

<input type="checkbox"/> CT Abdomen Wo Contrast	Routine, 1 time imaging For 1
---	-------------------------------

<input type="checkbox"/> iohexol (OMNIPACQUE) 300 mg iodine/mL oral solution	30 mL, oral, once
--	-------------------

<input type="checkbox"/> CT Abdomen WO Contrast (Readi-Cat)	<b>"And" Linked Panel</b>
---	---------------------------

Ordered as secondary option for those with iodine allergies.

<input type="checkbox"/> CT Abdomen Wo Contrast	Routine, 1 time imaging For 1
---	-------------------------------

<input type="checkbox"/> barium (READI-CAT 2) 2.1 % (w/v), 2.0 % (w/w) suspension	450 mL, oral, once in imaging, contrast, For 1 Doses
---	--

<input type="checkbox"/> CT Abdomen W/WO Contrast (Omnipaque)	<b>"And" Linked Panel</b>
---	---------------------------

For those with iodine allergies, please order the panel with Readi-Cat (barium sulfate).

<input type="checkbox"/> CT Abdomen W Wo Contrast	Routine, 1 time imaging For 1
---	-------------------------------

<input type="checkbox"/> iohexol (OMNIPACQUE) 300 mg iodine/mL oral solution	30 mL, oral, once
--	-------------------

## MRI

<input type="checkbox"/> MRI Abdomen W Wo Contrast	Routine, 1 time imaging For 1
--	-------------------------------

<input type="checkbox"/> MRI Cholangiogram	Routine, 1 time imaging For 1
--	-------------------------------

## Other Diagnostic Studies

### Respiratory

#### Respiratory

<input type="checkbox"/> Oxygen therapy - NC 2 Lpm	Routine, Continuous Device 1: Nasal Cannula Rate in liters per minute: 2 Lpm Rate in tenths of a liter per minute: O2 %: Titrate to keep O2 Sat Above: 92% Indications for O2 therapy: Device 2: Device 3: Indications for O2 therapy:
--	---

### Rehab

### Consults

#### Physician Consults

<input type="checkbox"/> Consult Gastroenterology	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated?
<input type="checkbox"/> Consult General Surgery	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated?
<input type="checkbox"/> Consult Pulmonary	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated?

<input type="checkbox"/> Consult Nephrology/Hyperten	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated?
<input type="checkbox"/> Consult Infectious Diseases	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated?

#### Ancillary Consults

<input type="checkbox"/> Consult to Case Management	Consult Reason:
<input type="checkbox"/> Consult to Social Work	Reason for Consult:
<input type="checkbox"/> Consult PT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult PT wound care	Special Instructions: Location of Wound?
<input type="checkbox"/> Consult OT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Purpose/Topic:
<input type="checkbox"/> Consult to Spiritual Care	Reason for consult?
<input type="checkbox"/> Consult to Speech Language Pathology	Routine, Once Reason for consult:
<input type="checkbox"/> Consult to Wound Ostomy Care nurse	Reason for consult: Reason for consult: Reason for consult: Reason for consult: Consult for NPWT: Reason for consult:
<input type="checkbox"/> Consult to Respiratory Therapy	Reason for Consult?

#### Additional Orders