

## General

## Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Details
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Details
<input type="checkbox"/>	Acute Renal Failure	Details
<input type="checkbox"/>	Acute Respiratory Failure	Details
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Details
<input type="checkbox"/>	Anemia	Details
<input type="checkbox"/>	Bacteremia	Details
<input type="checkbox"/>	Bipolar disorder, unspecified	Details
<input type="checkbox"/>	Cardiac Arrest	Details
<input type="checkbox"/>	Cardiac Dysrhythmia	Details
<input type="checkbox"/>	Cardiogenic Shock	Details
<input type="checkbox"/>	Decubitus Ulcer	Details
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Details
<input type="checkbox"/>	Disorder of Liver	Details
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Details
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Details
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Details
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Details
<input type="checkbox"/>	Other Alteration of Consciousness	Details
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Details
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Details
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Details
<input type="checkbox"/>	Protein-calorie Malnutrition	Details
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Details
<input type="checkbox"/>	Schizophrenia Disorder	Details
<input type="checkbox"/>	Sepsis	Details
<input type="checkbox"/>	Septic Shock	Details
<input type="checkbox"/>	Septicemia	Details
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Details
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Details

## Admission or Observation (Single Response)

<input type="checkbox"/> Admit to inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
---	--

<p>( ) Admit to IP- University Teaching Service</p>	<p>Diagnosis:          Admitting Physician:          Resident Physician:          Resident team assignment:          Level of Care:          Patient Condition:          Bed request comments:          Certification: I certify that based on my best clinical judgement and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.          To reach the team taking care of this patient please call the University Teaching Service Answering Service at (713) 363-9648 and ask for the team taking care of the patient to be paged. The team name is listed in both "Treatment Teams" and "Notes from Clinical Staff" sections in the Summary\Overview tab of Epic.</p>
<p>( ) Outpatient observation services under general supervision</p>	<p>Diagnosis:          Admitting Physician:          Patient Condition:          Bed request comments:</p>
<p>( ) UTS - Outpatient observation services under general supervision</p>	<p>Diagnosis:          Admitting Physician:          Resident Physician:          Resident team assignment:          Patient Condition:          Bed request comments:          To reach the team taking care of this patient please call the University Teaching Service Answering Service at (713) 363-9648 and ask for the team taking care of the patient to be paged. The team name is listed in both "Treatment Teams" and "Notes from Clinical Staff" sections in the Summary\Overview tab of Epic.</p>
<p>( ) Outpatient in a bed - extended recovery</p>	<p>Diagnosis:          Admitting Physician:          Bed request comments:</p>

**Admission or Observation (Single Response)**  
 Patient has active status order on file

<p>( ) Admit to inpatient</p>	<p>Diagnosis:          Admitting Physician:          Level of Care:          Patient Condition:          Bed request comments:          Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.</p>
-------------------------------	---

<input type="checkbox"/> Admit to IP- University Teaching Service	Diagnosis: Admitting Physician: Resident Physician: Resident team assignment: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgement and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. To reach the team taking care of this patient please call the University Teaching Service Answering Service at (713) 363-9648 and ask for the team taking care of the patient to be paged. The team name is listed in both "Treatment Teams" and "Notes from Clinical Staff" sections in the Summary\Overview tab of Epic.
<input type="checkbox"/> Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
<input type="checkbox"/> UTS - Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Resident Physician: Resident team assignment: Patient Condition: Bed request comments: To reach the team taking care of this patient please call the University Teaching Service Answering Service at (713) 363-9648 and ask for the team taking care of the patient to be paged. The team name is listed in both "Treatment Teams" and "Notes from Clinical Staff" sections in the Summary\Overview tab of Epic.
<input type="checkbox"/> Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

**Admission (Single Response)**

Patient has active status order on file.

<input type="checkbox"/> Admit to inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
---	--

**Admission or Observation (Single Response)**

<input type="checkbox"/> Admit to inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
---	--

<input type="checkbox"/> Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
<input type="checkbox"/> Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

**Admission or Observation (Single Response)**  
Patient has status order on file

<input type="checkbox"/> Admit to inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
<input type="checkbox"/> Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
<input type="checkbox"/> Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

**Code Status**

<input type="checkbox"/> Full code	Code Status decision reached by:
<input type="checkbox"/> DNR	
<input type="checkbox"/> DNR (Do Not Resuscitate)	Does patient have decision-making capacity?
<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult:
<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions:
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions:

**Isolation**

<input type="checkbox"/> Airborne isolation status	
<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.	Once, Sputum
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

**Precautions**

<input type="checkbox"/> Aspiration precautions	Details
<input type="checkbox"/> Fall precautions	Increased observation level needed:
<input type="checkbox"/> Latex precautions	Details
<input type="checkbox"/> Seizure precautions	Increased observation level needed:

**Nursing**

**Vital signs**

<input checked="" type="checkbox"/> Vital signs - T/P/R/BP (per unit protocol)	Routine, Per unit protocol
--	----------------------------

### Telemetry Order

<input type="checkbox"/> Telemetry	<b>"And" Linked Panel</b>
<input type="checkbox"/> Telemetry monitoring	Routine, Continuous Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box) Reason for telemetry: Chest pain syndrome Can be off of Telemetry for tests and baths? Yes
<input type="checkbox"/> Telemetry Additional Setup Information	Routine, Continuous High Heart Rate (BPM): 120 Low Heart Rate(BPM): 50 High PVC's (per minute): 10 High SBP(mmHg): 175 Low SBP(mmHg): 100 High DBP(mmHg): 95 Low DBP(mmHg): 40 Low Mean BP: 60 High Mean BP: 120 Low SPO2(%): 94

### Activity

<input type="checkbox"/> Up ad lib	Routine, Until discontinued, Starting S Specify: Up ad lib
<input type="checkbox"/> Strict bed rest	Routine, Until discontinued, Starting S
<input type="checkbox"/> Bed rest with bathroom privileges	Routine, Until discontinued, Starting S Bathroom Privileges: with bathroom privileges

### Nursing Care

<input type="checkbox"/> Daily weights	Routine, Daily
<input type="checkbox"/> Intake and Output Qshift	Routine, Every shift
<input type="checkbox"/> Nasogastric Tube Insert and Maintain	
<input type="checkbox"/> Nasogastric tube insertion	Routine, Once Type:
<input type="checkbox"/> Nasogastric tube maintenance	Routine, Until discontinued, Starting S Tube Care Orders:
<input type="checkbox"/> Insert and Maintain Foley	
<input type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed:
<input type="checkbox"/> Foley Catheter Care	Routine, Until discontinued, Starting S Orders: Maintain

### Notify Physician

<input checked="" type="checkbox"/> Notify Physician(vitals,output,pulse ox)	Routine, Until discontinued, Starting S Temperature greater than: 100.5 Temperature less than: Systolic BP greater than: 160 Systolic BP less than: 90 Diastolic BP greater than: 100 Diastolic BP less than: 50 Heart rate greater than (BPM): 100 Heart rate less than (BPM): 60 Respiratory rate greater than: 25 Respiratory rate less than: 8 SpO2 less than: 92 Urine Output less than: Output (Specify) greater than: Other:
--	---

**Notify Physician- UTS**

Notify Physician- Teaching Service

Routine, Until discontinued, Starting S, To reach the team taking care of this patient please call the University Teaching Service Answering Service at (713) 363-9648 and ask for the team taking care of the patient to be paged.

The team name is listed in both "Treatment Teams" and "Notes from Clinical Staff" sections in the Summary\Overview tab of Epic. If no response, page the Sr. Resident at 713-768-0403. If no response is obtained using second pager, page the attending assigned to the patient.

**Diet (Single Response)**

NPO

Diet effective now, Starting S  
NPO:  
Pre-Operative fasting options:

Diet

Diet effective now, Starting S  
Diet(s):  
Advance Diet as Tolerated?  
Liquid Consistency:  
Fluid Restriction:  
Foods to Avoid:

**Tube Feed**

Tube feeding - Continuous

Continuous  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Schedule: Continuous  
Tube Feeding Route:  
Initial Tube Feed rate (mL/hr):  
Advance Rate by (mL/hr):  
Goal Tube Feed Rate (mL/hr):  
Dietitian to manage Tube Feed?

Tube feeding - Bolus

Diet effective now, Starting S  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Schedule: Bolus  
Bolus Route:  
Tube Feeding Bolus (mL):  
Additional Bolus Schedule Instructions:  
Dietitian to manage Tube Feed?

Tube feeding - Cyclic

Cyclic  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Formula:  
Tube Feeding Schedule: Cyclic  
Tube Feeding Route:  
Tube Feeding Cyclic (start / stop time):  
Tube Feeding Cyclic Rate (mL/hr):  
Dietitian to manage Tube Feed?

## IV Fluids

### Peripheral IV Access

Initiate and maintain IV

Insert peripheral IV

Routine, Once

sodium chloride 0.9 % flush

10 mL, intravenous, every 12 hours scheduled

sodium chloride 0.9 % flush

10 mL, intravenous, PRN, line care

### IV Bolus (Single Response)

sodium chloride 0.9 % bolus 500 mL

500 mL, intravenous, for 15 Minutes, once, For 1 Doses

sodium chloride 0.9 % bolus 1000 mL

1,000 mL, intravenous, for 30 Minutes, once, For 1 Doses

lactated ringer's bolus 500 mL

500 mL, intravenous, for 15 Minutes, once, For 1 Doses

lactated ringers bolus 1000 mL

1,000 mL, intravenous, for 30 Minutes, once, For 1 Doses

### Maintenance IV Fluids (Single Response)

sodium chloride 0.9 % infusion

75 mL/hr, intravenous, continuous

lactated Ringer's infusion

75 mL/hr, intravenous, continuous

dextrose 5 % and sodium chloride 0.45 % with potassium chloride 20 mEq/L infusion

75 mL/hr, intravenous, continuous

sodium chloride 0.45 % infusion

75 mL/hr, intravenous, continuous

sodium chloride 0.45 % 1,000 mL with sodium bicarbonate 75 mEq/L infusion

75 mL/hr, intravenous, continuous

## Medications

For Analgesics, please refer to the General Pain Management order sets.

For Antihypertensives, please refer to the Hypertensive Urgency order set.

### Antibiotics

azithromycin (ZITHROMAX) IV

intravenous, for 60 Minutes

Reason for Therapy:

azithromycin (ZITHROMAX) tablet

oral, daily

Reason for Therapy:

cefepime (MAXIPIME) IV

intravenous

Reason for Therapy:

ceftriaxone (ROCEPHIN) IV

intravenous, for 30 Minutes

Reason for Therapy:

ciprofloxacin (CIPRO) IV

intravenous, for 60 Minutes

Reason for Therapy:

ciprofloxacin (CIPRO) tablet

100 mg, oral, 2 times daily

levofloxacin (LEVAQUIN) IV

intravenous

Reason for Therapy:

levofloxacin (LEVAQUIN) tablet

oral, daily at 0600 (time critical)

Reason for Therapy:

meropenem (MERREM) IV

intravenous

Reason for Therapy:

<input type="checkbox"/>	metroNIDAZOLE (FLAGYL) IV	intravenous Reason for Therapy:
<input type="checkbox"/>	metroNIDAZOLE (FLAGYL) tablet	500 mg, oral, 3 times daily Reason for Therapy:
<input type="checkbox"/>	piperacillin-tazobactam (ZOSYN) IV	intravenous Reason for Therapy:
<input type="checkbox"/>	vancomycin (VANCOCIN) IV	intravenous Type of Therapy:

### Shortness of Breath

<input type="checkbox"/>	albuterol (PROVENTIL) nebulizer solution	2.5 mg, nebulization, Respiratory Therapy - every 4 hours Aerosol Delivery Device:
<input type="checkbox"/>	albuterol (PROVENTIL) nebulizer solution	2.5 mg, nebulization, every 4 hours PRN, shortness of breath Aerosol Delivery Device:
<input type="checkbox"/>	ipratropium (ATROVENT) 0.02 % nebulizer solution	0.5 mg, nebulization, Respiratory Therapy - every 4 hours Aerosol Delivery Device:
<input type="checkbox"/>	ipratropium (ATROVENT) 0.02 % nebulizer solution	0.5 mg, nebulization, every 4 hours PRN, shortness of breath Aerosol Delivery Device:

### Cough (Single Response)

<input type="checkbox"/>	guaifenesin (MUCINEX) 12 hr tablet	600 mg, oral, 2 times daily PRN, cough
<input type="checkbox"/>	dextromethorphan-guaifenesin (MUCINEX DM REGULAR) 30-600 mg per 12 hr tablet	1 tablet, oral, every 12 hours PRN, cough
<input type="checkbox"/>	guaifenesin (ROBITUSSIN) 100 mg/5 mL syrup	100 mg, oral, every 4 hours PRN, cough
<input type="checkbox"/>	dextromethorphan-guaifenesin (ROBITUSSIN-DM) 10-100 mg/5 mL liquid	5 mL, oral, every 4 hours PRN, cough
<input type="checkbox"/>	codeine-guaifenesin (GUAIFENESIN AC) 10-100 mg/5 mL liquid	5 mL, oral, every 4 hours PRN, cough
<input type="checkbox"/>	benzonatate (TESSALON) capsule	100 mg, oral, every 6 hours PRN, cough

### Constipation - NOT HMSJ

<input type="checkbox"/>	docusate sodium (COLACE) capsule	100 mg, oral, 2 times daily PRN, constipation
<input type="checkbox"/>	senna (SENOKOT) tablet	1 tablet, oral, 2 times daily PRN, constipation, stool softening
<input type="checkbox"/>	sennosides-docusate sodium (SENOKOT-S) 8.6-50 mg per tablet	1 tablet, oral, daily PRN, constipation
<input type="checkbox"/>	polyethylene glycol (MIRALAX) packet	17 g, oral, daily PRN, constipation
<input type="checkbox"/>	magnesium hydroxide suspension	30 mL, oral, daily PRN, constipation
<input type="checkbox"/>	psyllium (METAMUCIL) 3.4 gram packet	1 packet, oral, daily PRN, constipation
<input type="checkbox"/>	bisacodyl (DULCOLAX) suppository	10 mg, rectal, daily PRN, constipation

### Constipation - HMSJ Only

<input type="checkbox"/>	docusate sodium (COLACE) capsule	100 mg, oral, 2 times daily PRN, constipation
<input type="checkbox"/>	sennosides-docusate sodium (SENOKOT-S) 8.6-50 mg per tablet	1 tablet, oral, daily PRN, constipation
<input type="checkbox"/>	polyethylene glycol (MIRALAX) packet	17 g, oral, daily PRN, constipation
<input type="checkbox"/>	magnesium hydroxide suspension	30 mL, oral, daily PRN, constipation
<input type="checkbox"/>	psyllium (METAMUCIL) 3.4 gram packet	1 packet, oral, daily PRN, constipation
<input type="checkbox"/>	bisacodyl (DULCOLAX) suppository	10 mg, rectal, daily PRN, constipation

### Insomnia: For Patients LESS than 70 years old (Single Response)

<input type="checkbox"/>	zolpidem (AMBIEN) tablet	5 mg, oral, nightly PRN, sleep
<input type="checkbox"/>	ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep

### Insomnia: For Patients GREATER than or EQUAL to 70 years old (Single Response)

<input type="checkbox"/>	ramelteon (ROZEREM) tablet	8 mg, oral, nightly PRN, sleep
--------------------------	----------------------------	--------------------------------

### Antiemetics

<input checked="" type="checkbox"/>	ondansetron (ZOFRAN) IV or Oral	"Or" Linked Panel
-------------------------------------	---------------------------------	-------------------



<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting Give if patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
<input checked="" type="checkbox"/> promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input checked="" type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

#### Itching: For Patients GREATER than 77 years old (Single Response)

<input type="checkbox"/> cetirizine (ZyrTEC) tablet	5 mg, oral, daily PRN, itching
---	--------------------------------

#### Itching: For Patients LESS than 70 years old (Single Response)

<input type="checkbox"/> diphenhydramINE (BENADRYL) tablet	25 mg, oral, every 6 hours PRN, itching
<input type="checkbox"/> hydrOXYzine (ATARAX) tablet	10 mg, oral, every 6 hours PRN, itching
<input type="checkbox"/> cetirizine (ZyrTEC) tablet	5 mg, oral, daily PRN, itching
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet - For eGFR LESS than 80 mL/min, reduce frequency to once daily as needed	60 mg, oral, 2 times daily PRN, itching

#### GI Drugs

<input type="checkbox"/> famotidine (PEPCID) IV or ORAL	<b>"Or" Linked Panel</b>
<input type="checkbox"/> famotidine (PEPCID) injection	20 mg, intravenous, 2 times daily IV or ORAL
<input type="checkbox"/> famotidine (PEPCID) tablet	20 mg, oral, 2 times daily IV or ORAL
<input type="checkbox"/> pantoprazole (PROTONIX) IV or ORAL	<b>"Or" Linked Panel</b>
<input type="checkbox"/> pantoprazole (PROTONIX) EC tablet	40 mg, oral, daily at 0600 Indication(s) for Proton Pump Inhibitor (PPI) Therapy:
<input type="checkbox"/> pantoprazole (PROTONIX) 40 mg in sodium chloride 0.9 % 10 mL injection	40 mg, intravenous, daily at 0600 Indication(s) for Proton Pump Inhibitor (PPI) Therapy:
<input type="checkbox"/> omeprazole (PriLOSEC) suspension	oral, daily Indication(s) for Proton Pump Inhibitor (PPI) Therapy:
<input type="checkbox"/> sucralfate (CARAFATE) TABLET or SUSPENSION	<b>"Or" Linked Panel</b>
<input type="checkbox"/> sucralfate (CARAFATE) tablet - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR GREATER	1 g, oral, 4 times daily with meals and nightly
<input type="checkbox"/> sucralfate (CARAFATE) 100 mg/mL suspension - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR GREATER	1 g, Nasogastric, 4 times daily with meals and nightly Use with Nasogastric tubing. Use if patient is unable to swallow tablet.
<input type="checkbox"/> alum-mag hydroxide-simeth (MAALOX) 200-200-20 mg/5 mL suspension - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR GREATER	30 mL, oral, every 4 hours PRN, indigestion Do not give if patient is on hemodialysis or in chronic renal failure.
<input type="checkbox"/> simethicone (MYLICON) chewable tablet	80 mg, oral, every 4 hours PRN, flatulence, For 2 Doses

#### GI Drugs

<input type="checkbox"/> famotidine (PEPCID) IV or ORAL	<b>"Or" Linked Panel</b>
<input type="checkbox"/> famotidine (PEPCID) injection	20 mg, intravenous, 2 times daily IV or ORAL
<input type="checkbox"/> famotidine (PEPCID) tablet	20 mg, oral, 2 times daily IV or ORAL
<input type="checkbox"/> pantoprazole (PROTONIX) IV or ORAL	<b>"Or" Linked Panel</b>

<input type="checkbox"/>	pantoprazole (PROTONIX) EC tablet	40 mg, oral, daily at 0600 Indication(s) for Proton Pump Inhibitor (PPI) Therapy:
<input type="checkbox"/>	pantoprazole (PROTONIX) 40 mg in sodium chloride 0.9 % 10 mL injection	40 mg, intravenous, daily at 0600 Indication(s) for Proton Pump Inhibitor (PPI) Therapy:
<input type="checkbox"/>	sucralfate (CARAFATE) TABLET or SUSPENSION	<b>"Or" Linked Panel</b>
<input type="checkbox"/>	sucralfate (CARAFATE) tablet - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR GREATER	1 g, oral, 4 times daily with meals and nightly
<input type="checkbox"/>	sucralfate (CARAFATE) 100 mg/mL suspension - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR GREATER	1 g, Nasogastric, 4 times daily with meals and nightly Use with Nasogastric tubing. Use if patient is unable to swallow tablet.
<input type="checkbox"/>	alum-mag hydroxide-simeth (MAALOX) 200-200-20 mg/5 mL suspension - NOT RECOMMENDED FOR CHRONIC KIDNEY DISEASE STAGE 3 OR GREATER	30 mL, oral, every 4 hours PRN, indigestion Do not give if patient is on hemodialysis or in chronic renal failure.
<input type="checkbox"/>	simethicone (MYLICON) chewable tablet	80 mg, oral, every 4 hours PRN, flatulence, For 2 Doses

## VTE

### DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.

Will encourage early ambulation

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM

<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

High Risk of DVT - Surgical (Hip/Knee)

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

High Risk

<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

### DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

#### Low Risk of DVT

##### Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.  
Will encourage early ambulation

#### Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

##### Moderate Risk

Moderate risk of VTE

Routine, Once

##### Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.  
Therapy for the following:

Contraindications exist for pharmacologic prophylaxis

Routine, Once

No pharmacologic VTE prophylaxis due to the following contraindication(s):

enoxaparin (LOVENOX) injection (Single Response)

enoxaparin (LOVENOX) syringe

40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min

30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1

For Patients with CrCL LESS than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1

For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S+1 For Patients with CrCl LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:



<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> High Risk of DVT - Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> High Risk of DVT - Non-Surgical	
Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):

<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily, Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (time critical) For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

High Risk of DVT - Surgical (Hip/Knee)

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

High Risk

High risk of VTE Routine, Once

High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)

<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

### DVT Risk and Prophylaxis Tool (Single Response)

Low Risk Definition Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

Age less than 60 years and NO other VTE risk factors One or more of the following medical conditions: One or more of the following medical conditions:

Patient already adequately anticoagulated CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Age 60 and above Severe fracture of hip, pelvis or leg

Central line Acute spinal cord injury with paresis

History of DVT or family history of VTE Multiple major traumas

Anticipated length of stay GREATER than 48 hours Abdominal or pelvic surgery for CANCER

Less than fully and independently ambulatory Acute ischemic stroke

Estrogen therapy History of PE

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Low Risk of DVT

Low Risk (Single Response)

Low risk of VTE

Routine, Once

Low risk: Due to low risk, no VTE prophylaxis is needed.  
Will encourage early ambulation

Moderate Risk of DVT - Surgical

Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.

Moderate Risk

Moderate risk of VTE

Routine, Once

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)

Patient is currently receiving therapeutic anticoagulation

Routine, Once

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.  
Therapy for the following:

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> Moderate Risk of DVT - Non-Surgical	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input type="checkbox"/> Moderate Risk	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> High Risk of DVT - Surgical Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCl LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> High Risk of DVT - Non-Surgical Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
<input type="checkbox"/> High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (time critical), Starting S
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (time critical), Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical) Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
( <input type="checkbox"/> High Risk of DVT - Surgical (Hip/Knee) Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.	
[ ] High Risk	
<input type="checkbox"/> High risk of VTE	Routine, Once
[ ] High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (time critical), Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (time critical), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (time critical), Starting S+1 To be Given on Post Op Day 1. Indications:

<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 (time critical), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

## Labs Today

### Hematology/Coagulation Today

<input type="checkbox"/> CBC	Once
<input type="checkbox"/> CBC and differential	Once
<input type="checkbox"/> Prothrombin time with INR	Once
<input type="checkbox"/> Partial thromboplastin time	Once

### Chemistry Today

<input type="checkbox"/> Albumin	Once
<input type="checkbox"/> Amylase	Once
<input type="checkbox"/> Basic metabolic panel	Once
<input type="checkbox"/> B-type natriuretic peptide	Once
<input type="checkbox"/> CK total	Once
<input type="checkbox"/> Comprehensive metabolic panel	Once
<input type="checkbox"/> Hemoglobin A1c	Once
<input type="checkbox"/> Hepatic function panel	Once
<input type="checkbox"/> Lactic acid level	Once
<input type="checkbox"/> Lipase	Once
<input type="checkbox"/> Lipid panel	Once
<input type="checkbox"/> Magnesium	Once
<input type="checkbox"/> Phosphorus	Once
<input type="checkbox"/> Prealbumin	Once
<input type="checkbox"/> TSH	Once
<input type="checkbox"/> T4, free	Once
<input type="checkbox"/> Uric acid	Once
<input type="checkbox"/> Urine drugs of abuse screen	Once

### Cardiac Enzymes

<input type="checkbox"/> Troponin I : STAT	STAT For 1 Occurrences
<input type="checkbox"/> Troponin I : Now and every 6 hours x 2	Now then every 6 hours For 2 Occurrences
<input type="checkbox"/> Troponin I : Now and every 8 hours x 2	Now then every 8 hours For 2 Occurrences

### Microbiology

<input type="checkbox"/> Blood culture x 2	<b>"And" Linked Panel</b>
<input type="checkbox"/> Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.
<input type="checkbox"/> Blood Culture (Aerobic & Anaerobic)	Once, Blood Collect before antibiotics given. Blood cultures should be ordered x2, with each set drawn from a different peripheral site. If unable to draw both sets from a peripheral site, please call the lab for assistance; an IV line should NEVER be used.



<input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site:
<input type="checkbox"/> Sputum culture	Once, Sputum

## Labs Tomorrow

### Hematology/Coagulation Tomorrow

<input type="checkbox"/> CBC	AM draw For 1 Occurrences
<input type="checkbox"/> CBC with differential	AM draw For 1 Occurrences
<input type="checkbox"/> Prothrombin time with INR	AM draw For 1 Occurrences
<input type="checkbox"/> Partial thromboplastin time	AM draw For 1 Occurrences

### Chemistry Tomorrow

<input type="checkbox"/> Albumin	AM draw For 1 Occurrences
<input type="checkbox"/> Amylase	AM draw For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	AM draw For 1 Occurrences
<input type="checkbox"/> B-type natriuretic peptide	AM draw For 1 Occurrences
<input type="checkbox"/> CK total	AM draw For 1 Occurrences
<input type="checkbox"/> Comprehensive metabolic panel	AM draw For 1 Occurrences
<input type="checkbox"/> Hepatic function panel	AM draw For 1 Occurrences
<input type="checkbox"/> Lactic acid level	AM draw For 1 Occurrences
<input type="checkbox"/> Lipase	AM draw For 1 Occurrences
<input type="checkbox"/> Lipid panel	AM draw For 1 Occurrences
<input type="checkbox"/> Magnesium	AM draw For 1 Occurrences
<input type="checkbox"/> Phosphorus	AM draw For 1 Occurrences
<input type="checkbox"/> Prealbumin	AM draw For 1 Occurrences
<input type="checkbox"/> TSH	AM draw For 1 Occurrences
<input type="checkbox"/> T4, free	AM draw For 1 Occurrences
<input type="checkbox"/> Uric acid	AM draw For 1 Occurrences
<input type="checkbox"/> Urine drugs of abuse screen	Once, Starting S+1

## Cardiology

### Cardiology

<input type="checkbox"/> Myocardial perfusion stress test	Routine, 1 time imaging Must order Stress Test ECG Only order in conjunction.
<input type="checkbox"/> Cv exercise treadmill stress (no imaging)	Routine, Once
<input type="checkbox"/> ECG 12 lead - Routine	Routine, Once Clinical Indications: Chest Pain Interpreting Physician:
<input type="checkbox"/> ECG 12 lead - STAT	STAT, Once Clinical Indications: Chest Pain Interpreting Physician:
<input type="checkbox"/> Echocardiogram complete w contrast and 3D if needed	Routine, 1 time imaging

## Imaging

### MRI/MRA

<input type="checkbox"/> MRI Brain Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/> MRI Brain W Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/> MRI Brain W Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/> MRA Head Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/> MRA Head W Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/> MRA Head W Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/> MRA Neck Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/> MRA Neck W Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/> MRA Neck W Wo Contrast	Routine, 1 time imaging For 1

### CT

<input type="checkbox"/>	CT Head Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/>	CT Head W Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/>	CT Head W Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/>	CT Chest Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/>	CT Abdomen W Contrast (Omnipaque)	<b>"And" Linked Panel</b>

For those with iodine allergies, please order the panel with Read-Cat (barium sulfate).

<input type="checkbox"/>	CT Abdomen W Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/>	iohexol (OMNIPAQUE) 300 mg iodine/mL oral solution	30 mL, oral, once
<input type="checkbox"/>	CT Abdomen Pelvis WO Contrast (Omnipaque)	<b>"And" Linked Panel</b>

For those with iodine allergies, please order the panel with Read-Cat (barium sulfate).

<input type="checkbox"/>	CT Abdomen Pelvis Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/>	iohexol (OMNIPAQUE) 300 mg iodine/mL oral solution	30 mL, oral, once
<input type="checkbox"/>	CT Abdomen Pelvis WO Contrast (Read-Cat)	<b>"And" Linked Panel</b>

Ordered as secondary option for those with iodine allergies.

<input type="checkbox"/>	CT Abdomen Pelvis Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/>	barium (READI-CAT 2) 2.1 % (w/v), 2.0 % (w/w) suspension	450 mL, oral, once in imaging, contrast, For 1 Doses
<input type="checkbox"/>	CT Pelvis W Contrast (Omnipaque)	<b>"And" Linked Panel</b>

For those with iodine allergies, please order the panel with Read-Cat (barium sulfate).

<input type="checkbox"/>	CT Pelvis W Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/>	iohexol (OMNIPAQUE) 300 mg iodine/mL oral solution	30 mL, oral, once
<input type="checkbox"/>	CT Abdomen Pelvis WO Contrast (Omnipaque)	<b>"And" Linked Panel</b>

For those with iodine allergies, please order the panel with Read-Cat (barium sulfate).

<input type="checkbox"/>	CT Abdomen Pelvis Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/>	iohexol (OMNIPAQUE) 300 mg iodine/mL oral solution	30 mL, oral, once
<input type="checkbox"/>	CT Abdomen Pelvis WO Contrast (Read-Cat)	<b>"And" Linked Panel</b>

Ordered as secondary option for those with iodine allergies.

<input type="checkbox"/>	CT Abdomen Pelvis Wo Contrast	Routine, 1 time imaging For 1
<input type="checkbox"/>	barium (READI-CAT 2) 2.1 % (w/v), 2.0 % (w/w) suspension	450 mL, oral, once in imaging, contrast, For 1 Doses

<input type="checkbox"/>	CT Sinus Wo Contrast	Routine, 1 time imaging For 1
--------------------------	----------------------	-------------------------------

### X-Ray

<input type="checkbox"/>	Chest 1 Vw Portable	Routine, 1 time imaging For 1
<input type="checkbox"/>	Chest 1 Vw Portable	STAT, 1 time imaging For 1 Occurrences
<input type="checkbox"/>	Chest 2 Vw	Routine, 1 time imaging For 1
<input type="checkbox"/>	Chest 2 Vw	STAT, 1 time imaging For 1 Occurrences
<input type="checkbox"/>	KUB Kidney Ureter Bladder	Routine, 1 time imaging For 1
<input type="checkbox"/>	KUB Kidney Ureter Bladder	STAT, 1 time imaging For 1
<input type="checkbox"/>	Abdomen 2 Vw Ap W Upright And/Or Decubitus	Routine, 1 time imaging For 1
<input type="checkbox"/>	Abdomen 2 Vw Ap W Upright And/Or Decubitus	STAT, 1 time imaging For 1

### Ultrasound

<input type="checkbox"/>	US Abdomen Complete	Routine, 1 time imaging For 1
<input type="checkbox"/>	US Gallbladder	Routine, 1 time imaging For 1
<input type="checkbox"/>	US Renal	Routine, 1 time imaging For 1
<input type="checkbox"/>	US Pelvis Complete	Routine, 1 time imaging For 1
<input type="checkbox"/>	US Pelvic Non Ob Limited	Routine, 1 time imaging For 1
<input type="checkbox"/>	US Pelvic Transvaginal	Routine, 1 time imaging For 1
<input type="checkbox"/>	Pv carotid duplex	Routine, 1 time imaging
<input type="checkbox"/>	Pv duplex arterial upper extremity	Routine, 1 time imaging
<input type="checkbox"/>	Pv duplex arterial lower extremity	Routine, 1 time imaging
<input type="checkbox"/>	Pv vascular screening	Routine, 1 time imaging

## Other Studies

## Respiratory

### Respiratory

<input type="checkbox"/> Oxygen therapy - NC 2 Lpm	Routine, Continuous Device 1: Nasal Cannula Rate in liters per minute: 2 lpm Rate in tenths of a liter per minute: O2 %: Titrate to keep O2 Sat Above: 92% Indications for O2 therapy: Hypoxemia Device 2: Device 3: Indications for O2 therapy:
--	---

## Rehab

## Consults

### Pharmacy Consult for Anticoagulation

<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	Routine, Until discontinued, Starting S Indication:
<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S
<input type="checkbox"/> Pharmacy consult to manage dosing of all oral anticoagulants	Routine, Until discontinued, Starting S Which drug do you need help dosing? Contact Number:
<input type="checkbox"/> Pharmacy Consult to Manage Heparin STANDARD dose protocol (DVT/PE) - initiation bolus and infusion with titration boluses	STAT, Until discontinued, Starting S Heparin Indication: Specify: Give initial Bolus Monitoring: Anti-Xa
<input type="checkbox"/> Pharmacy consult to manage Heparin LOW dose protocol (ACS/Stroke/Afib) - initiation bolus and infusion withOUT titration boluses	STAT, Until discontinued, Starting S Heparin Indication: Specify: Monitoring: Anti-Xa

### Ancillary Consults

<input type="checkbox"/> Consult to Case Management	Consult Reason:
<input type="checkbox"/> Consult to Social Work	Reason for Consult:
<input type="checkbox"/> Consult PT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult PT wound care	Special Instructions: Location of Wound?
<input type="checkbox"/> Consult OT eval and treat	Special Instructions: Weight Bearing Status:
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Purpose/Topic:
<input type="checkbox"/> Consult to Spiritual Care	Reason for consult?
<input type="checkbox"/> Consult to Speech Language Pathology	Routine, Once Reason for consult:
<input type="checkbox"/> Consult to Wound Ostomy Care nurse	Reason for consult: Reason for consult: Reason for consult: Reason for consult: Consult for NPWT: Reason for consult:
<input type="checkbox"/> Consult to Respiratory Therapy	Reason for Consult?

## Additional Orders

