

## CVIR Post PICC Line Insertion [694]

### General

### Nursing

#### Nursing

<input type="checkbox"/> Ok to use	Routine, Until discontinued, Starting S Device: PICC line PICC trim length to *** centimeters, PACU & Post-op
<input type="checkbox"/> Needleless Connector	Routine, Every Tue, Fri Change every Tuesday and Friday and as needed. Use positive flow needleless valve, caps only (NO EXCEPTIONS), Post-op
<input type="checkbox"/> Change dressing	Routine, Once And as needed., PACU & Post-op
<input type="checkbox"/> Apply dressing	Routine, Once Apply new dressing 24 hours post procedure., Post-op
<input type="checkbox"/> Use 10 milliliter syringe or greater (NO EXCEPTIONS)	Routine, Until discontinued, Starting S Use 10 milliliter syringe or greater (NO EXCEPTIONS), PACU & Post-op
<input type="checkbox"/> Discharge Home	Routine, Once For 1 Occurrences At *** , if stable. Confirm medication reconciliation is complete., PACU

### IV Fluids

### Medications

#### Post Procedure Medications

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25 mcg, intravenous, every 15 min PRN, severe pain (score 7-10), persistent pain, For 2 Doses, PACU & Post-op Hold for respirations LESS than 12 breath/minute. For persistent pain, call interventional MD.
<input type="checkbox"/> ondansetron (ZOFTRAN) IV	4 mg, intravenous, once PRN, nausea, vomiting, For 1 Doses, PACU & Post-op

### VTE

### Labs

### Cardiology

### Imaging

### Other Diagnostic Studies

### Respiratory

### Rehab

### Consults

### Additional Orders