

General Adult Total Parenteral Nutrition [976]

ADULT TOTAL PARENTERAL NUTRITION

General

Nursing Assessments

<input type="checkbox"/> Strict intake and output	Routine, Every hour For Until specified
<input type="checkbox"/> Measure height	Routine, Once For 1 Occurrences
<input type="checkbox"/> Daily Weights	Routine, Daily For Until specified
<input type="checkbox"/> Neuro checks	Routine, Every 4 hours
<input type="checkbox"/> Neurovascular checks	Routine, Every 4 hours
<input type="checkbox"/> Skin assessment	Routine, Every shift
<input type="checkbox"/> POCT glucose	Once

Nursing - TPN Management

<input type="checkbox"/> Pharmacy Time Parameters for Initiation of TPN	Routine, Continuous 1) New initiation orders for TPN will not be accepted after 7PM on weekdays or 3PM on weekends and holidays per ANW pharmacy policy. 2) If TPN ordered after these times, hang D10W to run at 25 mL/hr until TPN is initiated the following day. (Nurse to order D10W for MAR documentation if initiation after hours).
<input type="checkbox"/> Infuse TPN Through a Central Line	Routine, Continuous For Until specified RN to confirm central line placement prior to infusing TPN.

Ancillary Consults

<input type="checkbox"/> Consult to Nutritional Services	Reason for Consult?
<input type="checkbox"/> Consult to Pastoral/Spiritual Care	Reason for Consult?
<input type="checkbox"/> Consult to Respiratory Therapy	Reason for Consult?
<input type="checkbox"/> Consult to Social Work	Reason for Consult?
<input type="checkbox"/> Tobacco Cessation Education	Routine, Once

Labs

Labs Once

<input type="checkbox"/> Comprehensive metabolic panel	Once For 1 Occurrences
<input type="checkbox"/> Magnesium	Once For 1 Occurrences prior to starting parenteral nutrition
<input type="checkbox"/> Phosphorus	Once For 1 Occurrences prior to starting parenteral nutrition
<input type="checkbox"/> Prealbumin level	Once For 1 Occurrences prior to starting parenteral nutrition
<input type="checkbox"/> Triglycerides	Once For 1 Occurrences

Labs Repeat - All Facilities Except HMSJ

<input type="checkbox"/> Comprehensive Metabolic Panel	AM draw repeats, Starting S+7 For 2 Occurrences
<input type="checkbox"/> Phosphorus	AM draw repeats, Starting S+7 For 2 Occurrences
<input type="checkbox"/> Magnesium level	AM draw repeats, Starting S+7 For 2 Occurrences
<input type="checkbox"/> Prealbumin	AM draw repeats, Starting S+2 For 2 Occurrences
<input type="checkbox"/> Triglycerides	AM draw repeats, Starting S+2 For 2 Occurrences

Labs Repeat - HMSJ Only

<input type="checkbox"/> Comprehensive Metabolic Panel	AM draw repeats, Starting S+1 For 3 Occurrences
<input type="checkbox"/> Phosphorus	AM draw repeats, Starting S+1 For 3 Occurrences
<input type="checkbox"/> Magnesium level	AM draw repeats, Starting S+1 For 3 Occurrences
<input type="checkbox"/> Prealbumin	Every Monday
<input type="checkbox"/> Triglycerides	Every Monday
<input type="checkbox"/> Hepatic function panel	Every Monday

TPN

Pharmacy Consults (Single Response)

<input type="checkbox"/> Pharmacy consult to manage TPN therapy	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted:
<input type="checkbox"/> Pharmacy consult to monitor and provide recommendation ONLY for TPN therapy	STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy: Physician contact number:
<input type="checkbox"/> No pharmacy TPN consult needed	STAT, Once For 1 Occurrences

Pharmacy Consults (Single Response)

<input type="checkbox"/> Pharmacy consult to manage TPN therapy	STAT, Until discontinued, Starting S Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted:
<input type="checkbox"/> No pharmacy TPN consult needed	STAT, Once For 1 Occurrences

Pharmacy Consult (Single Response)

<input type="checkbox"/> Pharmacy consult to monitor and provide recommendation ONLY for TPN therapy	STAT, Once For 1 Occurrences Enteral Nutrition: Indication: Location of venous access: Is patient volume restricted: Instructions to pharmacy: Physician contact number:
--	--

Premixed TPN solutions (CLINIMIX) - HMSL Only

TPN Solutions - HMSL Only (Single Response)

<input type="checkbox"/> amino acids 4.25%-dextrose 5%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 4.25%-dextrose 25%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 5%-dextrose 15%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 5%-dextrose 20%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 5%-dextrose 25%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> fat emulsion (INTRALIPID,LIPOSYN) 20 % infusion	intravenous, nightly at 2100 (time critical)

Premixed TPN solutions (CLINIMIX) - HMW Only

TPN Solutions - HMW Only (Single Response)

<input type="checkbox"/> amino acids 4.25%-dextrose 5%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 5%-dextrose 15%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 5%-dextrose 20%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> fat emulsion (INTRALIPID,LIPOSYN) 20 % infusion	intravenous, nightly at 2100 (time critical)

Premixed TPN solutions (CLINIMIX) HMSJ Only

<input type="checkbox"/> Clinimix Premix TPN Solutions (Single Response)	
<input type="checkbox"/> amino acids 4.25%-dextrose 5%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 5%-dextrose 20%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acid 5 %-dextrose 20 % (CLINIMIX) 2,000 mL with multivitamin 10 mL, multitrace elements-5 concentrate (MTE-5 Concentrate) 1 mL	intravenous, nightly
<input type="checkbox"/> fat emulsion (INTRALIPID,LIPOSYN) 20 % infusion	intravenous, nightly at 2100 (time critical)

Premixed TPN solutions (CLINIMIX) HMSTC (Single Response)

<input checked="" type="checkbox"/> TPN Solutions (Single Response)	
<input type="checkbox"/> amino acids 4.25%-dextrose 5%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 4.25%-dextrose 25%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 5%-dextrose 15%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 5%-dextrose 25%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> amino acids 5%-dextrose 20%- lytes (CLINIMIX E) with MVI and trace	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> fat emulsion (INTRALIPID,LIPOSYN) 20 % infusion	intravenous, nightly at 2100 (time critical)

TPN (HMSTJ Only)

<input checked="" type="checkbox"/> TPN Solutions (Single Response)	
<input type="checkbox"/> Adult Standard Clinimix TPN	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours hang D10 if TPN is interrupted
<input type="checkbox"/> Adult Custom Clinimix TPN	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours hang D10 if TPN is interrupted
<input type="checkbox"/> Adult Custom Central TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM, For 24 Hours Indication:
<input type="checkbox"/> fat emulsion (INTRALIPID,LIPOSYN) 20 % infusion	intravenous, for 12 Hours, nightly at 2100 (time critical)

TPN (HMW Only)

<input checked="" type="checkbox"/> TPN Solutions (Single Response)	
<input type="checkbox"/> Adult Standard Clinimix TPN	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> Adult Custom Clinimix TPN	intravenous, continuous TPN, S at 9:00 PM, For 24 Hours
<input type="checkbox"/> Adult Custom Central TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM, For 24 Hours Indication:
<input type="checkbox"/> fat emulsion (INTRALIPID,LIPOSYN) 20 % infusion	intravenous, for 12 Hours, nightly at 2100 (time critical)

Adult TPN - NOT HMSTJ, HMTW, HMSTC (Single Response)

<input type="checkbox"/> Adult Central TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM, For 24 Hours Indication:
<input type="checkbox"/> Adult Peripheral TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM, For 24 Hours Indication:
<input type="checkbox"/> Adult CYCLIC Peripheral TPN	intravenous, cyclic TPN - see admin instructions, S at 9:00 PM, For 24 Hours Infuse TPN over a total of *** hrs/day. Start at *** ml/hr for the first hour Then increase to *** ml/hr for *** hrs Finally decrease the rate to *** ml/hr for the last hour No TPN is to be infused for the other *** hrs per day. Indication:

<input type="checkbox"/> Adult CYCLIC Central TPN	intravenous, cyclic TPN - see admin instructions, S at 9:00 PM, For 24 Hours Infuse TPN over a total of *** hrs/day. Start at *** ml/hr for the first hour Then increase to *** ml/hr for *** hrs Finally decrease the rate to *** ml/hr for the last hour No TPN is to be infused for the other *** hrs per day. Indication:
---	---

Adult TPN - HMTW Only (Single Response)

<input type="checkbox"/> Adult Central TPN + Fat Emulsion	"And" Linked Panel
<input type="checkbox"/> Adult Central TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM, For 24 Hours Indication:
<input type="checkbox"/> fat emulsion (INTRALIPID,LIPOSYN) 20 % infusion	intravenous, nightly at 2100 (time critical) Discard any unused volume AFTER 12 hours of hang time.
<input type="checkbox"/> Adult Peripheral TPN + Fat Emulsion	"And" Linked Panel
<input type="checkbox"/> Adult Peripheral TPN	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM, For 24 Hours Indication:
<input type="checkbox"/> fat emulsion (INTRALIPID,LIPOSYN) 20 % infusion	intravenous, nightly at 2100 (time critical) Discard any unused volume AFTER 12 hours of hang time.
<input type="checkbox"/> Adult Central TPN WITHOUT Fat Emulsion	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM, For 24 Hours Indication:
<input type="checkbox"/> Adult Peripheral TPN WITHOUT Fat Emulsion	intravenous, for 24 Hours, continuous TPN, S at 9:00 PM, For 24 Hours Indication:

dextrose 10% for Disruption of TPN Therapy

Start 10% dextrose IV during any interruption in TPN at previous TPN rate up to a MAXIMUM rate of 40 mL/hr.

<input checked="" type="checkbox"/> dextrose 10 % infusion	40 mL/hr, intravenous, continuous PRN, other, For disruption of TPN therapy. Start 10% dextrose IV prior to initiation or during any interruption in TPN at the previous TPN rate up to a MAXIMUM rate of 40 mL/hr.
--	--