

IP APML4 INDUCTION

Types: ONCOLOGY TREATMENT

Synonyms: ACUTE, PROMYEL, APL, ATRA, IDAR, ATO, ARSE, APML, ARSENIC

Cycle 1	Repeat 1 time	Cycle length: 45 days
Day 1	Perform every 1 day x1	
Labs	☒ CBC WITH PLATELET AND DIFFERENTIAL	
	Interval: Once	Occurrences: --
	☒ COMPREHENSIVE METABOLIC PANEL	
	Interval: Once	Occurrences: --
	☒ MAGNESIUM LEVEL	
	Interval: Once	Occurrences: --
	☐ LDH	
	Interval: Once	Occurrences: --
	☐ URIC ACID LEVEL	
	Interval: Once	Occurrences: --
	☒ ECG 12-LEAD	
	Interval: Once	Occurrences: --
Provider Communication		
ONC PROVIDER COMMUNICATION 5		
Interval: Once Occurrences: --		
Comments: Use baseline weight to calculate dose. Adjust dose for weight gains/losses of greater than or equal to 10%.		
Provider Communication		
ONC PROVIDER COMMUNICATION		
Interval: Once Occurrences: --		
Comments: Verify Ejection Fraction prior to Cycle 1. Ejection Fraction: ***% on *** (date).		
If patient has not had a recent MUGA or ECHO, order one via order entry. A baseline cardiac evaluation with a MUGA scan or an ECHO is recommended, especially in patients with risk factors for increased cardiac toxicity. Repeated MUGA or ECHO determinations of LVEF should be performed, particularly with higher, cumulative anthracycline doses.		
Supportive Care		
predniSONE (DELTASONE) tablet 1 mg/kg (Treatment Plan)		
Dose: 1 mg/kg Route: oral daily for 10 doses		
Instructions:		
Continue until WBC count is less than 1000, or resolution of differentiation syndrome.		
Chemotherapy		
tretinoin (VESANOID) chemo capsule 22.5 mg/m2 (Treatment Plan)		

Dose: 22.5 mg/m² Route: oral 2 times daily with meals
 Start: S
 Instructions:
 DO NOT CRUSH, CHEW, DISSOLVE, OR
 MANIPULATE CAPSULE IN ANY WAY.
 Administer with meals.

Round to nearest 10 mg capsule (total daily
 dose = 45 mg/m²).

Day 2

Perform every 1 day x1

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions:
 To keep vein open.

Pre-Medications



ondansetron (ZOFTRAN) 16 mg in sodium chloride 0.9% 50 mL IVPB

Dose: -- Route: intravenous once over 15 Minutes for 1 dose
 Start: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	No	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

☐ **ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg Route: oral once for 1 dose
 Start: S

☐ **dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg Route: oral once for 1 dose
 Start: S

☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
 Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN	Base	130 mL	Yes	Yes

WATER (D5W) IV SOLP (EXCEL; NON-PVC) SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes
--	------	--------	----	-----

Provider Communication

ONC PROVIDER COMMUNICATION 25

Interval: Once	Occurrences: --
Comments:	Reduce dose of idarubicin to 9 mg/m2 for patients ages 61 - 70. Reduce dose of idarubicin to 6 mg/m2 for patients ages greater than 70 years old.

Chemotherapy

IDarubicin (IDAmycin) 12 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 12 mg/m2	Route: intravenous	once over 15 Minutes for 1 dose Offset: 30 Minutes
----------------	--------------------	---

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	IDARUBICIN 1 MG/ML INTRAVENOUS SOLUTION	Medications	12 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: Until discontinued	Occurrences: --
Comments:	Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose) 1. Stop the infusion. 2. Place the patient on continuous monitoring. 3. Obtain vital signs. 4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing. 5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once. 6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once. 7. Notify the treating physician. 8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe). 9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: Until discontinued	Occurrences: --
Comments:	Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea,

vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: Until discontinued

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg
Start: S

Route: intravenous PRN

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg
Start: S

Route: oral PRN

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg
Start: S

Route: intravenous PRN

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg

Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg

Route: intravenous PRN

Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT
injection syringe 0.3 mg**

Dose: 0.3 mg

Route: subcutaneous

PRN

Start: S

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL

Route: intravenous

PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units

Route: intra-catheter

once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Day 4

Perform every 1 day x1

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once

Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once

Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once

Occurrences: --

☐ **LDH**

Interval: Once

Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once

Occurrences: --

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

☒ **ondansetron (ZOFTRAN) 16 mg in sodium
chloride 0.9% 50 mL IVPB**

Dose: --

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ONDANSETRON
HCL (PF) 4 MG/2
ML INJECTION
SOLUTION

Medications

16 mg

Yes

No

DEXAMETHASONE
4 MG/ML
INJECTION
SOLUTION

Medications

12 mg

No

No

SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Base

50 mL

Always

Yes

DEXTROSE 5 % IN

Base

No

Yes

WATER (D5W)
INTRAVENOUS
SOLUTION

☐ **ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg
Start: S

Route: oral

once for 1 dose

☐ **dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg
Start: S

Route: oral

once for 1 dose

☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg
Start: S

Route: intravenous
End: S

once over 30 Minutes for 1 dose

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

APREPITANT 7.2
MG/ML

Medications

130 mg

Main

Yes

Ingredient

INTRAVENOUS
EMULSION

DEXTROSE 5 % IN

Base

130 mL

Yes

Yes

WATER (D5W) IV
SOLP (EXCEL;

NON-PVC)

SODIUM
CHLORIDE 0.9 % IV

Base

130 mL

No

Yes

SOLP

(EXCEL;NON-PVC)

Chemotherapy

IDarubicin (IDAmycin) 12 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 12 mg/m2

Route: intravenous

once over 15 Minutes for 1 dose
Offset: 30 Minutes

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

IDARUBICIN 1
MG/ML

Medications

12 mg/m2

Main

Yes

Ingredient

INTRAVENOUS
SOLUTION

SODIUM
CHLORIDE 0.9 %

QS Base

50 mL

Yes

Yes

INTRAVENOUS
SOLUTION

DEXTROSE 5 % IN
WATER (D5W)

QS Base

No

Yes

INTRAVENOUS
SOLUTION

Day 6

Perform every 1 day x1

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once

Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once

Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once

Occurrences: --

☐ **LDH**

Interval: Once

Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once

Occurrences: --

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

☒ **ondansetron (ZOFTRAN) 16 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: --

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ONDANSETRON
HCL (PF) 4 MG/2
ML INJECTION
SOLUTION

Medications

16 mg

Yes

No

DEXAMETHASONE
4 MG/ML
INJECTION
SOLUTION

Medications

12 mg

No

No

SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Base

50 mL

Always

Yes

DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Base

No

Yes

☐ **ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg

Route: oral

once for 1 dose

Start: S

☐ **dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg

Route: oral

once for 1 dose

Start: S

☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg

Route: intravenous

once over 30 Minutes for 1 dose

Start: S

End: S

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

APREPITANT 7.2
MG/ML
INTRAVENOUS
EMULSION

Medications

130 mg

Main

Ingredient

Yes

DEXTROSE 5 % IN
WATER (D5W) IV
SOLP (EXCEL;
NON-PVC)

Base

130 mL

Yes

Yes

SODIUM
CHLORIDE 0.9 % IV
SOLP
(EXCEL;NON-PVC)

Base

130 mL

No

Yes

Chemotherapy

IDArubicin (IDAmycin) 12 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 12 mg/m2

Route: intravenous

once over 15 Minutes for 1 dose

		Ingredients:		Name	Type	Dose	Selected	Adds Vol.
				IDARUBICIN 1 MG/ML INTRAVENOUS SOLUTION	Medications	12 mg/m2	Main Ingredient	Yes
				SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	50 mL	Yes	Yes
				DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes
Offset: 30 Minutes								
Day 8								
Perform every 1 day x1								
Labs								
☒ CBC WITH PLATELET AND DIFFERENTIAL								
Interval: Once Occurrences: --								
☒ COMPREHENSIVE METABOLIC PANEL								
Interval: Once Occurrences: --								
☒ MAGNESIUM LEVEL								
Interval: Once Occurrences: --								
☐ LDH								
Interval: Once Occurrences: --								
☐ URIC ACID LEVEL								
Interval: Once Occurrences: --								
☒ ECG 12-LEAD								
Interval: Once Occurrences: --								
Nursing Orders								
sodium chloride 0.9 % infusion 250 mL								
Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose								
Start: S								
Instructions: To keep vein open.								
Pre-Medications								
☒ ondansetron (ZOFTRAN) 16 mg in sodium chloride 0.9% 50 mL IVPB								
Dose: -- Route: intravenous once over 15 Minutes for 1 dose								
Start: S								
		Ingredients:		Name	Type	Dose	Selected	Adds Vol.
				ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
				DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	No	No
				SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
				DEXTROSE 5 % IN WATER (D5W)	Base		No	Yes

INTRAVENOUS SOLUTION						
☐	ondansetron (ZOFTRAN) tablet 16 mg					
	Dose: 16 mg	Route: oral	once for 1 dose			
	Start: S					
☐	dexamethasone (DECADRON) tablet 12 mg					
	Dose: 12 mg	Route: oral	once for 1 dose			
	Start: S					
☐	aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB					
	Dose: 130 mg	Route: intravenous	once over 30 Minutes for 1 dose			
	Start: S	End: S				
	Ingredients:	Name	Type	Dose	Selected	Adds Vol.
		APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
		DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
		SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Chemotherapy

IDarubicin (IDAmycin) 12 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB						
Dose: 12 mg/m2		Route: intravenous		once over 15 Minutes for 1 dose		
				Offset: 30 Minutes		
Ingredients:		Name	Type	Dose	Selected	Adds Vol.
		IDARUBICIN 1 MG/ML INTRAVENOUS SOLUTION	Medications	12 mg/m2	Main Ingredient	Yes
		SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	50 mL	Yes	Yes
		DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

Day 9

Perform every 1 day x1

Labs

☒ CBC WITH PLATELET AND DIFFERENTIAL		
Interval: Once		Occurrences: --
☒ COMPREHENSIVE METABOLIC PANEL		
Interval: Once		Occurrences: --
☒ MAGNESIUM LEVEL		
Interval: Once		Occurrences: --
☐ LDH		
Interval: Once		Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once

Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once

Occurrences: --

Comments:

Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

☒ **ondansetron (ZOFTRAN) 16 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: --

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ONDANSETRON
HCL (PF) 4 MG/2
ML INJECTION
SOLUTION

Medications

16 mg

Yes

No

DEXAMETHASONE
4 MG/ML
INJECTION
SOLUTION

Medications

12 mg

No

No

SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Base

50 mL

Always

Yes

DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Base

No

Yes

☐ **ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg

Route: oral

once for 1 dose

Start: S

☐ **dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg

Route: oral

once for 1 dose

Start: S

☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg

Route: intravenous

once over 30 Minutes for 1 dose

Start: S

End: S

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

APREPITANT 7.2
MG/ML
INTRAVENOUS
EMULSION

Medications

130 mg

Main

Yes

DEXTROSE 5 % IN
WATER (D5W) IV
SOLP (EXCEL;
NON-PVC)

Base

130 mL

Yes

Yes

SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes
--	------	--------	----	-----

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg	Route: oral	once for 1 dose
Start: S		

Chemotherapy

arsenic trioxide (TRISENOX) 0.15 mg/kg in sodium chloride 0.9% 250 mL chemo IVPB

Dose: 0.15 mg/kg	Route: intravenous	once over 2 Hours for 1 dose
		Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ARSENIC TRIOXIDE 2 MG/ML INTRAVENOUS SOLUTION	Medications	0.15 mg/kg	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes

Electrolyte Replacement

TREATMENT CONDITIONS 39

Interval: --	Occurrences: --
Comments:	Potassium (Normal range 3.5 to 5.0mEq/L)
	o Serum potassium less than 2.9mEq/L, Call MD/NP
	o Serum potassium 3.0 to 3.5mEq/L, give 40mEq KCL IV and 40mEq KCL PO and notify MD/NP
	o Serum potassium 3.6 to 3.9mEq/L, give 40mEq KCL IV or PO
	o Serum potassium 4 mEq/L or greater, do not give potassium replacement

potassium chloride 20 mEq in 100 mL IVPB (FOR CENTRAL LINE ONLY)

Dose: 20 mEq	Route: intravenous	every 1 hour prn over 60 Minutes for 2 doses
Start: S		

potassium chloride (KLOR-CON) packet 40 mEq

Dose: 40 mEq	Route: oral	once PRN
Start: S		

Instructions:

Dissolve contents of each packet in 4 oz. of water or other beverage.

Electrolyte Replacement

TREATMENT CONDITIONS 40

Interval: --	Occurrences: --
Comments:	Magnesium (Normal range 1.6 to 2.6mEq/L)
	o Serum Magnesium less than 1.1mEq/L, Call MD/NP
	o Serum Magnesium 1.2 to 1.5mEq/L, give 4 gram magnesium sulfate IV

- o Serum Magnesium 1.6 to 1.9mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 2 mEq/L or greater, do not give magnesium replacement

magnesium sulfate 2 g/50 mL IVPB (premix)

Dose: 2 g Route: intravenous once PRN @ 25 mL/hr over 1 Hours
Start: S

magnesium sulfate 4 g IVPB (premix)

Dose: 4 g Route: intravenous once PRN over 2 Hours
Start: S

Day 10

Perform every 1 day x1

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --
Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose
Start: S

Chemotherapy

**arsenic trioxide (TRISENOX) 0.15 mg/kg in
sodium chloride 0.9% 250 mL chemo IVPB**

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ARSENIC	Medications	0.15	Main	Yes
	TRIOXIDE 2 MG/ML		mg/kg	Ingredient	

INTRAVENOUS SOLUTION SODIUM CHLORIDE 0.9 %	QS Base	250 mL	Yes	Yes
INTRAVENOUS SOLUTION DEXTROSE 5 % IN WATER (D5W)	QS Base	250 mL	No	Yes
INTRAVENOUS SOLUTION				

Day 11

Perform every 1 day x1

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

☒ **ECG 12-LEAD**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --
 Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose

Start: S

Chemotherapy

**arsenic trioxide (TRISENOX) 0.15 mg/kg in
sodium chloride 0.9% 250 mL chemo IVPB**

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
 Offset: 30 Minutes

Instructions:
 This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ARSENIC	Medications	0.15	Main	Yes

			TRIOXIDE 2 MG/ML INTRAVENOUS SOLUTION		mg/kg	Ingredient	
			SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
			DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes
Days 12,13,14						Perform every 1 day x3	
Labs							
			☒ CBC WITH PLATELET AND DIFFERENTIAL				
			Interval: Once		Occurrences: --		
			☒ COMPREHENSIVE METABOLIC PANEL				
			Interval: Once		Occurrences: --		
			☒ MAGNESIUM LEVEL				
			Interval: Once		Occurrences: --		
			☐ LDH				
			Interval: Once		Occurrences: --		
			☐ URIC ACID LEVEL				
			Interval: Once		Occurrences: --		
Nursing Orders							
			ONC NURSING COMMUNICATION 67				
			Interval: Once		Occurrences: --		
			Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.				
			Refer to Potassium Replacement Scale.				
Nursing Orders							
			sodium chloride 0.9 % infusion 250 mL				
			Dose: 250 mL		Route: intravenous		
			once @ 30 mL/hr for 1 dose				
			Start: S				
			Instructions:				
			To keep vein open.				
Pre-Medications							
			ondansetron (ZOFTRAN) tablet 8 mg				
			Dose: 8 mg		Route: oral		
			once for 1 dose				
			Start: S				
Chemotherapy							
			arsenic trioxide (TRISENOX) 0.15 mg/kg in sodium chloride 0.9% 250 mL chemo IVPB				
			Dose: 0.15 mg/kg		Route: intravenous		
			once over 2 Hours for 1 dose				
			Offset: 30 Minutes				
			Instructions:				
			This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.				
			Ingredients:	Name	Type	Dose	Selected Adds Vol.
				ARSENIC	Medications	0.15	Main Yes
				TRIOXIDE 2 MG/ML		mg/kg	Ingredient
				INTRAVENOUS			

SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes

Day 15

Perform every 1 day x1

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

☒ **ECG 12-LEAD**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --
Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions: To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose
Start: S

Chemotherapy

arsenic trioxide (TRISENOX) 0.15 mg/kg in sodium chloride 0.9% 250 mL chemo IVPB

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name

ARSENIC
TRIOXIDE 2 MG/ML

Type

Medications

Dose

0.15
mg/kg

Selected

Main Ingredient

Adds Vol.

Yes

INTRAVENOUS SOLUTION SODIUM CHLORIDE 0.9 %	QS Base	250 mL	Yes	Yes
INTRAVENOUS SOLUTION DEXTROSE 5 % IN WATER (D5W)	QS Base	250 mL	No	Yes
INTRAVENOUS SOLUTION				

Days 16,17

Perform every 1 day x2

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --
 Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions: To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose
 Start: S

Chemotherapy

arsenic trioxide (TRISENOX) 0.15 mg/kg in sodium chloride 0.9% 250 mL chemo IVPB

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
 Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ARSENIC	Medications	0.15	Main	Yes
	TRIOXIDE 2 MG/ML		mg/kg	Ingredient	
	INTRAVENOUS SOLUTION				

SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes

Day 18

Perform every 1 day x1

Labs

☒ CBC WITH PLATELET AND DIFFERENTIAL

Interval: Once Occurrences: --

☒ COMPREHENSIVE METABOLIC PANEL

Interval: Once Occurrences: --

☒ MAGNESIUM LEVEL

Interval: Once Occurrences: --

☐ LDH

Interval: Once Occurrences: --

☐ URIC ACID LEVEL

Interval: Once Occurrences: --

☒ ECG 12-LEAD

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --
 Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions: To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose
 Start: S

Chemotherapy

arsenic trioxide (TRISENOX) 0.15 mg/kg in sodium chloride 0.9% 250 mL chemo IVPB

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
 Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ARSENIC	Medications	0.15	Main	Yes
	TRIOXIDE 2 MG/ML		mg/kg	Ingredient	
	INTRAVENOUS				

SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes

Days 19,20,21

Perform every 1 day x3

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --
 Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions: To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose
 Start: S

Chemotherapy

arsenic trioxide (TRISENOX) 0.15 mg/kg in sodium chloride 0.9% 250 mL chemo IVPB

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
 Offset: 30 Minutes

Instructions:
 This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ARSENIC	Medications	0.15	Main	Yes
	TRIOXIDE 2 MG/ML		mg/kg	Ingredient	
	INTRAVENOUS				
	SOLUTION				
	SODIUM	QS Base	250 mL	Yes	Yes

CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION
DEXTROSE 5 % IN QS Base 250 mL No Yes
WATER (D5W)
INTRAVENOUS
SOLUTION

Day 22

Perform every 1 day x1

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

☒ **ECG 12-LEAD**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --

Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose

Start: S

Chemotherapy

**arsenic trioxide (TRISENOX) 0.15 mg/kg in
sodium chloride 0.9% 250 mL chemo IVPB**

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name
ARSENIC
TRIOXIDE 2 MG/ML
INTRAVENOUS
SOLUTION

Type
Medications

Dose
0.15
mg/kg

Selected Adds Vol.
Main Yes
Ingredient

SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes

Days 23,24

Perform every 1 day x2

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --
 Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions: To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose
 Start: S

Chemotherapy

arsenic trioxide (TRISENOX) 0.15 mg/kg in sodium chloride 0.9% 250 mL chemo IVPB

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
 Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ARSENIC TRIOXIDE 2 MG/ML INTRAVENOUS SOLUTION	Medications	0.15 mg/kg	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 %	QS Base	250 mL	Yes	Yes

INTRAVENOUS
SOLUTION
DEXTROSE 5 % IN QS Base 250 mL No Yes
WATER (D5W)
INTRAVENOUS
SOLUTION

Day 25

Perform every 1 day x1

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

☒ **ECG 12-LEAD**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --

Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose

Start: S

Chemotherapy

**arsenic trioxide (TRISENOX) 0.15 mg/kg in
sodium chloride 0.9% 250 mL chemo IVPB**

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name

ARSENIC
TRIOXIDE 2 MG/ML
INTRAVENOUS
SOLUTION
SODIUM

Type

Medications

QS Base

Dose

0.15
mg/kg

250 mL

Selected

Main
Ingredient

Adds Vol.

Yes

Yes

CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION
DEXTROSE 5 % IN QS Base 250 mL No Yes
WATER (D5W)
INTRAVENOUS
SOLUTION

Days 26,27,28

Perform every 1 day x3

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --

Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose

Start: S

Chemotherapy

**arsenic trioxide (TRISENOX) 0.15 mg/kg in
sodium chloride 0.9% 250 mL chemo IVPB**

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ARSENIC
TRIOXIDE 2 MG/ML
INTRAVENOUS
SOLUTION

Medications

0.15
mg/kg

Main

Ingredient

SODIUM
CHLORIDE 0.9 %
INTRAVENOUS

QS Base

250 mL

Yes

Yes

SOLUTION
DEXTROSE 5 % IN QS Base 250 mL No Yes
WATER (D5W)
INTRAVENOUS
SOLUTION

Day 29

Perform every 1 day x1

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

☒ **ECG 12-LEAD**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --

Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose

Start: S

Chemotherapy

**arsenic trioxide (TRISENOX) 0.15 mg/kg in
sodium chloride 0.9% 250 mL chemo IVPB**

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ARSENIC
TRIOXIDE 2 MG/ML
INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %

Medications

0.15
mg/kg

Main
Ingredient

QS Base

250 mL

Yes

Yes

INTRAVENOUS
SOLUTION
DEXTROSE 5 % IN QS Base 250 mL No Yes
WATER (D5W)
INTRAVENOUS
SOLUTION

Days 30,31

Perform every 1 day x2

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --

Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose

Start: S

Chemotherapy

**arsenic trioxide (TRISENOX) 0.15 mg/kg in
sodium chloride 0.9% 250 mL chemo IVPB**

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ARSENIC
TRIOXIDE 2 MG/ML
INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Medications

0.15
mg/kg

Main
Ingredient

QS Base

250 mL

Yes

Yes

DEXTROSE 5 % IN QS Base 250 mL No Yes
WATER (D5W)
INTRAVENOUS
SOLUTION

Day 32

Perform every 1 day x1

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

☒ **ECG 12-LEAD**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --

Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose

Start: S

Chemotherapy

**arsenic trioxide (TRISENOX) 0.15 mg/kg in
sodium chloride 0.9% 250 mL chemo IVPB**

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ARSENIC
TRIOXIDE 2 MG/ML
INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS

Medications

0.15
mg/kg

Main

Ingredient

QS Base

250 mL

Yes

Yes

SOLUTION
DEXTROSE 5 % IN QS Base 250 mL No Yes
WATER (D5W)
INTRAVENOUS
SOLUTION

Days 33,34,35

Perform every 1 day x3

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --

Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose

Start: S

Chemotherapy

**arsenic trioxide (TRISENOX) 0.15 mg/kg in
sodium chloride 0.9% 250 mL chemo IVPB**

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ARSENIC
TRIOXIDE 2 MG/ML
INTRAVENOUS
SOLUTION

Medications

0.15
mg/kg

Main Yes
Ingredient

SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

QS Base

250 mL

Yes Yes

DEXTROSE 5 % IN QS Base

250 mL

No Yes

WATER (D5W)
INTRAVENOUS
SOLUTION

Day 36

Perform every 1 day x1

Labs

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

☒ **ECG 12-LEAD**

Interval: Once Occurrences: --

Nursing Orders

ONC NURSING COMMUNICATION 67

Interval: Once Occurrences: --
Comments: Supplement to maintain magnesium GREATER than 2 and potassium GREATER than 4.

Refer to Potassium Replacement Scale.

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions: To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) tablet 8 mg

Dose: 8 mg Route: oral once for 1 dose
Start: S

Chemotherapy

**arsenic trioxide (TRISENOX) 0.15 mg/kg in
sodium chloride 0.9% 250 mL chemo IVPB**

Dose: 0.15 mg/kg Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:

This drug requires close monitoring of electrolytes. Check electrolytes twice weekly and supplement to keep magnesium greater than 2 and potassium greater than 4.

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ARSENIC TRIOXIDE 2 MG/ML INTRAVENOUS SOLUTION	Medications	0.15 mg/kg	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes

DEXTROSE 5 % IN QS Base	250 mL	No	Yes
WATER (D5W)			
INTRAVENOUS			
SOLUTION			