

IP BRENTUXIMAB VEDOTIN

Types: ONCOLOGY TREATMENT

Synonyms: ADCETRIS, LYMPHOMA, HODGKIN, BRENTUX, LIMP

Cycle 1	Repeat 1 time	Cycle length: 21 days
Day 1	Perform every 1 day x1	
Provider Communication	ONC PROVIDER COMMUNICATION 5 Interval: Once Occurrences: -- Comments: Use baseline weight to calculate dose. Adjust dose for weight gains/losses of greater than or equal to 10%.	
Labs	☒ CBC WITH PLATELET AND DIFFERENTIAL Interval: Once Occurrences: -- ☒ COMPREHENSIVE METABOLIC PANEL Interval: Once Occurrences: -- ☒ MAGNESIUM LEVEL Interval: Once Occurrences: -- ☐ LDH Interval: Once Occurrences: -- ☐ URIC ACID LEVEL Interval: Once Occurrences: --	
Nursing Orders	TREATMENT CONDITIONS 7 Interval: Until discontinued Occurrences: -- Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.	
Line Flush	sodium chloride 0.9 % flush 20 mL Dose: 20 mL Route: intravenous PRN Start: S	
Nursing Orders	sodium chloride 0.9 % infusion 250 mL Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose Start: S Instructions: To keep vein open.	
Hydration	sodium chloride 0.9 % infusion Dose: 100 mL/hr Route: intravenous continuous Start: S	
Pre-Medications		

☒ **acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg Route: oral once for 1 dose

Instructions:

Administer 30 minutes prior to chemotherapy.

☒ **diphenhydramine (BENADRYL) injection 25 mg**

Dose: 25 mg Route: intravenous once for 1 dose

Start: S

Instructions:

Administer via slow IV push 30 minutes prior to chemotherapy.

☐ **diphenhydramine (BENADRYL) tablet 25 mg**

Dose: 25 mg Route: oral once for 1 dose

Instructions:

Administer 30 minutes prior to chemotherapy.

☒ **hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**

Dose: 100 mg Route: intravenous once for 1 dose

Instructions:

Administer via slow IV push 30 minutes prior to chemotherapy.

Chemotherapy

brentuximab vedotin (ADCETRIS) 1.8 mg/kg in sodium chloride 0.9 % 100 mL IVPB

Dose: 1.8 mg/kg Route: intravenous once over 30 Minutes for 1 dose
Offset: 30 Minutes

Instructions:

Max dose = 180 mg.

Ingredients:

Name

BRENTUXIMAB
VEDOTIN 50 MG
INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Type

Medications

Dose

1.8 mg/kg

Selected

Main
Ingredient

Adds Vol.

No

Yes

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

☒ **HEparin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Post-Medications

☐ **TBO-FILGRASTIM INJECTION ORDERABLE solution**

Dose: -- Route: subcutaneous

Start: S

Rule-Based Template: RULE ONCBCN
NEUPOGEN WEIGHT BASED

Conditions:

Weight > 72 kg

Weight <= 72 kg

Modifications:

Set dose to 480 mcg

Set dose to 300 mcg