

IP CARBOPLATIN / NAB-PACLITAXEL

Types: ONCOLOGY TREATMENT

Synonyms: ABRAXANE, NAB-PACLITAXEL, NAP, ABE, BRAX, CARBO, LUNG, NON SMALL CELL, NSCLC

Cycle 1	Repeat 1 time	Cycle length: 28 days
Day 1	Perform every 1 day x1	
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: Once Occurrences: --		
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: Once Occurrences: --		
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: Once Occurrences: --		
<input type="checkbox"/> LDH		
Interval: Once Occurrences: --		
<input type="checkbox"/> URIC ACID LEVEL		
Interval: Once Occurrences: --		
<input type="checkbox"/> URINALYSIS, AUTOMATED WITH MICROSCOPY		
Interval: Once Occurrences: --		
Nursing Orders		
TREATMENT CONDITIONS 7		
Interval: Once Occurrences: --		
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.		
Line Flush		
sodium chloride 0.9 % flush 20 mL		
Dose: 20 mL Route: intravenous PRN		
Start: S		
Nursing Orders		
sodium chloride 0.9 % infusion 250 mL		
Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose		
Start: S		
Instructions: To keep vein open.		
Pre-Medications		
ondansetron (ZOFTRAN) 16 mg, dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB		
Dose: -- Route: intravenous once over 15 Minutes for 1 dose		
Start: S End: S 11:30 AM		
Ingredients:	Name	Type Dose Selected Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications 16 mg Yes No
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications 12 mg Yes No

SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
 Start: S End: S 11:30 AM

dexamethasone (DECADRON) tablet 12 mg

Dose: 12 mg Route: oral once for 1 dose
 Start: S

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
 Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Pre-Medications

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
 Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Chemotherapy

NAB-PAClitaxel (ABRAXANE) chemo infusion 100 mg/m2 (Treatment Plan)

Dose: 100 mg/m2 Route: intravenous once over 30 Minutes for 1 dose
 Offset: 30 Minutes

Chemotherapy

CARBOplatin (PARAplatin) in sodium chloride 0.9 % 250 mL chemo IVPB

AUC: 6 Use AUC Route: intravenous once over 60 Minutes for 1 dose

Ingredients:	Name	Offset: 1 Hours		Selected	Adds Vol.
		Type	Dose		
	CARBOPLATIN 10 MG/ML INTRAVENOUS SOLUTION	Medications		Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: Until discontinued

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: Until discontinued

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: Until discontinued
 Comments:

Occurrences: --

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
 Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
 Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
 Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
 Start: S

epinephrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
 Start: S

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEparin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
 Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Labs

CBC WITH PLATELET AND DIFFERENTIAL

Interval: Once Occurrences: --

Nursing Orders

TREATMENT CONDITIONS 7Interval: Once Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Nursing Orders

sodium chloride 0.9 % infusion 250 mLDose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions: To keep vein open.

Pre-Medications

 ondansetron (ZOFTRAN) injection 8 mgDose: 8 mg Route: intravenous once for 1 dose
Start: S End: S 11:15 AM **ondansetron (ZOFTRAN) tablet 16 mg**Dose: 16 mg Route: oral once for 1 dose
Start: S **ondansetron (ZOFTRAN) 16 mg in dextrose 5% 50 mL IVPB**Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Chemotherapy

NAB-PACLitaxel (ABRAXANE) chemo infusion 100 mg/m2 (Treatment Plan)Dose: 100 mg/m2 Route: intravenous once over 30 Minutes for 1 dose
Offset: 30 Minutes

Discharge Nursing Orders

 sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

 HEParin, porcine (PF) injection 500 UnitsDose: 500 Units Route: intra-catheter once PRN
Start: S
Instructions: Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.