

IP CISPLATIN / PACLITAXEL (INTRAPERITONEAL)

Types: ONCOLOGY TREATMENT

Synonyms: SIS, CIS, CISPLATIN, TAXOL, PLATINOL, PAX, TACK

Cycle 1	Repeat 1 time	Cycle length: 21 days
Day 1	Perform every 1 day x1	
Labs		
☒ COMPREHENSIVE METABOLIC PANEL	Interval: Once Occurrences: --	
☒ CBC WITH PLATELET AND DIFFERENTIAL	Interval: Once Occurrences: --	
☒ MAGNESIUM LEVEL	Interval: Once Occurrences: --	
☐ CANCER ANTIGEN 125	Interval: Once Occurrences: --	
☐ LDH	Interval: Once Occurrences: --	
☐ URIC ACID LEVEL	Interval: Once Occurrences: --	
Nursing Orders		
TREATMENT CONDITIONS 13	Interval: Once Occurrences: --	
Comments:	HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000; Serum Creatinine GREATER than 1.5, Total Bilirubin GREATER than 1.5	
Line Flush		
sodium chloride 0.9 % flush 20 mL	Dose: 20 mL Route: intravenous PRN	
Start: S		
Nursing Orders		
sodium chloride 0.9 % infusion 250 mL	Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose	
Start: S		
Instructions:	To keep vein open.	
Pre-Medications		
ondansetron (ZOFTRAN) 16 mg, dexamethasone (DECADRON) 20 mg in sodium chloride 0.9% 50 mL IVPB	Dose: -- Route: intravenous once over 15 Minutes for 1 dose	
Start: S	End: S 11:30 AM	
Ingredients:	Name	Type Dose Selected Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications 16 mg Yes No
	DEXAMETHASONE 4 MG/ML INJECTION	Medications 20 mg Yes No

	SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
		Base		No	Yes
☐ ondansetron (ZOFTRAN) tablet 16 mg					
Dose: 16 mg	Route: oral	once for 1 dose			
Start: S	End: S 11:30 AM				
☐ dexamethasone (DECADRON) tablet 20 mg					
Dose: 20 mg	Route: oral	once for 1 dose			
Start: S					
☐ aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB					
Dose: 130 mg	Route: intravenous	once over 30 Minutes for 1 dose			
Start: S	End: S				
Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Pre-Medications

☐

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg

Route: intravenous

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy via slow IVP.

☒

diphenhydrAMINE (BENADRYL) 50 mg in dextrose 5% 50 mL IVPB

Dose: 50 mg

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:00 AM

Instructions:

Administer 30 minutes prior to chemotherapy

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DIPHENHYDRAMIN E 50 MG/ML INJECTION SOLUTION	Medications	50 mg	Main Ingredient	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	No	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes

☐ **diphenhydramine (BENADRYL) tablet 25 mg**

Dose: 25 mg

Route: oral

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

☒ **famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg

Route: intravenous

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

☐ **famotidine (PEPCID) tablet 20 mg**

Dose: 20 mg

Route: oral

once for 1 dose

Offset: 0 Hours

Instructions:

Administer 30 minutes prior to chemotherapy.

☐ **dexamethasone (DECADRON) injection 20 mg**

Dose: 20 mg

Route: intravenous

once for 1 dose

Offset: 0 Hours

Chemotherapy

PAClitaxel (TAXOL) 135 mg/m2 in sodium chloride 0.9 % 1,000 mL chemo IVPB

Dose: 135 mg/m2

Route: intravenous

once over 24 Hours for 1 dose

Offset: 30 Minutes

Instructions:

Protect from light. Administer through a 0.22 micron filter and non-PVC tubing set.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

PACLITAXEL 6
MG/ML
CONCENTRATE, IN
TRAVENOUS
SODIUM
CHLORIDE 0.9 % IV
SOLP
(EXCEL;NON-PVC)
DEXTROSE 5 % IN
WATER (D5W) IV
SOLP (EXCEL;
NON-PVC)

Medications

135
mg/m2

Main
Ingredient

Yes

QS Base

1,000 mL

Yes

Yes

QS Base

1,000 mL

No

Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: Until
discontinued

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.

2. Place the patient on continuous monitoring.

3. Obtain vital signs.

4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.

5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.

6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

7. Notify the treating physician.

8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: Until discontinued

Occurrences: --

Comments:

- Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)
1. Stop the infusion.
 2. Notify the CERT team and treating physician immediately.
 3. Place the patient on continuous monitoring.
 4. Obtain vital signs.
 5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
 6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
 7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
 8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
 9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: Until discontinued

Occurrences: --

Comments:

- Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)
1. Stop the infusion.
 2. Notify the CERT team and treating physician immediately.
 3. Place the patient on continuous monitoring.
 4. Obtain vital signs.
 5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
 6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
 7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
 8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
 9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
 10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg
Start: S

Route: intravenous

PRN

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg
Start: S

Route: oral

PRN

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Day 2

Perform every 1 day x1

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S

Instructions:

To keep vein open.

Pre-Hydration

dextrose 5%-0.45% sodium chloride 1,000 mL with potassium chloride 20 mEq, magnesium sulfate 2 g infusion

Dose: -- Route: intravenous once @ 500 mL/hr for 1 dose
Start: S

Instructions:

Administer at hour 22 of 24 hour Paclitaxel
(Taxol) infusion.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

DEXTROSE 5 % IN

Base

1,000 mL

No

Yes

WATER (D5W)

INTRAVENOUS

SOLUTION

Pre-Medications

☒ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose

Start: S End: S

Instructions:

Administer 1 hour before Cisplatin.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

☒ **ondansetron (ZOFTRAN) 8 mg, dexamethasone (DECADRON) 20 mg in dextrose 5% 50 mL IVPB**

Dose: -- Route: intravenous once over 15 Minutes for 1 dose

Offset: 30 Minutes

Instructions:

Administer 30 minutes before Cisplatin.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL 2 MG/ML INTRAVENOUS SOLUTION	Medications	8 mg	Yes	No
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	20 mg	Yes	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	No	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes

☒ **LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg Route: intravenous once for 1 dose
Offset: 30 Minutes

Instructions:

Administer 30 minutes before Cisplatin.

Pre-Medications

- ☐ **diphenhydrAMINE (BENADRYL) injection 25 mg**

Dose: 25 mg Route: intravenous once for 1 dose
Start: S
Instructions:
Administer via slow IV push 30 minutes prior to chemotherapy.

- ☐ **diphenhydrAMINE (BENADRYL) 50 mg in sodium chloride 0.9 % 50 mL IVPB**

Dose: 50 mg Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:45 AM
Instructions:
Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DIPHENHYDRAMIN E 50 MG/ML INJECTION SOLUTION	Medications	50 mg	Main Ingredient	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

- ☐ **diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg Route: oral once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

- ☐ **diphenhydrAMINE (BENADRYL) tablet 50 mg**

Dose: 50 mg Route: oral once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

- ☐ **famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg Route: intravenous once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

- ☐ **famotidine (PEPCID) tablet 20 mg**

Dose: 20 mg Route: oral once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

- ☐ **acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg Route: oral once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

Nursing Orders

ONC WARM FLUIDS FOR INTRAPERITONEAL

Interval: Once
Comments:

Occurrences: --
ENSURE ALL FLUIDS ADMINISTERED VIA PERITONEAL ROUTE
ARE WARMED TO PREVENT PATIENT DISCOMFORT.

Hydration

sodium chloride 0.9 % bolus 500 mL

Dose: 500 mL Route: intraperitoneal once for 1 dose

Start: S End: S 5:31 AM

Instructions:

ENSURE THESE FLUIDS ARE WARMED
PRIOR TO ADMINISTERING INTO THE
PERITONEAL AREA!

Administer via gravity infusion.

Chemotherapy

CISplatin (PLATINOL) 100 mg/m2 in sodium chloride 0.9 % 1,000 mL chemo INTRAPERITONEAL

Dose: 100 mg/m2 Route: intraperitoneal once for 1 dose

Start: S End: S 5:53 AM

Instructions:

ENSURE THESE FLUIDS ARE WARMED
PRIOR TO ADMINISTERING INTO THE
PERITONEAL AREA!

Administer via gravity infusion.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

CISPLATIN 1
MG/ML
INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION
DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Medications

100
mg/m2

Main
Ingredient

Yes

QS Base

1,000 mL

Yes

Yes

QS Base

1,000 mL

No

Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: Until
discontinued

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms
only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.

2. Place the patient on continuous monitoring.

3. Obtain vital signs.

4. Administer Normal Saline at 50 mL per hour using a new bag and new
intravenous tubing.

5. If greater than or equal to 30 minutes since the last dose of
Diphenhydramine, administer Diphenhydramine 25 mg intravenous
once.

6. If less than 30 minutes since the last dose of Diphenhydramine,
administer Fexofenadine 180 mg orally and Famotidine 20 mg
intravenous once.

7. Notify the treating physician.

8. If no improvement after 15 minutes, advance level of care to Grade 2
(Moderate) or Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or
otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: Until
discontinued
Comments:

Occurrences: --

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: Until
discontinued
Comments:

Occurrences: --

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg
Start: S

Route: intravenous PRN

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg
Start: S

Route: oral PRN

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg
Start: S

Route: intravenous PRN

hydrocortisone sodium succinate

(Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mgDose: 4 mg Route: intravenous PRN
Start: S**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**Dose: 0.3 mg Route: subcutaneous PRN
Start: S**Day 8**

Perform every 1 day x1

Labs

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: Once Occurrences: --

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: Once Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: Once Occurrences: --

☐ **CANCER ANTIGEN 125**

Interval: Once Occurrences: --

☐ **LDH**

Interval: Once Occurrences: --

☐ **URIC ACID LEVEL**

Interval: Once Occurrences: --

Nursing Orders

TREATMENT CONDITIONS 13Interval: Once Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000; Serum Creatinine GREATER than 1.5, Total Bilirubin GREATER than 1.5

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:
To keep vein open.

Pre-Medications

☒ **dexamethasone (DECADRON) 20 mg in dextrose 5% 50 mL IVPB**

Dose: -- Route: intravenous once over 15 Minutes for 1 dose

Start: S End: S 5:02 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON	Medications		No	No
	HCL 2 MG/ML				
	INTRAVENOUS				
	SOLUTION				
	DEXAMETHASONE	Medications	20 mg	Yes	No
	4 MG/ML				
	INJECTION				
	SOLUTION				
	SODIUM	Base	50 mL	No	Yes
	CHLORIDE 0.9 %				
	INTRAVENOUS				

SOLUTION					
DEXTROSE 5 % IN	Base	50 mL	Yes	Yes	
WATER (D5W)					
INTRAVENOUS					
SOLUTION					

☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg	Route: intravenous	once over 30 Minutes for 1 dose
Start: S	End: S	

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION				
	DEXTROSE 5 % IN	Base	130 mL	Yes	Yes
	WATER (D5W) IV				
	SOLP (EXCEL; NON-PVC)				
	SODIUM CHLORIDE 0.9 % IV	Base	130 mL	No	Yes
	SOLP (EXCEL;NON-PVC)				

Pre-Medications

☐ **diphenhydrAMINE (BENADRYL) injection 25 mg**

Dose: 25 mg	Route: intravenous	once for 1 dose
		Offset: 0 Hours

Instructions:
Administer 30 minutes prior to chemotherapy via slow IVP.

☒ **diphenhydrAMINE (BENADRYL) 50 mg in dextrose 5% 50 mL IVPB**

Dose: 50 mg	Route: intravenous	once over 15 Minutes for 1 dose
Start: S	End: S 11:00 AM	

Instructions:
Administer 30 minutes prior to chemotherapy

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DIPHENHYDRAMIN E 50 MG/ML	Medications	50 mg	Main Ingredient	No
	INJECTION SOLUTION				
	SODIUM CHLORIDE 0.9 %	Base	50 mL	No	Yes
	INTRAVENOUS SOLUTION				
	DEXTROSE 5 % IN	Base	50 mL	Yes	Yes
	WATER (D5W)				
	INTRAVENOUS SOLUTION				

☐ **diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg	Route: oral	once for 1 dose
		Offset: 0 Hours

Instructions:
Administer 30 minutes prior to chemotherapy.

☒ **famotidine (PEPCID) 20 mg/2 mL injection 20 mg**
Dose: 20 mg Route: intravenous once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

☐ **famotidine (PEPCID) tablet 20 mg**
Dose: 20 mg Route: oral once for 1 dose
Offset: 0 Hours
Instructions:
Administer 30 minutes prior to chemotherapy.

☐ **dexamethasone (DECADRON) injection 20 mg**
Dose: 20 mg Route: intravenous once for 1 dose
Offset: 0 Hours

Nursing Orders

ONC WARM FLUIDS FOR INTRAPERITONEAL

Interval: Once Occurrences: --
Comments: ENSURE ALL FLUIDS ADMINISTERED VIA PERITONEAL ROUTE
ARE WARMED TO PREVENT PATIENT DISCOMFORT.

Hydration

sodium chloride 0.9 % bolus 500 mL
Dose: 500 mL Route: intraperitoneal once for 1 dose
Start: S End: S 5:31 AM
Instructions:
ENSURE THESE FLUIDS ARE WARMED
PRIOR TO ADMINISTERING INTO THE
PERITONEAL AREA!
Administer via gravity infusion.

Chemotherapy

PAClitaxel (TAXOL) 60 mg/m2 in sodium chloride 0.9 % 1,000 mL chemo INTRAPERITONEAL
Dose: 60 mg/m2 Route: intraperitoneal once for 1 dose
Start: S End: S 5:50 AM
Instructions:
ENSURE FLUIDS ARE WARMED IN
PHARMACY PRIOR TO MIXING AND
ADMINISTRATION.
Protect from light. Administer through a 0.22
micron filter and non-PVC tubing set.
Administer as rapidly as possible via gravity
infusion. Keep head of bed flat, reposition
patient every 15 minutes: right side, left side,
Trendelenberg, reverse Trendelenberg.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PACLITAXEL 6 MG/ML CONCENTRATE, INTRAVENOUS	Medications	60 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	QS Base	1,000 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	QS Base	1,000 mL	No	Yes

Hydration

sodium chloride 0.9 % bolus 500 mL

Dose: 500 mL

Route: intraperitoneal once for 1 dose

Start: S

End: S 5:31 AM

Instructions:

ENSURE THESE FLUIDS ARE WARMED
PRIOR TO ADMINISTERING INTO THE
PERITONEAL AREA!

Administer via gravity infusion.