

IP DECITABINE (42 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: MDS, DECITABINE, DACOGEN, DEE, LEUKEMIA

Cycle 1	Repeat 1 time	Cycle length: 42 days
Day 1 Perform every 1 day x1		
Provider Communication		
ONC PROVIDER COMMUNICATION 5 Interval: Once Occurrences: -- Comments: Use baseline weight to calculate dose. Adjust dose for weight gains/losses of greater than or equal to 10%.		
Labs		
<input checked="" type="checkbox"/> BASIC METABOLIC PANEL Interval: Once Occurrences: --		
<input type="checkbox"/> COMPREHENSIVE METABOLIC PANEL Interval: Once Occurrences: --		
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL Interval: Once Occurrences: --		
<input checked="" type="checkbox"/> MAGNESIUM LEVEL Interval: Once Occurrences: --		
<input type="checkbox"/> LDH Interval: Once Occurrences: --		
<input type="checkbox"/> URIC ACID LEVEL Interval: Once Occurrences: --		
<input type="checkbox"/> ECHOCARDIOGRAM COMPLETE W CONTRAST AND 3D IF NEEDED Interval: 1 time imaging Occurrences: --		
Line Flush		
sodium chloride 0.9 % flush 20 mL Dose: 20 mL Route: intravenous PRN Start: S		
Nursing Orders		
sodium chloride 0.9 % infusion 250 mL Dose: 250 mL Route: intravenous every 24 hours @ 30 mL/hr for 3 doses Start: S Instructions: To keep vein open.		
Pre-Medications		
<input checked="" type="radio"/> ondansetron (ZOFRAN) injection 8 mg Dose: 8 mg Route: intravenous every 24 hours for 3 doses Start: S		
<input type="radio"/> ondansetron (ZOFRAN) tablet 16 mg Dose: 16 mg Route: oral every 24 hours for 3 doses Start: S		

○ **ondansetron (ZOFTRAN) 16 mg in dextrose 5%
50 mL IVPB**

Dose: 16 mg

Route: intravenous

every 24 hours over 15 Minutes for 3 doses

Start: S

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ONDANSETRON HCL 2 MG/ML INTRAVENOUS SOLUTION	Medications	16 mg	Yes	No
DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications		No	No
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	No	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes

Chemotherapy

decitabine (DACOGEN) 15 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 15 mg/m2

Route: intravenous

every 8 hours over 1 Hours for 9 doses

Offset: 30 Minutes

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
DECITABINE 50 MG INTRAVENOUS SOLUTION	Medications	15 mg/m2	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base		Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: Until discontinued
Comments:

Occurrences: --

- Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
1. Stop the infusion.
 2. Place the patient on continuous monitoring.
 3. Obtain vital signs.
 4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
 5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
 6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
 7. Notify the treating physician.
 8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
 9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: Until discontinued
Comments:

Occurrences: --

- Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)
1. Stop the infusion.
 2. Notify the CERT team and treating physician immediately.
 3. Place the patient on continuous monitoring.
 4. Obtain vital signs.
 5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
 6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
 7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
 8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
 9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: Until discontinued
Comments:

Occurrences: --

- Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)
1. Stop the infusion.
 2. Notify the CERT team and treating physician immediately.
 3. Place the patient on continuous monitoring.
 4. Obtain vital signs.
 5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
 6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.

7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.