Protocol Index

IP FOLFIRI / PANITUMUMAB

Types: ONCOLOGY TREATMENT
Synonyms: FOLFIRI, IRINOTECAN, LEUCOVORIN, 5FU, 5-FLUOROURACIL, FLUOROURACIL, LEUCO, IRIN, CAMPTOSAR, CAMP, CET, PANITUMUMAB, PANT, VECTIBEX, VEX, COLORECTAL, GI

Cycle 1 Repeat 1 time Cycle length: 14 days

Day 1 Perform every 1 day x1

Provider Communication

ONC PROVIDER COMMUNICATION 2
Interval: Once Occurrences: --
Comments: Tumor KRAS gene status should be determined prior to initiation of therapy. KRAS type: Please Push F2:115540219.

Labs

☑ **COMPREHENSIVE METABOLIC PANEL**
Interval: Once Occurrences: --

☑ **CBC WITH PLATELET AND DIFFERENTIAL**
Interval: Once Occurrences: --

☑ **MAGNESIUM LEVEL**
Interval: Once Occurrences: --

Nursing Orders

TREATMENT CONDITIONS 4
Interval: Until discontinued Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000; Total Bilirubin GREATER than 1.5; ALT/AST GREATER than 3 times upper normal limit; or Serum Creatinine GREATER than 1.2

Line Flush

sodium chloride 0.9 % flush 20 mL
Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL
Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

☑ **ondansetron (ZOFRAN) 16 mg, dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB**
Dose: -- Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:30 AM

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Yes	No

SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

☐ **ondansetron (ZOFRAN) tablet 16 mg**

Dose: 16 mg Route: oral once for 1 dose
Start: S End: S 11:30 AM

☐ **dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg Route: oral once for 1 dose
Start: S

☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Start: S End: S

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Pre-Medications

atropine injection 0.25 mg

Dose: 0.25 mg Route: intravenous PRN
Start: S

Supportive Care

○ **LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg Route: intravenous PRN
Start: S

○ **LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg Route: oral PRN
Start: S

Chemotherapy

panitumumab (VECTIBIX) 6 mg/kg in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 6 mg/kg Route: intravenous once over 60 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
USE 0.2 OR 0.22 MICRON INLINE FILTER.

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
PANITUMUMAB 100 MG/5 ML (20 MG/ML) INTRAVENOUS SOLUTION	Medications	6 mg/kg	Main Ingredient	Yes
SODIUM	QS Base	100 mL	Yes	Yes

	CHLORIDE 0.9 % INTRAVENOUS SOLUTION				

leucovorin 400 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 400 mg/m2 Route: intravenous once over 90 Minutes for 1 dose
Offset: 1.5 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	LEUCOVORIN CALCIUM 350 MG SOLUTION FOR INJECTION	Medications	400 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes

irinotecan (CAMPTOSAR) 180 mg/m2 in dextrose 5% 500 mL chemo IVPB

Dose: 180 mg/m2 Route: intravenous once over 90 Minutes for 1 dose
Offset: 1.5 Hours

Instructions:
Protect from light

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	IRINOTECAN 100 MG/5 ML INTRAVENOUS SOLUTION	Medications	180 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	No	Yes

fluorouracil (ADRUCIL) 400 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 400 mg/m2 Route: intravenous once over 15 Minutes for 1 dose
Offset: 3 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	FLUOROURACIL 5 GRAM/100 ML INTRAVENOUS SOLUTION	Medications	400 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	50 mL	Yes	Yes

fluorouracil (ADRUCIL) 1,200 mg/m2 in sodium chloride 0.9 % 500 mL chemo IVPB

Dose: 1,200 mg/m2 Route: intravenous once over 23 Hours for 1 dose
Offset: 3.25 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.

FLUOROURACIL 500 MG/10 ML INTRAVENOUS SOLUTION	Medications	1,200 mg/m2	Main Ingredient	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	No	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: Until discontinued

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: Until discontinued

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)
1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: Until

Occurrences: --

discontinued
Comments: Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)
1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydrAMINE (BENADRYL) injection 25 mg
Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg
Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg
Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg
Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg
Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg
Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Nursing Orders

ONC NURSING COMMUNICATION 14
Interval: Until discontinued Occurrences: --
Comments: Contact Provider if drug-induced acneiform rash develops and covers more than 25 per cent of the body.

Discharge Nursing Orders

☑ **sodium chloride 0.9 % flush 20 mL**
Dose: 20 mL Route: intravenous PRN

☑ **HEParin, porcine (PF) injection 500 Units**
Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Day 2 Perform every 1 day x1

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFRAN) 16 mg in sodium chloride 0.9 % 50 mL IVPB

Dose: -- Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 10:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	No	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

Chemotherapy

fluorouracil (ADRUCIL) 1,200 mg/m2 in sodium chloride 0.9 % 500 mL chemo IVPB

Dose: 1,200 mg/m2 Route: intravenous once over 23 Hours for 1 dose

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	FLUOROURACIL 500 MG/10 ML INTRAVENOUS SOLUTION	Medications	1,200 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	No	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes