

IP LIPOSOMAL DOXORUBICIN (EVERY 14 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: DOXIL, LIP DOXO, DOCKS, LIPOSOMAL, BREAST

Cycle 1	Repeat 1 time	Cycle length: 14 days
Day 1 Perform every 1 day x1		
Labs		
<input checked="" type="checkbox"/>	COMPREHENSIVE METABOLIC PANEL	Interval: Once Occurrences: --
<input checked="" type="checkbox"/>	CBC WITH PLATELET AND DIFFERENTIAL	Interval: Once Occurrences: --
<input checked="" type="checkbox"/>	MAGNESIUM LEVEL	Interval: Once Occurrences: --
<input type="checkbox"/>	CARCINOEMBRYONIC ANTIGEN (CEA)	Interval: Once Occurrences: --
<input type="checkbox"/>	PROSTATE SPECIFIC ANTIGEN	Interval: Once Occurrences: --
<input type="checkbox"/>	URINALYSIS, AUTOMATED WITH MICROSCOPY	Interval: Once Occurrences: --
<input type="checkbox"/>	CANCER ANTIGEN 27-29 (CA BR)	Interval: Once Occurrences: --
Provider Communication		
	ONC PROVIDER COMMUNICATION	Interval: Until discontinued Occurrences: --
	Comments:	Verify Ejection Fraction prior to Cycle 1. Ejection Fraction: ***% on *** (date). If patient has not had a recent MUGA or ECHO, order one via order entry. A baseline cardiac evaluation with a MUGA scan or an ECHO is recommended, especially in patients with risk factors for increased cardiac toxicity. Repeated MUGA or ECHO determinations of LVEF should be performed, particularly with higher, cumulative anthracycline doses.
Nursing Orders		
	TREATMENT CONDITIONS 7	Interval: Until discontinued Occurrences: --
	Comments:	HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.
Line Flush		
	dextrose 5% flush syringe 20 mL	Dose: 20 mL Route: intravenous PRN
	Start: S	
	Instructions:	Administer ONLY for Liposomal Doxorubicin.

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S
Instructions:
Do NOT administer with Liposomal Doxorubicin.

Nursing Orders

dextrose 5% infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open for Liposomal Doxorubicin.

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open. Do NOT administer with Liposomal Doxorubicin.

Pre-Medications

ondansetron (ZOFRAN) injection 8 mg

Dose: 8 mg Route: intravenous once for 1 dose
Start: S End: S 11:15 AM

ondansetron (ZOFRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S

ondansetron (ZOFRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Chemotherapy

DOXOrubicin liposomal (DOXIL) 25 mg/m2 in dextrose 5% 250 mL chemo IVPB

Dose: 25 mg/m2 Route: intravenous once over 1 Hours for 1 dose
Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DOXORUBICIN, PEGYLATED LIPOSOMAL 2 MG/ML INTRAVENOUS SUSPENSION	Medications	25 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes

Interval: Until discontinued
Comments:

Occurrences: --

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: Until discontinued
Comments:

Occurrences: --

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: Until discontinued
Comments:

Occurrences: --

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new

bag and new intravenous tubing.

8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.

9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.

10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramINE (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Supportive Care

LORAZepam (ATIVAN) injection 1 mg

Dose: 1 mg Route: intravenous PRN
Start: S

LORAZepam (ATIVAN) tablet 1 mg

Dose: 1 mg Route: oral PRN
Start: S

Supportive Care

promethazine (PHENERGAN) injection 12.5 mg

Dose: 12.5 mg Route: injection every 6 hours PRN
Start: S

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.