

IP R-CEOP

Types: ONCOLOGY TREATMENT

Synonyms: CEOP, RCEOP, R-CEOP, LYMPHOMA, LYMPH, CYCLO, CYCLOPHOSPHAMIDE, VINC, VINCRISTINE, ETOP, ETOPOSIDE, NON, PRED, PREDNISONE

Cycle 1	Repeat 1 time	Cycle length: 21 days
Day 1	Perform every 1 day x1	
Provider Communication	ONC PROVIDER COMMUNICATION 5	
	Interval: Once	Occurrences: --
	Comments:	Use baseline weight to calculate dose. Adjust dose for weight gains/losses of greater than or equal to 10%.
	Labs	
	☒ COMPREHENSIVE METABOLIC PANEL	
	Interval: Once	Occurrences: --
	☒ CBC WITH PLATELET AND DIFFERENTIAL	
	Interval: Once	Occurrences: --
	☒ MAGNESIUM LEVEL	
	Interval: Once	Occurrences: --
	☐ LDH	
	Interval: Once	Occurrences: --
	☐ URIC ACID LEVEL	
	Interval: Once	Occurrences: --
	☐ ECHOCARDIOGRAM COMPLETE W CONTRAST AND 3D IF NEEDED	
	Interval: 1 time imaging	Occurrences: --
Provider Communication	ONC PROVIDER COMMUNICATION 58	
	Interval: Once	Occurrences: --
	Comments:	Prior to beginning Rituxan infusion, please check if a Hepatitis B and C serology has been performed within the past 6 months. Hepatitis B and C serologies results: Push F2:11554001 drawn on ***.
Nursing Orders	TREATMENT CONDITIONS 7	
	Interval: Once	Occurrences: --
	Comments:	HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.
Line Flush	sodium chloride 0.9 % flush 20 mL	
	Dose: 20 mL	Route: intravenous PRN
	Start: S	
Nursing Orders	sodium chloride 0.9 % infusion 250 mL	
	Dose: 250 mL	Route: intravenous once @ 30 mL/hr for 1 dose
	Start: S	
	Instructions:	To keep vein open.

Pre-Medications

☒ **ondansetron (ZOFTRAN) 16 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: --

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:30 AM

Ingredients:**Name****Type****Dose****Selected****Adds Vol.**ONDANSETRON
HCL (PF) 4 MG/2
ML INJECTION
SOLUTION

Medications

16 mg

Yes

No

DEXAMETHASONE
4 MG/ML
INJECTION
SOLUTION

Medications

12 mg

No

No

SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION
DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Base

50 mL

Always

Yes

Base

No

Yes

☐ **dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: --

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:30 AM

Instructions:Give ONLY if patient had NOT taken their
scheduled dose of ORAL dexamethasone on
day of chemotherapy treatment.**Ingredients:****Name****Type****Dose****Selected****Adds Vol.**ONDANSETRON
HCL (PF) 4 MG/2
ML INJECTION
SOLUTION

Medications

16 mg

No

No

DEXAMETHASONE
4 MG/ML
INJECTION
SOLUTION

Medications

12 mg

Yes

No

SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Base

50 mL

Always

Yes

DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Base

No

Yes

☐ **ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg

Route: oral

once for 1 dose

Start: S

End: S 11:30 AM

☐ **dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg

Route: oral

once for 1 dose

Start: S

☐ **palonosetron (ALOXI) injection 0.25 mg**

Dose: 250 mcg

Route: intravenous

once for 1 dose

Start: S

End: S 3:00 PM

☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg

Start: S

Ingredients:

Route: intravenous

End: S

once over 30 Minutes for 1 dose

Name

APREPITANT 7.2
MG/ML

Type

Medications

Dose

130 mg

Selected

Main

Ingredient

Adds Vol.

Yes

INTRAVENOUS
EMULSION

DEXTROSE 5 % IN
WATER (D5W) IV

Base

130 mL

Yes

Yes

SOLP (EXCEL;
NON-PVC)

SODIUM
CHLORIDE 0.9 % IV

Base

130 mL

No

Yes

SOLP
(EXCEL;NON-PVC)

Chemotherapy

cyclophosphamide (CYTOXAN) 750 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 750 mg/m2

Route: intravenous

once over 60 Minutes for 1 dose

Offset: 30 Minutes

Ingredients:

Name

CYCLOPHOSPHAM
IDE 1 GRAM

Type

Medications

Dose

750
mg/m2

Selected

Main

Ingredient

Adds Vol.

Yes

INTRAVENOUS
SOLUTION

SODIUM
CHLORIDE 0.9 %

QS Base

250 mL

Yes

Yes

INTRAVENOUS
SOLUTION

DEXTROSE 5 % IN
WATER (D5W)

QS Base

250 mL

No

Yes

INTRAVENOUS
SOLUTION

vinCRISTine (ONCOVIN) 1.4 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 1.4 mg/m2

Route: intravenous

once over 15 Minutes for 1 dose

Offset: 1.5 Hours

Instructions:

Max dose = 2 mg.

Rule-Based Template: RULE ONCBCN

VINCRIStINE 1.4 MG/M2

Conditions:

BSA < 1.43 m2

BSA >= 1.43 m2

Modifications:

Set dose to 1.4 mg/m2

Set dose to 2 mg

Ingredients:

Name

VINCRIStINE 1
MG/ML

Type

Medications

Dose

1.4
mg/m2

Selected

Main

Ingredient

Adds Vol.

Yes

INTRAVENOUS
SOLUTION

SODIUM
CHLORIDE 0.9 %

QS Base

50 mL

Yes

Yes

INTRAVENOUS
SOLUTION

etoposide (TOPOSAR) 50 mg/m2 in sodium chloride (NON-PVC) 0.9 % 1,000 mL chemo IVPB

Dose: 50 mg/m2

Route: intravenous

once over 24 Hours for 1 dose

Offset: 1.75 Hours

Instructions:

Administer through a 0.22 micron filter and
non-PVC tubing set. Final concentration: 0.2

mg/mL.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ETOPOSIDE 20
MG/ML

Medications

50 mg/m2 Main
Ingredient

INTRAVENOUS
SOLUTION

SODIUM
CHLORIDE 0.9 % IV
SOLP

QS Base

1,000 mL Yes

Yes

(EXCEL;NON-PVC)

predniSONE (DELTASONE) tablet 100 mg

Dose: 100 mg

Route: oral

daily for 5 doses

Start: S

Rituximab Pre-Medications

acetaminophen (TYLENOL) tablet 650 mg

Dose: 650 mg

Route: oral

once for 1 dose

Start: S

Instructions:

Give 30 minutes before rituximab infusion.

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg

Route: intravenous

once for 1 dose

Start: S

Instructions:

Give 30 minutes before rituximab infusion.

sodium chloride 0.9 % infusion 500 mL

Dose: 500 mL

Route: intravenous

continuous

Start: S

Pharmacy Consult

**PHARMACY CONSULT TO SCREEN FOR
RAPID RITUXIMAB INFUSION**

Interval: --

Occurrences: --

Rituximab Instructions

VITAL SIGNS - T/P/R/BP PER UNIT PROTOCOL

Interval: Once

Occurrences: --

Comments:

1) During Rituximab infusion:

-Vitals every 15 minutes during 1st hour of infusion, THEN

-Every 30 minutes for 1 hour, THEN

-Every hour until end of infusion

-Call MD if SBP less than 90, pulse less than 60 or greater than 120,
temperature greater than 38.5 degrees C

ONC NURSING COMMUNICATION 26

Interval: Once

Occurrences: --

Comments:

2) Infuse antibody via pump

3) If any of the following occurs: FEVER (T greater than 38.5 degrees C),
RIGORS, HYPOTENSION (30mm Hg decrease from baseline), and / or
MUCOSAL CONGESTION / EDEMA, HOLD infusion until improvement
of symptoms (When symptoms improve, resume infusion at HALF the
previous rate)

Chemotherapy

RiTUXimab (PF) (RITUXAN) 375 mg/m2 in

☒ **sodium chloride 0.9% INITIAL INFUSION RATE
IVPB**

Dose: 375 mg/m2

Route: intravenous

once for 1 dose

Offset: 2 Hours

Instructions:

Initiate infusion at rate of 50 mg/hour. In the

absence of infusion toxicity (SBP within 20 mmHG of baseline, PULSE between 60 and 120 and TEMP less than 38 degrees C, and no symptoms), then increase infusion rate by 50 mg/hour every 30 minutes, to a maximum rate of 400 mg/hour.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	RITUXIMAB 10 MG/ML CONCENTRATE, IN TRAVENOUS SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Medications	375 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		Yes	Yes
		Base		No	Yes

☐ **RiTUXImab (PF) (RITUXAN) in sodium chloride 0.9% NON-INITIAL INFUSION IVPB**

Dose: -- Route: intravenous once for 1 dose
Offset: 2 Hours

Instructions:

Initiate infusion rate at a 100 mg/hour. In the absence of infusion toxicity (SBP within 20 mmHG of baseline, PULSE between 60 and 120 and TEMP less than 38 degrees C, and no symptoms), increase rate by 100 mg/hour increments at 30 minute intervals, to a maximum rate of 400 mg/hour.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	RITUXIMAB 10 MG/ML CONCENTRATE, IN TRAVENOUS SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Medications		Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		Yes	Yes
		Base		No	Yes

☐ **RiTUXImab (PF) (RITUXAN) 375 mg/m2 in sodium chloride 0.9% 250 mL RAPID INFUSION RATE IVPB**

Dose: 375 mg/m2 Route: intravenous once over 90 Minutes for 1 dose
Offset: 2 Hours

Instructions:

RAPID INFUSION RATE: Initiate infusion at a rate of 100mL/hour. After 30 minutes, in the absence of infusion reactions (SBP within 20 mmHg of baseline, PULSE between 60 and 120 and Temp less than 38 degrees C, and no symptoms), increase the infusion rate to 200mL/hour. This infusion should take 90 minutes to administer.

Reaction grades:

Grade 3 Reaction: Prolonged (e.g., not rapidly responsive to symptomatic medication and/or brief interruption of infusion); recurrence of symptoms following initial improvement; hospitalization indicated for clinical sequelae (e.g., renal impairment, pulmonary infiltrates).

Grade 4 Reaction: Life-threatening consequences; urgent intervention indicated (e.g., vasopressors or ventilator support).

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	RITUXIMAB 10 MG/ML CONCENTRATE, IN TRAVENOUS SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Medications	375 mg/m2	Main Ingredient	Yes
	QS Base	250 mL	Yes	Yes	
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	No	Yes	

Rituximab Infusion Reaction Orders

meperidine (DEMEROL) injection 25 mg		
Dose: 25 mg	Route: intravenous	once PRN
Start: S		
diphenhydrAMINE (BENADRYL) injection 25 mg		
Dose: 25 mg	Route: intravenous	once PRN
Start: S		
hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg		
Dose: 100 mg	Route: intravenous	once PRN
famotidine (PEPCID) injection 20 mg		
Dose: 20 mg	Route: intravenous	once PRN
Start: S		

Rituximab Additional Orders

epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.3 mg		
Dose: 0.3 mg	Route: intramuscular	once PRN
Start: S		

Discharge Nursing Orders

☒ sodium chloride 0.9 % flush 20 mL		
Dose: 20 mL	Route: intravenous	PRN
☒ HEParin, porcine (PF) injection 500 Units		
Dose: 500 Units	Route: intra-catheter	once PRN
Start: S		
Instructions:		
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.		

Days 2,3

Perform every 1 day x2

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

☒ **ondansetron (ZOFRAN) injection 8 mg**

Dose: 8 mg Route: intravenous once for 1 dose
Start: S

☐ **ondansetron (ZOFRAN) tablet 16 mg**

Dose: 16 mg Route: oral once for 1 dose
Start: S

☐ **ondansetron (ZOFRAN) 16 mg in dextrose 5% 50 mL IVPB**

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S

Ingredients:

Name

Type

Dose
16 mg

Selected
Yes

Adds Vol.
No

ONDANSETRON
HCL 2 MG/ML
INTRAVENOUS
SOLUTION

Medications

DEXAMETHASONE
4 MG/ML
INJECTION
SOLUTION

Medications

No

No

SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Base

50 mL

No

Yes

DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Base

50 mL

Yes

Yes

Chemotherapy

etoposide (TOPOSAR;VEPESID) chemo capsule 100 mg/m2 (Treatment Plan)

Dose: 100 mg/m2 Route: oral once for 1 dose
Start: S