

NHL IP BENDAMUSTINE 120 / RITUXIMAB 375 (START ON DAY 2)

Types: ONCOLOGY TREATMENT

Synonyms: BENDAMUSTINE, BEND, LYMPH, BENDEKA, NON, NHL, 21 DAYS, EVERY 21, Q21D, RITUX, BR

Cycle 1		Repeat 1 time	Cycle length: 28 days		
Day 1		Perform every 1 day x1			
Labs					
<input checked="" type="checkbox"/>	COMPREHENSIVE METABOLIC PANEL		Interval: Once	Occurrences: --	
<input checked="" type="checkbox"/>	CBC WITH PLATELET AND DIFFERENTIAL		Interval: Once	Occurrences: --	
<input checked="" type="checkbox"/>	MAGNESIUM LEVEL		Interval: Once	Occurrences: --	
<input type="checkbox"/>	LDH		Interval: Once	Occurrences: --	
<input type="checkbox"/>	URIC ACID LEVEL		Interval: Once	Occurrences: --	
<input type="checkbox"/>	ECHOCARDIOGRAM COMPLETE W CONTRAST AND 3D IF NEEDED		Interval: 1 time imaging	Occurrences: --	
Nursing Orders					
TREATMENT CONDITIONS 7					
Interval: Until discontinued			Occurrences: --		
Comments:			HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.		
Line Flush					
sodium chloride 0.9 % flush 20 mL					
Dose: 20 mL		Route: intravenous		PRN	
Start: S					
Nursing Orders					
sodium chloride 0.9 % infusion 250 mL					
Dose: 250 mL		Route: intravenous		once @ 30 mL/hr for 1 dose	
Instructions: To keep vein open.					
Pre-Medications					
ondansetron (ZOFTRAN) 16 mg, dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB					
Dose: --		Route: intravenous		once over 15 Minutes for 1 dose	
Start: S					
Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
	DEXAMETHASONE 4 MG/ML INJECTION	Medications	12 mg	Yes	No

SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

ondansetron (ZOFRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S

dexamethasone (DECADRON) tablet 12 mg

Dose: 12 mg Route: oral once for 1 dose
Start: S

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Chemotherapy

bendamustine (BENDEKA) 120 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 120 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
VESICANT

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	BENDAMUSTINE 25 MG/ML INTRAVENOUS SOLUTION	Medications	120 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:
Concentration: 100 units/mL. Heparin flush for

Implanted Vascular Access Device maintenance.

Day 2

Perform every 1 day x1

Labs

BASIC METABOLIC PANEL

Interval: Once Occurrences: --

Provider Communication

ONC PROVIDER COMMUNICATION 58

Interval: Once Occurrences: --

Comments: Prior to beginning Rituxan infusion, please check if a Hepatitis B and C serology has been performed within the past 6 months. Hepatitis B and C serologies results: Push F2:11554001 drawn on ***.

Nursing Orders

TREATMENT CONDITIONS 7

Interval: Until Occurrences: --

discontinued

Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Instructions:

To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg, dexamethasone

- (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: -- Route: intravenous once over 15 Minutes for 1 dose

Start: S

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Yes	No
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

- ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg Route: oral once for 1 dose
Start: S

- dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg Route: oral once for 1 dose
Start: S

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Chemotherapy

bendamustine (BENDEKA) 120 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 120 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
VESICANT

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	BENDAMUSTINE 25 MG/ML	Medications	120 mg/m2	Main Ingredient	Yes
	INTRAVENOUS SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes

Rituximab Pre-Medications

acetaminophen (TYLENOL) tablet 650 mg

Dose: 650 mg Route: oral once for 1 dose
Start: S

Instructions:
Give 30 minutes before rituximab infusion.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous once for 1 dose
Start: S

Instructions:
Give 30 minutes before rituximab infusion.

sodium chloride 0.9 % infusion 500 mL

Dose: 500 mL Route: intravenous continuous
Start: S

Instructions:
Give 30 minutes before rituximab infusion.

Pharmacy Consult

PHARMACY CONSULT TO SCREEN FOR RAPID RITUXIMAB INFUSION

Interval: -- Occurrences: --

Chemotherapy

RiTUXimab (PF) (RITUXAN) 375 mg/m2 in sodium chloride 0.9% INITIAL INFUSION RATE

IVPB

Dose: 375 mg/m² Route: intravenous once for 1 dose
 Offset: 60 Minutes

Instructions:

Administer on Day 2.
 Initiate infusion at rate of 50 mg/hour. In the absence of infusion toxicity (SBP within 20 mmHG of baseline, PULSE between 60 and 120 and TEMP less than 38 degrees C, and no symptoms), then increase infusion rate by 50 mg/hour every 30 minutes, to a maximum rate of 400 mg/hour.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	RITUXIMAB 10 MG/ML CONCENTRATE, INTRAVENOUS	Medications	375 mg/m ²	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base		Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

RiTUXimab (PF) (RITUXAN) in sodium chloride 0.9% NON-INITIAL INFUSION IVPB

Dose: -- Route: intravenous once for 1 dose
 Offset: 60 Minutes

Instructions:

Administer on Day 2.
 Initiate infusion rate at a 100 mg/hour. In the absence of infusion toxicity (SBP within 20 mmHG of baseline, PULSE between 60 and 120 and TEMP less than 38 degrees C, and no symptoms), increase rate by 100 mg/hour increments at 30 minute intervals, to a maximum rate of 400 mg/hour.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	RITUXIMAB 10 MG/ML CONCENTRATE, INTRAVENOUS	Medications		Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base		Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

RiTUXimab (PF) (RITUXAN) 375 mg/m² in sodium chloride 0.9% 250 mL RAPID INFUSION RATE IVPB

Dose: 375 mg/m² Route: intravenous once over 90 Minutes for 1 dose
 Offset: 60 Minutes

Instructions:

Administer on Day 2.
 RAPID INFUSION RATE: Initiate infusion at a rate of 100mL/hour. After 30 minutes, in the absence of infusion reactions (SBP within 20

mmHg of baseline, PULSE between 60 and 120 and Temp less than 38 degrees C, and no symptoms), increase the infusion rate to 200mL/hour. This infusion should take 90 minutes to administer.

Reaction grades:

Grade 3 Reaction: Prolonged (e.g., not rapidly responsive to symptomatic medication and/or brief interruption of infusion); recurrence of symptoms following initial improvement; hospitalization indicated for clinical sequelae (e.g., renal impairment, pulmonary infiltrates).

Grade 4 Reaction: Life-threatening consequences; urgent intervention indicated (e.g., vasopressors or ventilator support).

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	RITUXIMAB 10 MG/ML CONCENTRATE, IN TRAVENOUS	Medications	375 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

Rituximab Instructions

VITAL SIGNS - T/P/R/BP PER UNIT PROTOCOL

Interval: Until discontinued
Comments:

Occurrences: --

- 1) During Rituximab infusion:
 - Vitals every 15 minutes during 1st hour of infusion, THEN
 - Every 30 minutes for 1 hour, THEN
 - Every hour until end of infusion
 - Call MD if SBP less than 90, pulse less than 60 or greater than 120, temperature greater than 38.5 degrees C

ONC NURSING COMMUNICATION 26

Interval: Until discontinued
Comments:

Occurrences: --

- 2) Infuse antibody via pump
- 3) If any of the following occurs: FEVER (temperature greater than 38.5 degrees Celsius), RIGORS, HYPOTENSION (30mm Hg decrease from baseline), and/or MUCOSAL CONGESTION / EDEMA, HOLD infusion until improvement of symptoms (When symptoms improve, resume infusion at HALF the previous rate)

Rituximab Infusion Reaction Orders

meperidine (DEMEROL) injection 25 mg

Dose: 25 mg Route: intravenous once PRN
Start: S

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous once PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous once PRN

famotidine (PEPCID) injection 20 mg

Dose: 20 mg Route: intravenous once PRN
Start: S

Rituximab Additional Orders

epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.3 mg

Dose: 0.3 mg Route: intramuscular once PRN
Start: S

Cycles 2,3

Repeat 2 times

Cycle length: 28 days

Day 1

Perform every 1 day x1

Labs

COMPREHENSIVE METABOLIC PANEL

Interval: Once Occurrences: --

CBC WITH PLATELET AND DIFFERENTIAL

Interval: Once Occurrences: --

MAGNESIUM LEVEL

Interval: Once Occurrences: --

LDH

Interval: Once Occurrences: --

URIC ACID LEVEL

Interval: Once Occurrences: --

ECHOCARDIOGRAM COMPLETE W CONTRAST AND 3D IF NEEDED

Interval: 1 time imaging Occurrences: --

Nursing Orders

TREATMENT CONDITIONS 7

Interval: Until discontinued Occurrences: --

Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg, dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB

Dose: -- Route: intravenous once over 15 Minutes for 1 dose
Start: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2	Medications	16 mg	Yes	No

ML INJECTION
SOLUTION
DEXAMETHASONE Medications 12 mg Yes No
4 MG/ML
INJECTION
SOLUTION
SODIUM Base 50 mL Always Yes
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION
DEXTROSE 5 % IN Base No Yes
WATER (D5W)
INTRAVENOUS
SOLUTION

ondansetron (ZOFRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S

dexamethasone (DECADRON) tablet 12 mg

Dose: 12 mg Route: oral once for 1 dose
Start: S

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Chemotherapy

bendamustine (BENDEKA) 120 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 120 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
VESICANT

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	BENDAMUSTINE 25 MG/ML	Medications	120 mg/m2	Main Ingredient	Yes
	INTRAVENOUS SOLUTION SODIUM CHLORIDE 0.9 %	Base	50 mL	Yes	Yes
	INTRAVENOUS SOLUTION				

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Day 2

Perform every 1 day x1

Labs

BASIC METABOLIC PANEL

Interval: Once Occurrences: --

Nursing Orders

TREATMENT CONDITIONS 7

Interval: Until discontinued Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg, dexamethasone

(DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB

Dose: -- Route: intravenous once over 15 Minutes for 1 dose
Start: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Yes	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S

dexamethasone (DECADRON) tablet 12 mg

Dose: 12 mg Route: oral once for 1 dose
Start: S

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
 Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Chemotherapy

bendamustine (BENDEKA) 120 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 120 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
 Offset: 30 Minutes

Instructions:
VESICANT

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	BENDAMUSTINE 25 MG/ML	Medications	120 mg/m2	Main Ingredient	Yes
	INTRAVENOUS SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes

Rituximab Pre-Medications

acetaminophen (TYLENOL) tablet 650 mg

Dose: 650 mg Route: oral once for 1 dose
 Start: S

Instructions:
Give 30 minutes before rituximab infusion.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous once for 1 dose
 Start: S

Instructions:
Give 30 minutes before rituximab infusion.

sodium chloride 0.9 % infusion 500 mL

Dose: 500 mL Route: intravenous continuous
 Start: S

Instructions:
Give 30 minutes before rituximab infusion.

Pharmacy Consult

PHARMACY CONSULT TO SCREEN FOR RAPID RITUXIMAB INFUSION

Interval: -- Occurrences: --

Chemotherapy

RiTUXimab (PF) (RITUXAN) in sodium chloride 0.9% NON-INITIAL INFUSION IVPB

Dose: -- Route: intravenous once for 1 dose
 Offset: 60 Minutes

Instructions:

Administer on Day 2.

Initiate infusion rate at a 100 mg/hour. In the absence of infusion toxicity (SBP within 20 mmHG of baseline, PULSE between 60 and 120 and TEMP less than 38 degrees C, and no symptoms), increase rate by 100 mg/hour increments at 30 minute intervals, to a maximum rate of 400 mg/hour.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	RITUXIMAB 10 MG/ML CONCENTRATE, INTRAVENOUS	Medications		Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base		Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

RiTUXImab (PF) (RITUXAN) 375 mg/m2 in

sodium chloride 0.9% 250 mL RAPID INFUSION RATE IVPB

Dose: 375 mg/m2

Route: intravenous

once over 90 Minutes for 1 dose
Offset: 60 Minutes

Instructions:

Administer on Day 2.

RAPID INFUSION RATE: Initiate infusion at a rate of 100mL/hour. After 30 minutes, in the absence of infusion reactions (SBP within 20 mmHg of baseline, PULSE between 60 and 120 and Temp less than 38 degrees C, and no symptoms), increase the infusion rate to 200mL/hour. This infusion should take 90 minutes to administer.

Reaction grades:

Grade 3 Reaction: Prolonged (e.g., not rapidly responsive to symptomatic medication and/or brief interruption of infusion); recurrence of symptoms following initial improvement; hospitalization indicated for clinical sequelae (e.g., renal impairment, pulmonary infiltrates).

Grade 4 Reaction: Life-threatening consequences; urgent intervention indicated (e.g., vasopressors or ventilator support).

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	RITUXIMAB 10 MG/ML CONCENTRATE, INTRAVENOUS	Medications	375 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W)	QS Base		No	Yes

INTRAVENOUS
SOLUTION

Rituximab Instructions

VITAL SIGNS - T/P/R/BP PER UNIT PROTOCOL

Interval: Until discontinued

Occurrences: --

Comments:

- 1) During Rituximab infusion:
 - Vitals every 15 minutes during 1st hour of infusion, THEN
 - Every 30 minutes for 1 hour, THEN
 - Every hour until end of infusion
 - Call MD if SBP less than 90, pulse less than 60 or greater than 120, temperature greater than 38.5 degrees C

ONC NURSING COMMUNICATION 26

Interval: Until discontinued

Occurrences: --

Comments:

- 2) Infuse antibody via pump
- 3) If any of the following occurs: FEVER (temperature greater than 38.5 degrees Celsius), RIGORS, HYPOTENSION (30mm Hg decrease from baseline), and/or MUCOSAL CONGESTION / EDEMA, HOLD infusion until improvement of symptoms (When symptoms improve, resume infusion at HALF the previous rate)

Rituximab Infusion Reaction Orders

meperidine (DEMEROL) injection 25 mg

Dose: 25 mg

Route: intravenous

once PRN

Start: S

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg

Route: intravenous

once PRN

Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg

Route: intravenous

once PRN

famotidine (PEPCID) injection 20 mg

Dose: 20 mg

Route: intravenous

once PRN

Start: S

Rituximab Additional Orders

epINEPHrine (ADRENALIN) 1 mg/1 mL injection 0.3 mg

Dose: 0.3 mg

Route: intramuscular

once PRN

Start: S