

OP APML4 MAINTENANCE

Types: ONCOLOGY TREATMENT

Synonyms: ACUTE, PROMYEL, APL, ATRA, IDAR, ATO, ARSE, TRIO, ALL, TRANS, RETINO, MAIN

Take-Home Medications	Repeat 1 time	Cycle length: 1 day
Labs and Take-Home Medications Perform every 1 day x1		
Take-Home Medications Prior to Treatment		
tretinoin (VESANOID) 10 mg chemo capsule		
Dose: 22.5 mg/m2	Route: oral	2 times daily
Dispense: --	Refills: --	
Start: S		
Instructions: DO NOT CRUSH, CHEW, OR DISSOLVE CAPSULE. Administer with meals.		
Take-Home Medications Prior to Treatment		
methotrexate 2.5 MG tablet		
Dose: 5 mg/m2	Route: oral	weekly
Dispense: --	Refills: --	
Start: S+14	End: S+85	
Instructions: CYTOTOXIC AGENT DO NOT OPEN OR CRUSH. Swallow whole.		
Take-Home Medications Prior to Treatment		
mercaptopurine (PURINETHOL) 50 mg chemo tablet		
Dose: 50 mg/m2	Route: oral	daily
Dispense: --	Refills: --	
Start: S+14	End: S+90	
Comments: On days 1 through 14 of treatment in all cycles. Take a least 1 hour after evening meal without milk or citrus products.		
Instructions: Take a least 1 hour after evening meal without milk or citrus products.		
Provider Communication		
ONC PROVIDER COMMUNICATION 26		
Interval: --	Occurrences: --	
Comments:	Adjust oral methotrexate and mercaptopurine to target a neutrophil count of 1000 -2000.	