

## OP ATEZOLIZUMAB / NAB-PACLITAXEL EVERY 28 DAYS

*Types:* ONCOLOGY TREATMENT

*Synonyms:* BREAST, ATEZ, ATAZ, TEC, TAC, TECENTRIQ, ATEZOLIZUMAB, ABRAX, ABRAXANE, NAB, NAB-PACLITAXEL

Cycle 1	Repeat 1 time	Cycle length: 28 days
Day 1	Perform every 1 day x1	
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: -- Occurrences: --		
Labs		
CBC WITH PLATELET AND DIFFERENTIAL		
Interval: -- Occurrences: --		
COMPREHENSIVE METABOLIC PANEL		
Interval: -- Occurrences: --		
THYROID STIMULATING HORMONE		
Interval: -- Occurrences: --		
T3, FREE		
Interval: -- Occurrences: --		
T4, FREE		
Interval: -- Occurrences: --		
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
TREATMENT CONDITIONS 40		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6 mg/dL)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum Magnesium less than 1.0 mg/dL, give 2 gram magnesium sulfate IV and contact MD/NP		
o Serum Magnesium 1.0 to 1.2 mg/dL, give 2 gram magnesium sulfate IV		
o Serum Magnesium 1.3 to 1.5 mg/dL, give 1 gram magnesium sulfate IV		
o Serum Magnesium 1.6 mg/dL or greater, do not give magnesium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		

o Sign electrolyte replacement order as Per protocol: cosign required

#### Nursing Orders

##### TREATMENT CONDITIONS 7

Interval: -- Occurrences: --  
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

#### Nursing Orders

##### TREATMENT CONDITIONS 30

Interval: -- Occurrences: --  
Comments: HOLD and notify provider if AST or ALT GREATER than 3 times ULN or total bilirubin GREATER than 1.5 times ULN.

#### Line Flush

##### sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN  
Start: S

#### Nursing Orders

##### sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
Start: S  
Instructions: To keep vein open.

#### Chemotherapy

##### atezolizumab (TECENTRIQ) 840 mg in sodium chloride 0.9% 250 mL chemo IVPB

Dose: 840 mg Route: intravenous once over 60 Minutes for 1 dose  
Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ATEZOLIZUMAB 840 MG/14 ML (60 MG/ML) INTRAVENOUS SOLUTION	Medications	840 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	236 mL	Yes	Yes

#### Pre-Medications

##### ondansetron (ZOFTRAN) IV 8 mg

Dose: 8 mg Route: intravenous once for 1 dose  
Start: S

#### Chemotherapy

##### NAB-PAClitaxel (ABRAXANE) chemo infusion 100 mg/m2 (Treatment Plan)

Dose: 100 mg/m2 Route: intravenous once over 30 Minutes for 1 dose  
Offset: 1.5 Hours

#### Hematology & Oncology Hypersensitivity Reaction Standing Order

##### ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --  
Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)  
1. Stop the infusion.  
2. Place the patient on continuous monitoring.  
3. Obtain vital signs.  
4. Administer Normal Saline at 50 mL per hour using a new bag and new

intravenous tubing.

5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.

6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

7. Notify the treating physician.

8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 4**

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.

2. Notify the CERT team and treating physician immediately.

3. Place the patient on continuous monitoring.

4. Obtain vital signs.

5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.

6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.

7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 83**

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.

2. Notify the CERT team and treating physician immediately.

3. Place the patient on continuous monitoring.

4. Obtain vital signs.

5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.

6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.

7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.

8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.

9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.

10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**diphenhydrAMINE (BENADRYL) injection 25  
ma**

Dose: 25 mg                      Route: intravenous                      PRN  
Start: S

**fexofenadine (ALLEGRA) tablet 180 mg**  
Dose: 180 mg                      Route: oral                      PRN  
Start: S

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**  
Dose: 20 mg                      Route: intravenous                      PRN  
Start: S

**hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**  
Dose: 100 mg                      Route: intravenous                      PRN

**dexamethasone (DECADRON) injection 4 mg**  
Dose: 4 mg                      Route: intravenous                      PRN  
Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**  
Dose: 0.3 mg                      Route: subcutaneous                      PRN  
Start: S

Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --                      Occurrences: --  
Comments:                      Discontinue IV.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**  
Dose: 20 mL                      Route: intravenous                      PRN

☒ **HEParin, porcine (PF) injection 500 Units**  
Dose: 500 Units                      Route: intra-catheter                      once PRN  
Start: S  
Instructions:  
Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.

**Day 8**

Perform every 1 day x1

Appointment Requests

**INFUSION APPOINTMENT REQUEST**  
Interval: --                      Occurrences: --

Labs

**CBC WITH PLATELET AND DIFFERENTIAL**  
Interval: --                      Occurrences: --

Labs

**COMPREHENSIVE METABOLIC PANEL**  
Interval: --                      Occurrences: --

Outpatient Electrolyte Replacement Protocol

**TREATMENT CONDITIONS 39**

Interval: --                      Occurrences: --  
Comments:                      Potassium (Normal range 3.5 to 5.0mEq/L)  
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP  
o Protocol applies only to same day lab value.  
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP  
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO

- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

#### **TREATMENT CONDITIONS 40**

Interval: --

Occurrences: --

Comments:

Magnesium (Normal range 1.6 to 2.6 mg/dL)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0 mg/dL, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2 mg/dL, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5 mg/dL, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mg/dL or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

#### **Nursing Orders**

##### **TREATMENT CONDITIONS 7**

Interval: --

Occurrences: --

Comments:

HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

#### **Line Flush**

##### **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL

Route: intravenous

PRN

Start: S

#### **Nursing Orders**

##### **sodium chloride 0.9 % infusion 250 mL**

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

#### **Pre-Medications**

##### **ondansetron (ZOFTRAN) IV 8 mg**

Dose: 8 mg

Route: intravenous

once for 1 dose

Start: S

#### **Chemotherapy**

##### **NAB-PACLitaxel (ABRAXANE) chemo infusion 100 mg/m2 (Treatment Plan)**

Dose: 100 mg/m2

Route: intravenous

once over 30 Minutes for 1 dose

Offset: 30 Minutes

#### **Hematology & Oncology Hypersensitivity Reaction Standing Order**

##### **ONC NURSING COMMUNICATION 82**

Interval: --

Occurrences: --

Comments:

- Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
1. Stop the infusion.
  2. Place the patient on continuous monitoring.

3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 4**

Interval: --

Occurrences: --

Comments:

- Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)
1. Stop the infusion.
  2. Notify the CERT team and treating physician immediately.
  3. Place the patient on continuous monitoring.
  4. Obtain vital signs.
  5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
  6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
  7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
  8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
  9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 83**

Interval: --

Occurrences: --

Comments:

- Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)
1. Stop the infusion.
  2. Notify the CERT team and treating physician immediately.
  3. Place the patient on continuous monitoring.
  4. Obtain vital signs.
  5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
  6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
  7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
  8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
  9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
  10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**diphenhydramine (BENADRYL) injection 25 mg**

Dose: 25 mg                      Route: intravenous                      PRN  
Start: S

**fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg                      Route: oral                      PRN  
Start: S

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg                      Route: intravenous                      PRN  
Start: S

**hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**

Dose: 100 mg                      Route: intravenous                      PRN

**dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg                      Route: intravenous                      PRN  
Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**

Dose: 0.3 mg                      Route: subcutaneous                      PRN  
Start: S

## Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --                      Occurrences: --  
Comments:                      Discontinue IV.

## Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL                      Route: intravenous                      PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units                      Route: intra-catheter                      once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.

**Day 15**

Perform every 1 day x1

## Appointment Requests

**INFUSION APPOINTMENT REQUEST**

Interval: --                      Occurrences: --

## Labs

**CBC WITH PLATELET AND DIFFERENTIAL**

Interval: --                      Occurrences: --

**COMPREHENSIVE METABOLIC PANEL**

Interval: --                      Occurrences: --

**THYROID STIMULATING HORMONE**

Interval: --                      Occurrences: --

**T3, FREE**

Interval: --                      Occurrences: --

**T4, FREE**

Interval: --                      Occurrences: --

## Outpatient Electrolyte Replacement Protocol

**TREATMENT CONDITIONS 39**

Interval: --                      Occurrences: --

Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

#### TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

Comments: Magnesium (Normal range 1.6 to 2.6 mg/dL)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0 mg/dL, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2 mg/dL, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5 mg/dL, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mg/dL or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

#### Nursing Orders

##### TREATMENT CONDITIONS 7

Interval: -- Occurrences: --

Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

#### Nursing Orders

##### TREATMENT CONDITIONS 30

Interval: -- Occurrences: --

Comments: HOLD and notify provider if AST or ALT GREATER than 3 times ULN or total bilirubin GREATER than 1.5 times ULN.

#### Line Flush

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

Start: S

#### Nursing Orders

**sodium chloride 0.9 % infusion 250 mL**

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:  
To keep vein open.

#### Chemotherapy

**atezolizumab (TECENTRIQ) 840 mg in sodium chloride 0.9% 250 mL chemo IVPB**



Dose: 840 mg

Route: intravenous

once over 60 Minutes for 1 dose

Offset: 30 Minutes

**Ingredients:**

**Name**

**Type**

**Dose**

**Selected**

**Adds Vol.**

ATEZOLIZUMAB  
840 MG/14 ML (60  
MG/ML)  
INTRAVENOUS  
SOLUTION  
SODIUM  
CHLORIDE 0.9 %  
INTRAVENOUS  
SOLUTION

Medications

840 mg

Main

Yes

Ingredient

QS Base

236 mL

Yes

Yes

**Pre-Medications**

**ondansetron (ZOFTRAN) IV 8 mg**

Dose: 8 mg

Route: intravenous

once for 1 dose

Start: S

**Chemotherapy**

**NAB-PACLitaxel (ABRAXANE) chemo infusion**

**100 mg/m2 (Treatment Plan)**

Dose: 100 mg/m2

Route: intravenous

once over 30 Minutes for 1 dose

Offset: 1.5 Hours

**Hematology & Oncology Hypersensitivity Reaction Standing Order**

**ONC NURSING COMMUNICATION 82**

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**ONC NURSING COMMUNICATION 4**

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous). Fexofenadine 180 mg orally and Famotidine 20 mg

intravenous once.

8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.

2. Notify the CERT team and treating physician immediately.

3. Place the patient on continuous monitoring.

4. Obtain vital signs.

5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.

6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.

7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.

8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.

9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.

10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg

Route: intravenous

PRN

Start: S

#### fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg

Route: oral

PRN

Start: S

#### famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg

Route: intravenous

PRN

Start: S

#### hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg

Route: intravenous

PRN

#### dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg

Route: intravenous

PRN

Start: S

#### epinephrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg

Route: subcutaneous

PRN

Start: S

#### Discharge Nursing Orders

#### ONC NURSING COMMUNICATION 76

Interval: --

Occurrences: --

Comments:

Discontinue IV.

#### Discharge Nursing Orders

#### ☒ sodium chloride 0.9 % flush 20 mL

Dose: 20 mL

Route: intravenous

PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units      Route: intra-catheter      once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.