

OP AYA - COURSE 1 (REMISSION INDUCTION)

Types: ONCOLOGY TREATMENT

Synonyms: ALL, ACUTE, LYMPHO, ALL, INDUCT, PRO, PROMYEL, CALGB, REMISSION, INDUC, INTRATHECAL, COURSE 1, COURSE I, AYA

Take-Home Medications	Repeat 1 time	Cycle length: 1 day
Day 1		Perform every 1 day x1
Take-Home Medications Prior to Treatment		
allopurinol (ZYLOPRIM) 300 MG tablet		
Dose: 300 mg	Route: oral	daily
Dispense: --	Refills: --	
Start: S		
Instructions:		
Continue until peripheral blasts and extramedullary disease are reduced.		
Take-Home Medications Prior to Treatment		
predniSONE (DELTASONE) 20 MG tablet		
Dose: 30 mg/m2	Route: oral	2 times daily
Dispense: --	Refills: --	
Start: S		
Comments:		
On day 1 through day 14 of all cycles. Take after meals or with food or milk.		
Instructions:		
On day 1 through day 28. Take after meals or with food or milk. Do not taper.		
Cycle 1	Repeat 1 time	Cycle length: 29 days
Day 1		Perform every 1 day x1
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: --	Occurrences: --	
Labs		
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: --	Occurrences: --	
<input type="checkbox"/> LDH		
Interval: --	Occurrences: --	
<input type="checkbox"/> URIC ACID LEVEL		
Interval: --	Occurrences: --	
Labs		
<input checked="" type="checkbox"/> LIPASE LEVEL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> AMYLASE LEVEL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> FIBRINOGEN		

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

- Comments: Potassium (Normal range 3.5 to 5.0mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
 - o Protocol applies only to same day lab value.
 - o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
 - o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
 - o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
 - o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
 - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
 - o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

- Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
 - o Protocol applies only to same day lab value.
 - o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
 - o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
 - o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
 - o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
 - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
 - o Sign electrolyte replacement order as Per protocol: cosign required

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg in sodium chloride 0.9% 50 mL IVPB

Dose: -- Route: intravenous once over 15 Minutes for 1 dose
Start: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
	DEXAMETHASONE 4 MG/ML	Medications	12 mg	No	No

INJECTION SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

ondansetron (ZOFTRAN) injection 8 mg

Dose: 8 mg Route: intravenous once for 1 dose
Start: S

ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S

dexamethasone (DECADRON) tablet 12 mg

Dose: 12 mg Route: oral once for 1 dose
Start: S

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

LORAZepam (ATIVAN) tablet 1 mg

Dose: 1 mg Route: oral once for 1 dose
Start: S

LORAZepam (ATIVAN) injection 1 mg

Dose: 1 mg Route: intravenous once for 1 dose
Start: S

Provider Communication

ONC PROVIDER COMMUNICATION 28

Interval: -- Occurrences: --
Comments: Voriconazole and posaconazole are contra-indicated with vincristine.

Chemotherapy

vinCRISTine (ONCOVIN) 1.5 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 1.5 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
Start: S

Instructions:
DRUG IS A VESICANT. FATAL IF GIVEN
INTRATHECALLY. Maximum dose = 2 mg.

Rule-Based Template: RULE ONCBCN
 VINCRISTINE 1.5 MG/M2

Conditions:
 BSA < 1.33 m2
 BSA >= 1.33 m2

Modifications:
 Set dose to 1.5 mg/m2
 Set dose to 2 mg

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	VINCRISTINE 1 MG/ML	Medications	1.5 mg/m2	Main Ingredient	Yes
	INTRAVENOUS SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	50 mL	Yes	Yes

Intrathecal Injections

cytarabine PF (CYSTOSAR) 70 mg in sodium chloride 0.9% 5 mL chemo PF INTRATHECAL injection

Dose: 70 mg Route: intrathecal once over 5 Minutes for 1 dose
 Offset: 30 Minutes

Instructions:
 HAZARDOUS - Handle with care. Administer on Day 1.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	CYTARABINE (PF) 100 MG/5 ML (20 MG/ML) INJECTION SOLUTION	Medications	70 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	1.5 mL	Yes	Yes

Chemotherapy

DAUNORubicin (CERUBIDINE) 25 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 25 mg/m2 Route: intravenous once over 15 Minutes for 1 dose
 Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DAUNORUBICIN 20 MG INTRAVENOUS SOLUTION	Medications	25 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	100 mL	No	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --
 Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
 1. Stop the infusion.
 2. Place the patient on continuous monitoring.
 3. Obtain vital signs.
 4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
 5. If greater than or equal to 30 minutes since the last dose of

Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.

6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

7. Notify the treating physician.

8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.

2. Notify the CERT team and treating physician immediately.

3. Place the patient on continuous monitoring.

4. Obtain vital signs.

5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.

6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.

7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.

2. Notify the CERT team and treating physician immediately.

3. Place the patient on continuous monitoring.

4. Obtain vital signs.

5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.

6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.

7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.

8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.

9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.

10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg

Route: intravenous

PRN

Start: S

fexofenadine (ALLEGRA) tablet 180 mgDose: 180 mg Route: oral PRN
Start: S**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**Dose: 20 mg Route: intravenous PRN
Start: S**hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mgDose: 4 mg Route: intravenous PRN
Start: S**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

 sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

 HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.**Day 4**

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

 CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

 COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

 MAGNESIUM LEVEL

Interval: -- Occurrences: --

 LDH

Interval: -- Occurrences: --

 URIC ACID LEVEL

Interval: -- Occurrences: --

Labs

 LIPASE LEVEL

Interval: -- Occurrences: --

AMYLASE LEVEL

Interval: -- Occurrences: --

FIBRINOGEN

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

- Comments: Potassium (Normal range 3.5 to 5.0mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
 - o Protocol applies only to same day lab value.
 - o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
 - o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
 - o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
 - o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
 - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
 - o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

- Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
 - o Protocol applies only to same day lab value.
 - o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
 - o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
 - o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
 - o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
 - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
 - o Sign electrolyte replacement order as Per protocol: cosign required

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

acetaminophen (TYLENOL) tablet 650 mg

Dose: 650 mg Route: oral once for 1 dose

Instructions:

Administer 30 minutes prior to Pegaspargase.

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous once for 1 dose
Start: S
Instructions:
Administer 30 minutes prior to Pegaspargase.

diphenhydrAMINE (BENADRYL) tablet 25 mg

Dose: 25 mg Route: oral once for 1 dose

Instructions:
Administer 30 minutes prior to Pegaspargase.

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous once for 1 dose

Instructions:
Administer 30 minutes prior to Pegaspargase.

Nursing Orders

TREATMENT CONDITIONS 28

Interval: -- Occurrences: --
Comments: MD must review results of amylase, lipase, and fibrinogen prior to patient receiving peg-asparaginase.

Chemotherapy

pegaspargase (ONCASPAR) 2,500 Units/m2 in dextrose 5% 100 mL chemo IVPB

Dose: 2,500 Units/m2 Route: intravenous once over 2 Hours for 1 dose
Offset: 30 Minutes

Instructions:
Infuse into a flowing IV line. Observe for 1 hour post infusion

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PEGASPARGASE 750 UNIT/ML INJECTION SOLUTION	Medications	2,500 Units/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	100 mL	No	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	100 mL	Yes	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --
Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine,

administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

7. Notify the treating physician.

8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.

2. Notify the CERT team and treating physician immediately.

3. Place the patient on continuous monitoring.

4. Obtain vital signs.

5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.

6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.

7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.

2. Notify the CERT team and treating physician immediately.

3. Place the patient on continuous monitoring.

4. Obtain vital signs.

5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.

6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.

7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.

8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.

9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.

10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg

Route: intravenous

PRN

Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg

Route: oral

PRN

Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Day 8

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

LDH

Interval: -- Occurrences: --

URIC ACID LEVEL

Interval: -- Occurrences: --

Labs

LIPASE LEVEL

Interval: -- Occurrences: --

AMYLASE LEVEL

Interval: -- Occurrences: --

FIBRINOGEN

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Provider Communication

ONC PROVIDER COMMUNICATION 28

Interval: -- Occurrences: --

Comments: Voriconazole and posaconazole are contra-indicated with vincristine.

Chemotherapy

vinCRISTine (ONCOVIN) 1.5 mg/m2 in sodium

chloride 0.9 % 50 mL chemo IVPB

Dose: 1.5 mg/m2 Route: intravenous once over 10 Minutes for 1 dose

Start: S

Instructions:

DRUG IS A VESICANT. FATAL IF GIVEN INTRATHECALLY. Maximum dose = 2 mg.

Rule-Based Template: RULE ONCBCN

VINCRISTINE 1.5 MG/M2

Conditions:

BSA < 1.33 m2

BSA >= 1.33 m2

Modifications:

Set dose to 1.5 mg/m2

Set dose to 2 mg

Ingredients:**Name****Type****Dose****Selected****Adds Vol.**VINCRISTINE 1
MG/ML
INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Medications

1.5
mg/m2Main
Ingredient

QS Base

50 mL

Yes Yes

Intrathecal Injections

methotrexate PF 15 mg in sodium chloride 0.9% 5 mL chemo PF INTRATHECAL injection

Dose: 15 mg Route: intrathecal once over 5 Minutes for 1 dose

Start: S End: S

Instructions:

INTRATHECAL VIA DIRECT LUMBAR PUNCTURE. Total volume with preservative free sodium chloride 0.9% for intrathecal is 3-5 mL. This dose is NOT to be combined with cytarabine. To be used as single agent chemotherapy only. May be combined with hydrocortisone, but may not be combined with another chemotherapy agent. Patient should remain in horizontal position for 30 minutes.

Ingredients:**Name****Type****Dose****Selected****Adds Vol.**METHOTREXATE
SODIUM (PF) 25
MG/ML INJECTION
SOLUTION
SODIUM
CHLORIDE 0.9 %
INJECTION
SOLUTION

Medications

15 mg

Main
Ingredient

QS Base

4.4 mL

Yes Yes

ONC PROVIDER COMMUNICATION 29

Interval: -- Occurrences: --

Comments: Send CSF for cell count and cytology with every intrathecal administration.

Chemotherapy

DAUNORubicin (CERUBIDINE) 25 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 25 mg/m2 Route: intravenous once over 15 Minutes for 1 dose

Offset: 30 Minutes

Ingredients:**Name****Type****Dose****Selected****Adds Vol.**DAUNORUBICIN 20
MG INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %

Medications

25 mg/m2

Main
Ingredient

QS Base

100 mL

Yes Yes

INTRAVENOUS
SOLUTION

DEXTROSE 5 % IN QS Base 100 mL No Yes
WATER (D5W)
INTRAVENOUS
SOLUTION

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.

5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25

mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Day 15

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

LDH

Interval: -- Occurrences: --

URIC ACID LEVEL

Interval: -- Occurrences: --

Labs

LIPASE LEVEL

Interval: -- Occurrences: --

AMYLASE LEVEL

Interval: -- Occurrences: --

FIBRINOGEN

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Provider Communication

ONC PROVIDER COMMUNICATION 30

Interval: -- Occurrences: --
 Comments: Bone marrow biopsy and aspirate specimen must be obtained for all patients on Day 15 to assess initial response and on Day 29 to assess induction response and minimal residual disease.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions: To keep vein open.

Provider Communication

ONC PROVIDER COMMUNICATION 28

Interval: -- Occurrences: --
 Comments: Voriconazole and posaconazole are contra-indicated with vincristine.

Chemotherapy

vinCRISTine (ONCOVIN) 1.5 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 1.5 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
 Start: S

Instructions: DRUG IS A VESICANT. FATAL IF GIVEN INTRATHECALLY. Maximum dose = 2 mg.

Rule-Based Template: RULE ONCBCN

VINCRIStINE 1.5 MG/M2

Conditions:

BSA < 1.33 m2
 BSA >= 1.33 m2

Modifications:

Set dose to 1.5 mg/m2
 Set dose to 2 mg

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	VINCRIStINE 1 MG/ML INTRAVENOUS SOLUTION	Medications	1.5 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	50 mL	Yes	Yes

Chemotherapy

DAUNORubicin (CERUBIDINE) 25 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 25 mg/m2 Route: intravenous once over 15 Minutes for 1 dose
 Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DAUNORUBICIN 20 MG INTRAVENOUS SOLUTION	Medications	25 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes
	DEXTROSE 5 % IN	QS Base	100 mL	No	Yes

WATER (D5W)
INTRAVENOUS
SOLUTION

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --
Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: -- Occurrences: --
Comments: Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: -- Occurrences: --
Comments: Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to

maintain O2 saturation of greater than or equal to 92%.
 7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
 8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
 9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
 10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
 Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
 Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
 Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
 Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
 Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
 Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
 Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Day 22

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

LDH

Interval: -- Occurrences: --

URIC ACID LEVEL

Interval: -- Occurrences: --

Labs

LIPASE LEVEL

Interval: -- Occurrences: --

AMYLASE LEVEL

Interval: -- Occurrences: --

FIBRINOGEN

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

Comments:

Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

Comments:

Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S

Nursing Orders**sodium chloride 0.9 % infusion 250 mL**

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions:
 To keep vein open.

Provider Communication**ONC PROVIDER COMMUNICATION 28**

Interval: -- Occurrences: --
 Comments: Voriconazole and posaconazole are contra-indicated with vincristine.

Chemotherapy**vinCRISTine (ONCOVIN) 1.5 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB**

Dose: 1.5 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
 Start: S

Instructions:
 DRUG IS A VESICANT. FATAL IF GIVEN INTRATHECALLY. Maximum dose = 2 mg.

Rule-Based Template: RULE ONCBCN
 VINCRISTINE 1.5 MG/M2

Conditions:
 BSA < 1.33 m2
 BSA >= 1.33 m2

Modifications:
 Set dose to 1.5 mg/m2
 Set dose to 2 mg

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	VINCRISTINE 1 MG/ML	Medications	1.5 mg/m2	Main Ingredient	Yes
	INTRAVENOUS SOLUTION				
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	50 mL	Yes	Yes

Chemotherapy**DAUNORubicin (CERUBIDINE) 25 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB**

Dose: 25 mg/m2 Route: intravenous once over 15 Minutes for 1 dose
 Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DAUNORUBICIN 20 MG INTRAVENOUS SOLUTION	Medications	25 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	100 mL	No	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order**ONC NURSING COMMUNICATION 82**

Interval: -- Occurrences: --
 Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
 1. Stop the infusion.
 2. Place the patient on continuous monitoring.

3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fenofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Day 29

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

LDH

Interval: -- Occurrences: --

URIC ACID LEVEL

Interval: -- Occurrences: --

Labs

LIPASE LEVEL

Interval: -- Occurrences: --

AMYLASE LEVEL

Interval: -- Occurrences: --

FIBRINOGEN

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --

Comments:

- Potassium (Normal range 3.5 to 5.0mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
 - o Protocol applies only to same day lab value.
 - o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
 - o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
 - o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
 - o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
 - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
 - o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

Comments:

- Magnesium (Normal range 1.6 to 2.6mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
 - o Protocol applies only to same day lab value.
 - o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
 - o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
 - o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
 - o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
 - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
 - o Sign electrolyte replacement order as Per protocol: cosign required

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Intrathecal Injections

methotrexate PF 15 mg in sodium chloride 0.9% 5 mL chemo PF INTRATHECAL injection

Dose: 15 mg Route: intrathecal once over 5 Minutes for 1 dose
Start: S End: S

Instructions:

INTRATHECAL VIA DIRECT LUMBAR PUNCTURE. Total volume with preservative

free sodium chloride 0.9% for intrathecal is 3-5 mL. This dose is NOT to be combined with cytarabine. To be used as single agent chemotherapy only. May be combined with hydrocortisone, but may not be combined with another chemotherapy agent. Patient should remain in horizontal position for 30 minutes.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	METHOTREXATE SODIUM (PF) 25 MG/ML INJECTION SOLUTION	Medications	15 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INJECTION SOLUTION	QS Base	4.4 mL	Yes	Yes

ONC PROVIDER COMMUNICATION 29

Interval: -- Occurrences: --
 Comments: Send CSF for cell count and cytology with every intrathecal administration.

Provider Communication

ONC PROVIDER COMMUNICATION 30

Interval: -- Occurrences: --
 Comments: Bone marrow biopsy and aspirate specimen must be obtained for all patients on Day 15 to assess initial response and on Day 29 to assess induction response and minimal residual disease.

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --
 Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
 1. Stop the infusion.
 2. Place the patient on continuous monitoring.
 3. Obtain vital signs.
 4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
 5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
 6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
 7. Notify the treating physician.
 8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
 9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: -- Occurrences: --
 Comments: Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)
 1. Stop the infusion.
 2. Notify the CERT team and treating physician immediately.
 3. Place the patient on continuous monitoring.
 4. Obtain vital signs.

5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epinephrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.