

OP CARBOPLATIN / ETOPOSIDE (EVERY 21 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: NSCLC, NON, ETOP, CARBO, SMALL, LUNG, SCLC, PARA

Cycles 1 to 4	Repeat 4 times	Cycle length: 21 days
Day 1		Perform every 1 day x1
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: -- Occurrences: --		
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: -- Occurrences: --		
<input type="checkbox"/> URINALYSIS, AUTOMATED WITH MICROSCOPY		
Interval: -- Occurrences: --		
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
TREATMENT CONDITIONS 40		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP		
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV		
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV		
o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		

o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL
Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL
Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg, dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB
Dose: -- Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:30 AM
Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Yes	No
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

ondansetron (ZOFTRAN) tablet 16 mg
Dose: 16 mg Route: oral once for 1 dose
Start: S End: S 11:30 AM

dexamethasone (DECADRON) tablet 12 mg
Dose: 12 mg Route: oral once for 1 dose
Start: S

palonosetron (ALOXI) injection 0.25 mg
Dose: 250 mcg Route: intravenous once for 1 dose
Start: S End: S 3:00 PM
Instructions:
For OUTPATIENT use only.

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB
Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION				
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Pre-Medications

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
 Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION				
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Pre-Medications

netupitant-palonosetron (AKYNZEO) 300-0.5 mg 1 capsule

Dose: 1 capsule Route: oral once for 1 dose
 Start: S End: S 5:30 PM

Instructions:
 Administer approximately 1 hour prior to chemotherapy.

Chemotherapy

etoposide (TOPOSAR) 100 mg/m2 in sodium chloride (NON-PVC) 0.9 % 500 mL chemo IVPB

Dose: 100 mg/m2 Route: intravenous once over 1 Hours for 1 dose
 Offset: 30 Minutes

Instructions:
 Administer through a 0.22 micron filter and non-PVC tubing set.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ETOPOSIDE 20 MG/ML	Medications	100 mg/m2	Main Ingredient	Yes
	INTRAVENOUS SOLUTION				
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	QS Base	500 mL	Yes	Yes

Chemotherapy

CARBOplatin (PARAplatin) in dextrose 5% 250 mL chemo IVPB

AUC: 6 Use AUC Route: intravenous once over 60 Minutes for 1 dose

Ingredients:	Name	Offset: 1.5 Hours		Selected	Adds Vol.
		Type	Dose		
	CARBOPLATIN 10 MG/ML INTRAVENOUS SOLUTION	Medications		Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --
 Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
 1. Stop the infusion.
 2. Place the patient on continuous monitoring.
 3. Obtain vital signs.
 4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
 5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
 6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
 7. Notify the treating physician.
 8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
 9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: -- Occurrences: --
 Comments: Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)
 1. Stop the infusion.
 2. Notify the CERT team and treating physician immediately.
 3. Place the patient on continuous monitoring.
 4. Obtain vital signs.
 5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
 6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
 7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
 8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
 9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: -- Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device

maintenance.

Day 2

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

ondansetron (ZOFRAN) injection 8 mg

Dose: 8 mg Route: intravenous once for 1 dose
Start: S End: S 11:15 AM

ondansetron (ZOFRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S

ondansetron (ZOFRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Chemotherapy

etoposide (TOPOSAR) 100 mg/m2 in sodium chloride (NON-PVC) 0.9 % 500 mL chemo IVPB

Dose: 100 mg/m2 Route: intravenous once over 1 Hours for 1 dose
Offset: 30 Minutes

Instructions:
Administer through a 0.22 micron filter and non-PVC tubing set.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ETOPOSIDE 20 MG/ML INTRAVENOUS SOLUTION	Medications	100 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	QS Base	500 mL	Yes	Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Day 3

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

ondansetron (ZOFRAN) injection 8 mg

Dose: 8 mg Route: intravenous once for 1 dose

Start: S End: S 11:15 AM

ondansetron (ZOFRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose

Start: S

ondansetron (ZOFRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose

Start: S End: S 11:00 AM

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Chemotherapy

etoposide (TOPOSAR) 100 mg/m2 in sodium chloride (NON-PVC) 0.9 % 500 mL chemo IVPB

Dose: 100 mg/m2 Route: intravenous once over 1 Hours for 1 dose
Offset: 30 Minutes

Instructions:

Administer through a 0.22 micron filter and non-PVC tubing set.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ETOPOSIDE 20 MG/ML INTRAVENOUS SOLUTION	Medications	100 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	QS Base	500 mL	Yes	Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Post-Medications

**pegfilgrastim (NEULASTA) on-body injection
kit 6 mg**

Dose: 6 mg Route: subcutaneous once for 1 dose

Start: S

End: S

Instructions:

Apply to intact, nonirritated skin on the back of
the arm or abdomen (only use the back of the
arm if caregiver is available to monitor
On-body injection status).