

OP CISPLATIN / PEMETREXED W/ CONCURRENT THORACIC RT

Types: ONCOLOGY TREATMENT

Synonyms: PEMETREXED, PEM, ALIMTA, CISPLATIN, PLATINOL, CIS, PLAT, LUNG, NSCLC, NON-SMALL CELL, ALIM

Cycle 1	Repeat 1 time	Cycle length: 21 days
Day 1	Perform every 1 day x1	
Appointment Requests		
<input type="checkbox"/> INFUSION APPOINTMENT REQUEST	Interval: --	Occurrences: --
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL	Interval: --	Occurrences: --
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL	Interval: --	Occurrences: --
<input checked="" type="checkbox"/> MAGNESIUM LEVEL	Interval: --	Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

ONC NURSING COMMUNICATION 80

Interval: -- Occurrences: --
Comments: Confirm that patient has started taking Folic Acid 350-1000mcg PO daily beginning 1-2 weeks prior to the first dose of pemetrexed and continuing until 3 weeks after the last dose of pemetrexed.

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Pre-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL Route: intravenous once @ 500 mL/hr for 1 dose
Start: S

Pre-Medications

palonosetron (ALOXI) injection 0.25 mg

Dose: 0.25 mg Route: intravenous once for 1 dose
Offset: 1 Hours

dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% IVPB

Dose: 12 mg Route: intravenous once over 15 Minutes for 1 dose
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

netupitant-palonosetron (AKYNZEO) 300-0.5 mg per capsule 1 capsule

Dose: 1 capsule Route: oral once for 1 dose
Start: S End: S 5:30 PM
Instructions:
Administer approximately 1 hour prior to chemotherapy.

Chemotherapy

PEMEtrexed (ALIMTA) 500 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 500 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PEMETREXED 500 MG INTRAVENOUS SOLUTION	Medications	500 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes

CISplatin (PLATINOL) 75 mg/m2 in sodium chloride 0.9 % 500 mL chemo IVPB

Dose: 75 mg/m2

Route: intravenous

once over 2 Hours for 1 dose

Offset: 2 Hours

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

CISPLATIN 1
MG/ML
INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION
DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Medications

QS Base

QS Base

75 mg/m2

500 mL

Main
Ingredient

Yes

No

Yes

Yes

Post-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL

Route: intravenous

once @ 500 mL/hr for 1 dose

Offset: 3 Hours

Instructions:

Following chemotherapy.

Supportive Care

cyanocobalamin injection 1,000 mcg

Dose: 1,000 mcg

Route: intramuscular

once for 1 dose

Start: S

End: S 2:30 PM

Instructions:

Begin week 1 of Pemetrexed dose, then every
9 weeks thereafter.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: --

Occurrences: --

Comments:

Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL

Route: intravenous

PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units

Route: intra-catheter

once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Cycles 2,3

Repeat 2 times

Cycle length: 21 days

Day 1

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: --

Occurrences: --

Labs

COMPREHENSIVE METABOLIC PANEL

Interval: --

Occurrences: --

CBC WITH PLATELET AND DIFFERENTIAL

Interval: --

Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

ONC NURSING COMMUNICATION 80

Interval: -- Occurrences: --
Comments: Confirm that patient has started taking Folic Acid 350-1000mcg PO daily beginning 1-2 weeks prior to the first dose of pemetrexed and continuing until 3 weeks after the last dose of pemetrexed.

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL
Dose: 20 mL Route: intravenous PRN
Start: S

Pre-Hydration

sodium chloride 0.9 % infusion 1,000 mL
Dose: 1,000 mL Route: intravenous once @ 500 mL/hr for 1 dose

Start: S

Pre-Medications

palonosetron (ALOXI) injection 0.25 mg

Dose: 0.25 mg Route: intravenous once for 1 dose
Offset: 1 Hours

dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% IVPB

Dose: 12 mg Route: intravenous once over 15 Minutes for 1 dose
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

netupitant-palonosetron (AKYNZEO) 300-0.5 mg per capsule 1 capsule

Dose: 1 capsule Route: oral once for 1 dose
Start: S End: S 5:30 PM
Instructions:
Administer approximately 1 hour prior to chemotherapy.

Chemotherapy

PEMEtrexed (ALIMTA) 500 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 500 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PEMETREXED 500 MG INTRAVENOUS SOLUTION	Medications	500 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes

CISplatin (PLATINOL) 75 mg/m2 in sodium chloride 0.9 % 500 mL chemo IVPB

Dose: 75 mg/m2 Route: intravenous once over 2 Hours for 1 dose
Offset: 2 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	CISPLATIN 1 MG/ML INTRAVENOUS SOLUTION	Medications	75 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

Post-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL Route: intravenous once @ 500 mL/hr for 1 dose
Offset: 3 Hours

Instructions:
Following chemotherapy.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Cycle 4

Repeat 1 time

Cycle length: 21 days

Day 1

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --
 Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --
 Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

ONC NURSING COMMUNICATION 80

Interval: -- Occurrences: --
 Comments: Confirm that patient has started taking Folic Acid 350-1000mcg PO daily beginning 1-2 weeks prior to the first dose of pemetrexed and continuing until 3 weeks after the last dose of pemetrexed.

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL
 Dose: 20 mL Route: intravenous PRN
 Start: S

Pre-Hydration

sodium chloride 0.9 % infusion 1,000 mL
 Dose: 1,000 mL Route: intravenous once @ 500 mL/hr for 1 dose
 Start: S

Pre-Medications

palonosetron (ALOXI) injection 0.25 mg
 Dose: 0.25 mg Route: intravenous once for 1 dose

Offset: 1 Hours

dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% IVPB

Dose: 12 mg Route: intravenous once over 15 Minutes for 1 dose
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

netupitant-palonosetron (AKYNZEO) 300-0.5 mg per capsule 1 capsule

Dose: 1 capsule Route: oral once for 1 dose
Start: S End: S 5:30 PM
Instructions:
Administer approximately 1 hour prior to chemotherapy.

Chemotherapy

PEMExred (ALIMTA) 500 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 500 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PEMETREXED 500 MG INTRAVENOUS SOLUTION	Medications	500 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes

CISplatin (PLATINOL) 75 mg/m2 in sodium chloride 0.9 % 500 mL chemo IVPB

Dose: 75 mg/m2 Route: intravenous once over 2 Hours for 1 dose
Offset: 2 Hours

Inredients:	Name	Type	Dose	Selected	Adds Vol.
-------------	------	------	------	----------	-----------

CISPLATIN 1 MG/ML INTRAVENOUS SOLUTION	Medications	75 mg/m2	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

Post-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL Route: intravenous once @ 500 mL/hr for 1 dose
Offset: 3 Hours

Instructions:
Following chemotherapy.

Supportive Care

cyanocobalamin injection 1,000 mcg

Dose: 1,000 mcg Route: intramuscular once for 1 dose
Start: S End: S 2:30 PM

Instructions:
Begin week 1 of Pemetrexed dose, then every
9 weeks thereafter.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:
Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Cycles 5,6

Repeat 2 times

Cycle length: 21 days

Day 1

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

ONC NURSING COMMUNICATION 80

Interval: -- Occurrences: --
Comments: Confirm that patient has started taking Folic Acid 350-1000mcg PO daily beginning 1-2 weeks prior to the first dose of pemetrexed and continuing until 3 weeks after the last dose of pemetrexed.

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Pre-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL Route: intravenous once @ 500 mL/hr for 1 dose
Start: S

Pre-Medications

palonosetron (ALOXI) injection 0.25 mg

Dose: 0.25 mg Route: intravenous once for 1 dose
Offset: 1 Hours

dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% IVPB

Dose: 12 mg Route: intravenous once over 15 Minutes for 1 dose
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

netupitant-palonosetron (AKYNZEO) 300-0.5 mg per capsule 1 capsule

Dose: 1 capsule Route: oral once for 1 dose
Start: S End: S 5:30 PM
Instructions:
Administer approximately 1 hour prior to chemotherapy.

Chemotherapy

PEMEtrexed (ALIMTA) 500 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 500 mg/m2 Route: intravenous once over 10 Minutes for 1 dose
Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	PEMETREXED 500 MG INTRAVENOUS SOLUTION	Medications	500 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes

CISplatin (PLATINOL) 75 mg/m2 in sodium chloride 0.9 % 500 mL chemo IVPB

Dose: 75 mg/m2

Route: intravenous

once over 2 Hours for 1 dose

Offset: 2 Hours

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

CISPLATIN 1

Medications

75 mg/m2

Main

Yes

MG/ML

Ingredient

INTRAVENOUS

SOLUTION

SODIUM

QS Base

500 mL

Yes

Yes

CHLORIDE 0.9 %

INTRAVENOUS

SOLUTION

DEXTROSE 5 % IN

QS Base

No

Yes

WATER (D5W)

INTRAVENOUS

SOLUTION

Post-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL

Route: intravenous

once @ 500 mL/hr for 1 dose

Offset: 3 Hours

Instructions:

Following chemotherapy.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: --

Occurrences: --

Comments:

Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL

Route: intravenous

PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units

Route: intra-catheter

once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.