

## OP CISPLATIN W/ CONCURRENT RT (EVERY 7 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: CIS, PLAT, CISPLATIN, PLATINOL, CONCURRENT RT, WEEKLY

<b>Cycles 1,2</b>	Repeat 2 times	Cycle length: 42 days
<b>Days 1,8,15,22,29,36</b>	Perform every 7 days x6	
Appointment Requests		
<b>INFUSION APPOINTMENT REQUEST</b>		
Interval: -- Occurrences: --		
Labs		
<input checked="" type="checkbox"/> <b>COMPREHENSIVE METABOLIC PANEL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>CBC WITH PLATELET AND DIFFERENTIAL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>MAGNESIUM LEVEL</b>		
Interval: -- Occurrences: --		
Outpatient Electrolyte Replacement Protocol		
<b>TREATMENT CONDITIONS 39</b>		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
<b>TREATMENT CONDITIONS 40</b>		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP		
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV		
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV		
o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		

Nursing Orders

**TREATMENT CONDITIONS 7**

Interval: -- Occurrences: --  
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN  
 Start: S

Pre-Hydration

**sodium chloride 0.9 % infusion 1,000 mL**

Dose: 1,000 mL Route: intravenous once @ 500 mL/hr for 1 dose  
 Start: S

Pre-Medications

**palonosetron (ALOXI) injection 0.25 mg**

Dose: 0.25 mg Route: intravenous once for 1 dose  
 Start: S End: S 1:45 PM

**dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% IVPB**

Dose: 12 mg Route: intravenous once over 15 Minutes for 1 dose  
 Start: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

**aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose  
 Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

**netupitant-palonosetron (AKYNZEO) 300-0.5 mg per capsule 1 capsule**

Dose: 1 capsule Route: oral once for 1 dose  
 Start: S End: S 5:30 PM

Instructions:  
 Administer approximately 1 hour prior to chemotherapy.

Chemotherapy

**CISplatin (PLATINOL) 40 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB**

Dose: 40 mg/m2      Route: intravenous      once over 1 Hours for 1 dose  
Offset: 2 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	CISPLATIN 1 MG/ML	Medications	40 mg/m2	Main Ingredient	Yes
	INTRAVENOUS SOLUTION SODIUM CHLORIDE 0.9 %	QS Base		Yes	Yes
	INTRAVENOUS SOLUTION DEXTROSE 5 % IN WATER (D5W)	QS Base		No	Yes
	INTRAVENOUS SOLUTION				

Post-Hydration

**sodium chloride 0.9 % infusion 1,000 mL**

Dose: 1,000 mL      Route: intravenous      once @ 500 mL/hr for 1 dose  
Offset: 3 Hours

Instructions:  
Following chemotherapy.

Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --      Occurrences: --  
Comments:      Discontinue IV.

Discharge Nursing Orders

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL      Route: intravenous      PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units      Route: intra-catheter      once PRN

Start: S  
Instructions:  
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.