

# OP DARATUMUMAB

Types: ONCOLOGY TREATMENT

Synonyms: MULTIPLE, MYELOMA, DARA, DAZALEX, DARZ, DARS

<b>Take-Home Medications</b>	Repeat 1 time	Cycle length: 1 day
<b>Day 1</b>		Perform every 1 day x1
Take-Home Medications Prior to Treatment		
<b>valACYclovir (VALTREX) 500 MG tablet</b>		
Dose: 500 mg	Route: oral	daily
Dispense: --	Refills: --	
Start: S	End: S+90	
Instructions: For shingles prophylaxis: continue for 3 months following treatment.		

<b>Weeks 1 - 8</b>	Repeat 1 time	Cycle length: 56 days
<b>Day 1</b>		Perform every 1 day x1
Appointment Requests		
<b>INFUSION APPOINTMENT REQUEST</b>		
Interval: --	Occurrences: --	
Labs		
<input checked="" type="checkbox"/> <b>CBC WITH PLATELET AND DIFFERENTIAL</b>		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> <b>COMPREHENSIVE METABOLIC PANEL</b>		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> <b>MAGNESIUM LEVEL</b>		
Interval: --	Occurrences: --	
<input type="checkbox"/> <b>LDH</b>		
Interval: --	Occurrences: --	
<input type="checkbox"/> <b>URIC ACID LEVEL</b>		
Interval: --	Occurrences: --	
Labs		
<b>TYPE AND SCREEN</b>		
Interval: --	Occurrences: --	
Comments:	Draw type and screen BEFORE each infusion.	
Outpatient Electrolyte Replacement Protocol		
<b>TREATMENT CONDITIONS 39</b>		
Interval: --	Occurrences: --	
Comments:	Potassium (Normal range 3.5 to 5.0mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
	o Protocol applies only to same day lab value.	
	o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP	
	o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO	
	o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO	
	o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement	
	o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"	
	o Sign electrolyte replacement order as Per protocol: cosign	

required

**TREATMENT CONDITIONS 40**

Interval: -- Occurrences: --  
 Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

**Nursing Orders**

**ONC NURSING COMMUNICATION 73**

Interval: -- Occurrences: --  
 Comments: Draw type and screen BEFORE each infusion.

**Line Flush**

**sodium chloride 0.9 % flush 20 mL**  
 Dose: 20 mL Route: intravenous PRN  
 Start: S

**Nursing Orders**

**sodium chloride 0.9 % infusion 250 mL**  
 Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
 Start: S  
 Instructions:  
 To keep vein open.

**Pre-Medications**

**methylPREDNISolone sodium succinate (Solu-MEDROL) injection 100 mg**  
 Dose: 100 mg Route: intravenous once for 1 dose  
 Start: S  
 Instructions:  
 Administer 30 minutes prior to chemotherapy.

**dexamethasone (DECADRON) 40 mg in sodium chloride 0.9 % IVPB**  
 Dose: 40 mg Route: intravenous once over 30 Minutes for 1 dose  
 Start: S  
 Instructions:  
 Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	40 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN	Base	50 mL	No	Yes

WATER (D5W)  
INTRAVENOUS  
SOLUTION

Pre-Medications

**acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg      Route: oral      once for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

Pre-Medications

**diphenhydramine (BENADRYL) tablet 25 mg**

Dose: 25 mg      Route: oral      once for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

Chemotherapy

**daratumumab 16 mg/kg in sodium chloride 0.9 % 1,000 mL IVPB**

Dose: 16 mg/kg      Route: intravenous      once @ 50 mL/hr for 1 dose  
Offset: 30 Minutes

Instructions:

First infusion (1,000 mL volume): Infuse at 50 mL/hour for the first hour. If no infusion reactions occur, may increase the rate by 50 mL/hour every hour (maximum rate: 200 mL/hour).

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DARATUMUMAB 20 MG/ML INTRAVENOUS SOLUTION	Medications	16 mg/kg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base		Yes	Yes

Delayed infusion reactions

**methylPREDNISolone sodium succinate (Solu-MEDROL) injection 20 mg**

Dose: 20 mg      Route: intravenous      once for 1 dose

Start: S

Hematology & Oncology Hypersensitivity Reaction Standing Order

**ONC NURSING COMMUNICATION 82**

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or

otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 4**

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 83**

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **diphenhydrAMINE (BENADRYL) injection 25**

**mg**

Dose: 25 mg

Route: intravenous

PRN

Start: S

#### **fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg

Route: oral

PRN

Start: S

#### **famotidine (PEPCID) 20 mg/2 mL injection 20**

**mg**

Dose: 20 mg

Route: intravenous

PRN

Start: S

#### **hydrocortisone sodium succinate**

**(Solu-CORTEF) injection 100 mg**

Dose: 100 mg      Route: intravenous      PRN

**dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg      Route: intravenous      PRN  
Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**

Dose: 0.3 mg      Route: subcutaneous      PRN  
Start: S

Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --      Occurrences: --  
Comments:      Discontinue IV.

Discharge Nursing Orders

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL      Route: intravenous      PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units      Route: intra-catheter      once PRN  
Start: S  
Instructions:  
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

**Days 8,15,22,29,36,43,50**

Perform every 7 days x7

Appointment Requests

**INFUSION APPOINTMENT REQUEST**

Interval: --      Occurrences: --

Labs

**CBC WITH PLATELET AND DIFFERENTIAL**

Interval: --      Occurrences: --

**COMPREHENSIVE METABOLIC PANEL**

Interval: --      Occurrences: --

**MAGNESIUM LEVEL**

Interval: --      Occurrences: --

**LDH**

Interval: --      Occurrences: --

**URIC ACID LEVEL**

Interval: --      Occurrences: --

Labs

**TYPE AND SCREEN**

Interval: --      Occurrences: --  
Comments:      Draw type and screen BEFORE each infusion.

Outpatient Electrolyte Replacement Protocol

**TREATMENT CONDITIONS 39**

Interval: --      Occurrences: --  
Comments:      Potassium (Normal range 3.5 to 5.0mEq/L)  
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP

- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

**TREATMENT CONDITIONS 40**

- Interval: -- Occurrences: --
- Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
  - o Protocol applies only to same day lab value.
  - o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
  - o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
  - o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
  - o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
  - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
  - o Sign electrolyte replacement order as Per protocol: cosign required

**Nursing Orders**

**ONC NURSING COMMUNICATION 73**

- Interval: -- Occurrences: --
- Comments: Draw type and screen BEFORE each infusion.

**Line Flush**

- sodium chloride 0.9 % flush 20 mL**
- Dose: 20 mL Route: intravenous PRN
- Start: S

**Nursing Orders**

- sodium chloride 0.9 % infusion 250 mL**
- Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
- Start: S
- Instructions: To keep vein open.

**Pre-Medications**

- methyIPREDNISolone sodium succinate (Solu-MEDROL) injection 100 mg**  
 Dose: 100 mg Route: intravenous once for 1 dose  
 Start: S  
 Instructions: Administer 30 minutes prior to chemotherapy.
- dexamethasone (DECADRON) 40 mg in sodium chloride 0.9 % IVPB**  
 Dose: 40 mg Route: intravenous once over 30 Minutes for 1 dose  
 Start: S  
 Instructions: Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	40 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

#### Pre-Medications

##### **acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg      Route: oral      once for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

#### Pre-Medications

##### **diphenhydramine (BENADRYL) tablet 25 mg**

Dose: 25 mg      Route: oral      once for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

#### Chemotherapy

##### **daratumumab 16 mg/kg in sodium chloride 0.9 % 500 mL IVPB**

Dose: 16 mg/kg      Route: intravenous      once @ 50 mL/hr for 1 dose  
Offset: 30 Minutes

Instructions:

Second infusion (500 mL volume): Infuse at 50 mL/hour for the first hour. Escalate the rate only if there were no grade 1 or greater infusion reactions during the first 3 hours of the first infusion. If no infusion reactions occur, may increase the rate by 50 mL/hour every hour (maximum rate: 200 mL/hour).

Subsequent infusions (500 mL volume): Escalate the rate only if there were no grade 1 or greater infusion reactions during a final infusion rate of =100 mL/hour in the first 2 infusions. Infuse at 100 mL/hour for the first hour. If no infusion reactions occur, may increase the rate by 50 mL/hour every hour (maximum rate: 200 mL/hour).

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DARATUMUMAB 20 MG/ML INTRAVENOUS SOLUTION	Medications	16 mg/kg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base		Yes	Yes

#### Delayed infusion reactions

##### **methylPREDNISolone sodium succinate (Solu-MEDROL) injection 20 mg**

Dose: 20 mg      Route: intravenous      once for 1 dose

Start: S

Hematology & Oncology Hypersensitivity Reaction Standing Order

**ONC NURSING COMMUNICATION 82**

Interval: -- Occurrences: --  
Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**ONC NURSING COMMUNICATION 4**

Interval: -- Occurrences: --  
Comments: Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**ONC NURSING COMMUNICATION 83**

Interval: -- Occurrences: --  
Comments: Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new



bag and new intravenous tubing.  
 8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.  
 9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.  
 10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**diphenhydramine (BENADRYL) injection 25 mg**

Dose: 25 mg                      Route: intravenous                      PRN  
 Start: S

**fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg                      Route: oral                      PRN  
 Start: S

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg                      Route: intravenous                      PRN  
 Start: S

**hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**

Dose: 100 mg                      Route: intravenous                      PRN

**dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg                      Route: intravenous                      PRN  
 Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**

Dose: 0.3 mg                      Route: subcutaneous                      PRN  
 Start: S

Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --                      Occurrences: --  
 Comments:                      Discontinue IV.

Discharge Nursing Orders

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL                      Route: intravenous                      PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units                      Route: intra-catheter                      once PRN  
 Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Weeks 9 - 24

Repeat 1 time

Cycle length: 112 days

Days 57,71,85,99,113,127,141,155

Perform every 14 days x8

Appointment Requests

**INFUSION APPOINTMENT REQUEST**

Interval: --                      Occurrences: --

Labs

**CBC WITH PLATELET AND DIFFERENTIAL**

Interval: --                      Occurrences: --

**COMPREHENSIVE METABOLIC PANEL**

Interval: -- Occurrences: --

**MAGNESIUM LEVEL**

Interval: -- Occurrences: --

**LDH**

Interval: -- Occurrences: --

**URIC ACID LEVEL**

Interval: -- Occurrences: --

Labs

**TYPE AND SCREEN**

Interval: -- Occurrences: --

Comments: Draw type and screen BEFORE each infusion.

Outpatient Electrolyte Replacement Protocol

**TREATMENT CONDITIONS 39**

Interval: -- Occurrences: --

Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

**TREATMENT CONDITIONS 40**

Interval: -- Occurrences: --

Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

**ONC NURSING COMMUNICATION 73**

Interval: -- Occurrences: --

Comments: Draw type and screen BEFORE each infusion.

Line Flush

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

Start: S

Nursing Orders

**sodium chloride 0.9 % infusion 250 mL**

Dose: 250 mL      Route: intravenous      once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

**methyIPREDNISolone sodium succinate (Solu-MEDROL) injection 100 mg**

Dose: 100 mg      Route: intravenous      once for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

**dexamethasone (DECADRON) 40 mg in sodium chloride 0.9 % IVPB**

Dose: 40 mg      Route: intravenous      once over 30 Minutes for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	40 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

Pre-Medications

**acetaminophen (TYLENOL) tablet 650 mg**

Dose: 650 mg      Route: oral      once for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

Pre-Medications

**diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg      Route: oral      once for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

Chemotherapy

**daratumumab 16 mg/kg in sodium chloride 0.9 % 500 mL IVPB**

Dose: 16 mg/kg      Route: intravenous      once @ 50 mL/hr for 1 dose  
Offset: 30 Minutes

Instructions:

Second infusion (500 mL volume): Infuse at 50 mL/hour for the first hour. Escalate the rate only if there were no grade 1 or greater infusion reactions during the first 3 hours of the first infusion. If no infusion reactions occur, may increase the rate by 50 mL/hour every hour (maximum rate: 200 mL/hour).

Subsequent infusions (500 mL volume):  
 Escalate the rate only if there were no grade 1 or greater infusion reactions during a final infusion rate of =100 mL/hour in the first 2 infusions. Infuse at 100 mL/hour for the first hour. If no infusion reactions occur, may increase the rate by 50 mL/hour every hour (maximum rate: 200 mL/hour).

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DARATUMUMAB MG/ML INTRAVENOUS SOLUTION	20 Medications	16 mg/kg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base		Yes	Yes

#### Delayed infusion reactions

##### **methyIPREDNISolone sodium succinate (Solu-MEDROL) injection 20 mg**

Dose: 20 mg      Route: intravenous      once for 1 dose  
 Start: S

#### Hematology & Oncology Hypersensitivity Reaction Standing Order

##### **ONC NURSING COMMUNICATION 82**

Interval: --      Occurrences: --  
 Comments:      Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

##### **ONC NURSING COMMUNICATION 4**

Interval: --      Occurrences: --  
 Comments:      Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy)

to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

### **ONC NURSING COMMUNICATION 83**

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.

2. Notify the CERT team and treating physician immediately.

3. Place the patient on continuous monitoring.

4. Obtain vital signs.

5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.

6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.

7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.

8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.

9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.

10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **diphenhydRAMINE (BENADRYL) injection 25 mg**

Dose: 25 mg

Route: intravenous

PRN

Start: S

#### **fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg

Route: oral

PRN

Start: S

#### **famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg

Route: intravenous

PRN

Start: S

#### **hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**

Dose: 100 mg

Route: intravenous

PRN

#### **dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg

Route: intravenous

PRN

Start: S

#### **epINEPhrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**

Dose: 0.3 mg

Route: subcutaneous

PRN

Start: S

Discharge Nursing Orders

### **ONC NURSING COMMUNICATION 76**

Interval: --

Occurrences: --

Comments:

Discontinue IV.

Discharge Nursing Orders

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL                      Route: intravenous                      PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units                      Route: intra-catheter                      once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.

**Weeks 25+ (until disease progression)**

Repeat 1 time

Cycle length: 630 days

**Days 169,197,225,253,281,309,337,365**

Perform every 28 days x8

Appointment Requests

**INFUSION APPOINTMENT REQUEST**

Interval: --                      Occurrences: --

Labs

**CBC WITH PLATELET AND DIFFERENTIAL**

Interval: --                      Occurrences: --

**COMPREHENSIVE METABOLIC PANEL**

Interval: --                      Occurrences: --

**MAGNESIUM LEVEL**

Interval: --                      Occurrences: --

**LDH**

Interval: --                      Occurrences: --

**URIC ACID LEVEL**

Interval: --                      Occurrences: --

Labs

**TYPE AND SCREEN**

Interval: --                      Occurrences: --

Comments:                      Draw type and screen BEFORE each infusion.

Outpatient Electrolyte Replacement Protocol

**TREATMENT CONDITIONS 39**

Interval: --                      Occurrences: --

Comments:                      Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

**TREATMENT CONDITIONS 40**

Interval: --                      Occurrences: --

Comments:                      Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

**Nursing Orders**

**ONC NURSING COMMUNICATION 73**

Interval: -- Occurrences: --  
 Comments: Draw type and screen BEFORE each infusion.

**Line Flush**

**sodium chloride 0.9 % flush 20 mL**  
 Dose: 20 mL Route: intravenous PRN  
 Start: S

**Nursing Orders**

**sodium chloride 0.9 % infusion 250 mL**  
 Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
 Start: S  
 Instructions: To keep vein open.

**Pre-Medications**

**dexamethasone (DECADRON) 40 mg in sodium chloride 0.9 % IVPB**  
 Dose: 40 mg Route: intravenous once over 30 Minutes for 1 dose  
 Start: S  
 Instructions: Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	40 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

**Pre-Medications**

**acetaminophen (TYLENOL) tablet 650 mg**  
 Dose: 650 mg Route: oral once for 1 dose  
 Start: S  
 Instructions: Administer 30 minutes prior to chemotherapy.

**Pre-Medications**

**diphenhydrAMINE (BENADRYL) tablet 25 mg**

Dose: 25 mg                      Route: oral                      once for 1 dose  
 Start: S  
 Instructions:  
 Administer 30 minutes prior to chemotherapy.

**Chemotherapy**

**daratumumab 16 mg/kg in sodium chloride 0.9 % 500 mL IVPB**

Dose: 16 mg/kg                      Route: intravenous                      once @ 50 mL/hr for 1 dose  
 Offset: 30 Minutes

Instructions:  
 Second infusion (500 mL volume): Infuse at 50 mL/hour for the first hour. Escalate the rate only if there were no grade 1 or greater infusion reactions during the first 3 hours of the first infusion. If no infusion reactions occur, may increase the rate by 50 mL/hour every hour (maximum rate: 200 mL/hour).

Subsequent infusions (500 mL volume): Escalate the rate only if there were no grade 1 or greater infusion reactions during a final infusion rate of =100 mL/hour in the first 2 infusions. Infuse at 100 mL/hour for the first hour. If no infusion reactions occur, may increase the rate by 50 mL/hour every hour (maximum rate: 200 mL/hour).

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DARATUMUMAB 20 MG/ML INTRAVENOUS SOLUTION	20 Medications	16 mg/kg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base		Yes	Yes

**Delayed infusion reactions**

**methyIPREDNISolone sodium succinate (Solu-MEDROL) injection 20 mg**

Dose: 20 mg                      Route: intravenous                      once for 1 dose  
 Start: S

**Hematology & Oncology Hypersensitivity Reaction Standing Order**

**ONC NURSING COMMUNICATION 82**

Interval: --                      Occurrences: --  
 Comments:                      Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)  
 1. Stop the infusion.  
 2. Place the patient on continuous monitoring.  
 3. Obtain vital signs.  
 4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.  
 5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.  
 6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.  
 7. Notify the treating physician.  
 8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).  
 9. Assess vital signs every 15 minutes until resolution of symptoms or



otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 4**

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O<sub>2</sub> saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 83**

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O<sub>2</sub> saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O<sub>2</sub> saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **diphenhydrAMINE (BENADRYL) injection 25**

**mg**

Dose: 25 mg

Route: intravenous

PRN

Start: S

#### **fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg

Route: oral

PRN

Start: S

#### **famotidine (PEPCID) 20 mg/2 mL injection 20**

**mg**

Dose: 20 mg

Route: intravenous

PRN

Start: S

#### **hydrocortisone sodium succinate**

**(Solu-CORTEF) injection 100 mg**

Dose: 100 mg      Route: intravenous      PRN

**dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg      Route: intravenous      PRN  
Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT  
injection syringe 0.3 mg**

Dose: 0.3 mg      Route: subcutaneous      PRN  
Start: S

Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --      Occurrences: --  
Comments:      Discontinue IV.

Discharge Nursing Orders

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL      Route: intravenous      PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units      Route: intra-catheter      once PRN  
Start: S  
Instructions:  
Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.