

OP DECITABINE (28 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: MDS, DECITABINE, DACOGEN, LEUKEMIA

Cycles 1 to 3	Repeat 3 times	Cycle length: 28 days
Days 1,2,3,4,5	Perform every 1 day x5	
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: -- Occurrences: --		
Labs		
☑ CBC WITH PLATELET AND DIFFERENTIAL		
Interval: -- Occurrences: --		
☑ COMPREHENSIVE METABOLIC PANEL		
Interval: -- Occurrences: --		
☐ BASIC METABOLIC PANEL		
Interval: -- Occurrences: --		
☑ MAGNESIUM LEVEL		
Interval: -- Occurrences: --		
☐ LDH		
Interval: -- Occurrences: --		
☐ URIC ACID LEVEL		
Interval: -- Occurrences: --		
☐ ECHOCARDIOGRAM COMPLETE W CONTRAST AND 3D IF NEEDED		
Interval: -- Occurrences: --		
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
TREATMENT CONDITIONS 40		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum Magnesium less than 1.0mEq/L. give 2 gram magnesium		

- sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
Start: S
Instructions:
To keep vein open.

Pre-Medications

☒ **ondansetron (ZOFTRAN) injection 8 mg**

Dose: 8 mg Route: intravenous once for 1 dose
Start: S End: S 11:15 AM

☐ **ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg Route: oral once for 1 dose
Start: S

☐ **ondansetron (ZOFTRAN) 16 mg in dextrose 5% 50 mL IVPB**

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Chemotherapy

decitabine (DACOGEN) 20 mg/m2 in sodium chloride 0.9 % 100 mL chemo IVPB

Dose: 20 mg/m2 Route: intravenous once over 1 Hours for 1 dose
Offset: 30 Minutes

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DECITABINE 50 MG INTRAVENOUS SOLUTION	Medications	20 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W)	QS Base		No	Yes

INTRAVENOUS
SOLUTION

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.