

OP DOCETAXEL (EVERY 21 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: DOCE, DOCETAXEL, TAXOTERE, TAX, TAXO, PROSTATE, GYNECOLOGIC, NSCLC, NON SQUAMOUS CELL LUNG CANCER

Take-Home Medications	Repeat 1 time	Cycle length: 1 day
Day 1		Perform every 1 day x1
Pre-Medications		
<input type="radio"/> dexamethasone (DECADRON) 4 MG tablet		
Dose: 8 mg	Route: oral	2 times daily
Dispense: 12 tablet	Refills: 5	
Start: S	End: S+3	
Instructions: Start day prior to chemotherapy administration.		
Cycles 1 to 4	Repeat 4 times	Cycle length: 21 days
Day 1		Perform every 1 day x1
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: --	Occurrences: --	
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: --	Occurrences: --	
<input type="checkbox"/> CARCINOEMBRYONIC ANTIGEN (CEA)		
Interval: --	Occurrences: --	
<input type="checkbox"/> PROSTATE SPECIFIC ANTIGEN		
Interval: --	Occurrences: --	
<input type="checkbox"/> URINALYSIS, AUTOMATED WITH MICROSCOPY		
Interval: --	Occurrences: --	
<input type="checkbox"/> CANCER ANTIGEN 27-29 (CA BR)		
Interval: --	Occurrences: --	
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: --	Occurrences: --	
Comments:		
	Potassium (Normal range 3.5 to 5.0mEq/L)	
	o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
	o Protocol applies only to same day lab value.	
	o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP	
	o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO	
	o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO	
	o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement	
	o If patient meets criteria, order SmartSet called "Outpatient	

Electrolyte Replacement"

- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: --

Occurrences: --

Comments:

Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

ONC NURSING COMMUNICATION 8

Interval: --

Occurrences: --

Comments:

Verify that patient took DEXAMETHASONE orally prior to chemotherapy. Otherwise, please contact physician to order Dexamethasone IV.

Nursing Orders

TREATMENT CONDITIONS 7

Interval: --

Occurrences: --

Comments:

HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL

Route: intravenous

PRN

Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) injection 8 mg

Dose: 8 mg

Route: intravenous

once for 1 dose

Start: S

End: S 11:15 AM

ondansetron (ZOFTRAN) tablet 16 mg

Dose: 16 mg

Route: oral

once for 1 dose

Start: S

ondansetron (ZOFTRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Chemotherapy

DOCEtaxel (TAXOTERE) 75 mg/m2 in sodium chloride (NON-PVC) 0.9 % 250 mL chemo IVPB

Dose: 75 mg/m2 Route: intravenous once over 60 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
Administer through non-DEHP tubing; Use within 4 hours of preparation; Protect from light.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DOCETAXEL 80 MG/4 ML (20 MG/ML) INTRAVENOUS SOLUTION	Medications	75 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	QS Base	250 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	QS Base	250 mL	No	Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Supportive Care

pegfilgrastim (NEULASTA) on-body injection kit 6 mg

Dose: 6 mg Route: subcutaneous once for 1 dose
Start: S End: S

Instructions:
Apply to intact, nonirritated skin on the back of the arm or abdomen (only use the back of the arm if caregiver is available to monitor



On-body injection status).