

## OP DVD (LOW DOSE)

*Types:* ONCOLOGY TREATMENT

*Synonyms:* MM, MULTIPLE, MYELO, DEXAM, DECA, BORTE, DOXOR, LIPOS, DOXI, LENAL, VELCAD, REVLAM, DVD, LOW

<b>Take-Home Medications</b>		Repeat 1 time	Cycle length: 1 day
<b>Day 1</b>		Perform every 1 day x1	
Take-Home Medications Prior to Treatment			
		<b>ONC PROVIDER COMMUNICATION 6</b>	
Interval: --		Occurrences: --	
Comments:		Prescribe antibiotic, anti-viral, and DVT prophylaxis as needed.	
Take-Home Medications Prior to Treatment			
		<b>lenalidomide (REVLIMID) 10 mg capsule</b>	
Dose: --		Route: oral	daily
Dispense: --		Refills: --	
Start: S			
<b>Cycles 1 to 8</b>		Repeat 8 times	Cycle length: 28 days
<b>Days 1,4</b>		Perform every 3 days x2	
Appointment Requests			
		<b>INFUSION APPOINTMENT REQUEST</b>	
Interval: --		Occurrences: --	
Labs			
<input checked="" type="checkbox"/> <b>COMPREHENSIVE METABOLIC PANEL</b>			
Interval: --		Occurrences: --	
<input checked="" type="checkbox"/> <b>CBC WITH PLATELET AND DIFFERENTIAL</b>			
Interval: --		Occurrences: --	
<input checked="" type="checkbox"/> <b>MAGNESIUM LEVEL</b>			
Interval: --		Occurrences: --	
<input checked="" type="checkbox"/> <b>LDH</b>			
Interval: --		Occurrences: --	
<input checked="" type="checkbox"/> <b>URIC ACID LEVEL</b>			
Interval: --		Occurrences: --	
<input checked="" type="checkbox"/> <b>PHOSPHORUS LEVEL</b>			
Interval: --		Occurrences: --	
Outpatient Electrolyte Replacement Protocol			
		<b>TREATMENT CONDITIONS 39</b>	
Interval: --		Occurrences: --	
Comments:		Potassium (Normal range 3.5 to 5.0mEq/L)	
		o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP	
		o Protocol applies only to same day lab value.	
		o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP	
		o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO	
		o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO	
		o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement	
		o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"	

- o Sign electrolyte replacement order as Per protocol: cosign required

#### **TREATMENT CONDITIONS 40**

Interval: -- Occurrences: --  
 Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

#### **Nursing Orders**

#### **TREATMENT CONDITIONS 7**

Interval: -- Occurrences: --  
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

#### **Vitals**

#### **ONC NURSING COMMUNICATION 50**

Interval: -- Occurrences: --  
 Comments: 1) Check vital signs (BP, temperature, pulse, and respirations) prior to and 30 minutes after Bortezomib administration.

2) If systolic BP, 30 minutes after Bortezomib infusion, drops more than 30 mmHg, or if systolic BP is less than 90, please contact MD.

#### **Nursing Orders**

#### **ONC NURSING COMMUNICATION 51**

Interval: -- Occurrences: --  
 Comments: HOLD Bortezomib and notify provider if Hgb is LESS than or equal to \*\*\* g/dL.

#### **Line Flush**

#### **dextrose 5% flush syringe 20 mL**

Dose: 20 mL Route: intravenous PRN  
 Start: S  
 Instructions:  
 Administer ONLY for Liposomal Doxorubicin.

#### **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN  
 Start: S  
 Instructions:  
 Do NOT administer with Liposomal Doxorubicin.

#### **Nursing Orders**

#### **dextrose 5% infusion 250 mL**

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
 Start: S

Instructions:

To keep vein open for Liposomal Doxorubicin.

**sodium chloride 0.9 % infusion 250 mL**

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open. Do NOT administer with Liposomal Doxorubicin.

Nursing Orders

**ONC NURSING COMMUNICATION 60**

Interval: --

Occurrences: --

Comments:

Verify patient taking adequate prophylaxis medications and Lenalidomide.

Pre-Medications

**dexamethasone (DECADRON) 40 mg in sodium chloride 0.9 % IVPB**

Dose: 40 mg

Route: intravenous

once over 30 Minutes for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

**Ingredients:**

**Name**

**Type**

**Dose**

**Selected**

**Adds Vol.**

DEXAMETHASONE  
4 MG/ML  
INJECTION  
SOLUTION  
SODIUM  
CHLORIDE 0.9 %  
INTRAVENOUS  
SOLUTION  
DEXTROSE 5 % IN  
WATER (D5W)  
INTRAVENOUS  
SOLUTION

Medications

40 mg

Main  
Ingredient

Yes

Base

50 mL

Yes

Yes

Base

50 mL

No

Yes

Chemotherapy

**bortezomib (VelCADE) 1 mg/m2 in sodium chloride 0.9 % chemo injection**

Dose: 1 mg/m2

Route: subcutaneous

once for 1 dose

Offset: 30 Minutes

Instructions:

DRUG IS AN IRRITANT. Administer drug slowly to prevent burning upon administration. 72 hours between doses is recommended.

**Ingredients:**

**Name**

**Type**

**Dose**

**Selected**

**Adds Vol.**

BORTEZOMIB 3.5  
MG SOLUTION  
FOR INJECTION  
SODIUM  
CHLORIDE 0.9 %  
INJECTION  
SOLUTION

Medications

1 mg/m2

Main  
Ingredient

No

Base

Always

Yes

Chemotherapy

**DOXOrubicin liposomal (DOXIL) 4 mg/m2 in dextrose 5% 250 mL chemo IVPB**

Dose: 4 mg/m2

Route: intravenous

once over 1 Hours for 1 dose

Offset: 30 Minutes

Instructions:

DRUG IS AN IRRITANT. Initial infusion infused at 1 mg/min, but no faster than 1 hour to prevent infusion related reactions. Monitor

vital signs 15 minutes, 30 minutes, and one hour into infusion, then hourly for remainder of initial infusion. Stay with patient for the first 15 minutes of the initial infusion. If patient tolerated initial infusion, subsequent infusions to be given over 1 hour.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DOXORUBICIN, PEGYLATED LIPOSOMAL 2 MG/ML INTRAVENOUS SUSPENSION	Medications	4 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes

#### Hematology & Oncology Hypersensitivity Reaction Standing Order

##### ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --  
 Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)  
 1. Stop the infusion.  
 2. Place the patient on continuous monitoring.  
 3. Obtain vital signs.  
 4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.  
 5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.  
 6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.  
 7. Notify the treating physician.  
 8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).  
 9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

##### ONC NURSING COMMUNICATION 4

Interval: -- Occurrences: --  
 Comments: Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)  
 1. Stop the infusion.  
 2. Notify the CERT team and treating physician immediately.  
 3. Place the patient on continuous monitoring.  
 4. Obtain vital signs.  
 5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.  
 6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.  
 7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.  
 8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).  
 9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**ONC NURSING COMMUNICATION 83**

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**diphenhydramine (BENADRYL) injection 25 mg**

Dose: 25 mg

Route: intravenous

PRN

Start: S

**fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg

Route: oral

PRN

Start: S

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg

Route: intravenous

PRN

Start: S

**hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**

Dose: 100 mg

Route: intravenous

PRN

**dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg

Route: intravenous

PRN

Start: S

**epinephrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**

Dose: 0.3 mg

Route: subcutaneous

PRN

Start: S

## Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --

Occurrences: --

Comments:

Discontinue IV.

## Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL

Route: intravenous

PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units

Route: intra-catheter

once PRN

Start: S

Instructions:  
Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.

**Days 8,11**

Perform every 3 days x2

Appointment Requests

**INFUSION APPOINTMENT REQUEST**

Interval: -- Occurrences: --

Labs

☒ **COMPREHENSIVE METABOLIC PANEL**

Interval: -- Occurrences: --

☒ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: -- Occurrences: --

☒ **MAGNESIUM LEVEL**

Interval: -- Occurrences: --

☒ **LDH**

Interval: -- Occurrences: --

☒ **URIC ACID LEVEL**

Interval: -- Occurrences: --

☒ **PHOSPHORUS LEVEL**

Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

**TREATMENT CONDITIONS 39**

Interval: -- Occurrences: --

Comments:

Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

**TREATMENT CONDITIONS 40**

Interval: -- Occurrences: --

Comments:

Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria. order SmartSet called "Outpatient

Electrolyte Replacement"

o Sign electrolyte replacement order as Per protocol: cosign required

#### Nursing Orders

##### **TREATMENT CONDITIONS 7**

Interval: --

Occurrences: --

Comments:

HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

#### Vitals

##### **ONC NURSING COMMUNICATION 50**

Interval: --

Occurrences: --

Comments:

1) Check vital signs (BP, temperature, pulse, and respirations) prior to and 30 minutes after Bortezomib administration.

2) If systolic BP, 30 minutes after Bortezomib infusion, drops more than 30 mmHg, or if systolic BP is less than 90, please contact MD.

#### Nursing Orders

##### **ONC NURSING COMMUNICATION 51**

Interval: --

Occurrences: --

Comments:

HOLD Bortezomib and notify provider if Hgb is LESS than or equal to \*\*\* g/dL.

#### Line Flush

##### **dextrose 5% flush syringe 20 mL**

Dose: 20 mL

Route: intravenous

PRN

Start: S

Instructions:

Administer ONLY for Liposomal Doxorubicin.

##### **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL

Route: intravenous

PRN

Start: S

Instructions:

Do NOT administer with Liposomal Doxorubicin.

#### Nursing Orders

##### **dextrose 5% infusion 250 mL**

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open for Liposomal Doxorubicin.

##### **sodium chloride 0.9 % infusion 250 mL**

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open. Do NOT administer with Liposomal Doxorubicin.

#### Nursing Orders

##### **ONC NURSING COMMUNICATION 60**

Interval: --

Occurrences: --

Comments:

Verify patient taking adequate prophylaxis medications and Lenalidomide.

#### Pre-Medications

**dexamethasone (DECADRON) 40 mg in sodium chloride 0.9 % IVPB**

Dose: 40 mg      Route: intravenous      once over 30 Minutes for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	40 mg	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

#### Chemotherapy

##### **bortezomib (VelCADE) 1 mg/m2 in sodium chloride 0.9 % chemo injection**

Dose: 1 mg/m2      Route: subcutaneous      once for 1 dose  
Offset: 30 Minutes

Instructions:

DRUG IS AN IRRITANT. Administer drug slowly to prevent burning upon administration. 72 hours between doses is recommended.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	BORTEZOMIB 3.5 MG SOLUTION FOR INJECTION	Medications	1 mg/m2	Main Ingredient	No
	SODIUM CHLORIDE 0.9 % INJECTION SOLUTION	Base		Always	Yes

#### Chemotherapy

##### **DOXOrubicin liposomal (DOXIL) 4 mg/m2 in dextrose 5% 250 mL chemo IVPB**

Dose: 4 mg/m2      Route: intravenous      once over 1 Hours for 1 dose  
Offset: 30 Minutes

Instructions:

DRUG IS AN IRRITANT. Initial infusion infused at 1 mg/min, but no faster than 1 hour to prevent infusion related reactions. Monitor vital signs 15 minutes, 30 minutes, and one hour into infusion, then hourly for remainder of initial infusion. Stay with patient for the first 15 minutes of the initial infusion. If patient tolerated initial infusion, subsequent infusions to be given over 1 hour.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DOXORUBICIN, PEGYLATED LIPOSOMAL 2 MG/ML INTRAVENOUS SUSPENSION	Medications	4 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes



**ONC NURSING COMMUNICATION 82**

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**ONC NURSING COMMUNICATION 4**

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**ONC NURSING COMMUNICATION 83**

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.

8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.  
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.  
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**diphenhydramine (BENADRYL) injection 25 mg**

Dose: 25 mg                      Route: intravenous                      PRN  
Start: S

**fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg                      Route: oral                      PRN  
Start: S

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg                      Route: intravenous                      PRN  
Start: S

**hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**

Dose: 100 mg                      Route: intravenous                      PRN

**dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg                      Route: intravenous                      PRN  
Start: S

**epinephrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**

Dose: 0.3 mg                      Route: subcutaneous                      PRN  
Start: S

Discharge Nursing Orders

**ONC NURSING COMMUNICATION 76**

Interval: --                      Occurrences: --  
Comments:                      Discontinue IV.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL                      Route: intravenous                      PRN

☒ **HEparin, porcine (PF) injection 500 Units**

Dose: 500 Units                      Route: intra-catheter                      once PRN  
Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.