

OP FOLFIRI / BEVACIZUMAB (EVERY 14 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: FOLFIRI, IRINOTECAN, LEUCOVORIN, 5FU, 5-FLUOROURACIL, FLUOROURACIL, LEUCO, IRIN, CAMPTOSAR, CAMP, BEV, BEVACIZUMAB, AVAS, AVASTIN, COLORECTAL

Cycles 1 to 6	Repeat 6 times	Cycle length: 14 days
Day 1	Perform every 1 day x1	
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: -- Occurrences: --		
Labs		
URINALYSIS, AUTOMATED WITH MICROSCOPY		
Interval: -- Occurrences: --		
CBC WITH PLATELET AND DIFFERENTIAL		
Interval: -- Occurrences: --		
COMPREHENSIVE METABOLIC PANEL		
Interval: -- Occurrences: --		
MAGNESIUM LEVEL		
Interval: -- Occurrences: --		
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
TREATMENT CONDITIONS 40		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP		
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV		
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV		
o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement		
o If patient meets criteria. order SmartSet called "Outpatient		

Electrolyte Replacement"

o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS

Interval: --

Occurrences: --

Comments:

Do NOT administer within 28 days of surgery/procedure and until the surgical wound is fully healed or within 14 days of port placement.

Nursing Orders

TREATMENT CONDITIONS 5

Interval: --

Occurrences: --

Comments:

HOLD and notify provider if PROTEIN 2+ is detected in UA.

Nursing Orders

TREATMENT CONDITIONS 4

Interval: --

Occurrences: --

Comments:

HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000; Total Bilirubin GREATER than 1.5; ALT/AST GREATER than 3 times upper normal limit; or Serum Creatinine GREATER than 1.2

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL

Route: intravenous

PRN

Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg, dexamethasone☒ **(DECADRON) 12 mg in sodium chloride 0.9%****50 mL IVPB**

Dose: --

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:30 AM

Ingredients:**Name****Type****Dose****Selected****Adds Vol.**

ONDANSETRON

Medications

16 mg

Yes

No

HCL (PF) 4 MG/2

ML INJECTION

SOLUTION

DEXAMETHASONE Medications

12 mg

Yes

No

4 MG/ML

INJECTION

SOLUTION

SODIUM

Base

50 mL

Always

Yes

CHLORIDE 0.9 %

INTRAVENOUS

SOLUTION

DEXTROSE 5 % IN Base

No

Yes

WATER (D5W)

INTRAVENOUS

SOLUTION

☐ **ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg

Route: oral

once for 1 dose

Start: S

End: S 11:30 AM

☐ **dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg
Start: S

Route: oral

once for 1 dose

☐ **palonosetron (ALOXI) injection 0.25 mg**

Dose: 250 mcg
Start: S
Instructions:

Route: intravenous
End: S 3:00 PM

once for 1 dose

For OUTPATIENT use only.

☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg
Start: S

Route: intravenous
End: S

once over 30 Minutes for 1 dose

Ingredients:

Name

Type

Dose

Selected
Main
Ingredient

Adds Vol.
Yes

APREPITANT 7.2
MG/ML

Medications

130 mg

INTRAVENOUS
EMULSION

DEXTROSE 5 % IN
WATER (D5W) IV

Base

130 mL

Yes

Yes

SOLP (EXCEL;
NON-PVC)

SODIUM
CHLORIDE 0.9 % IV
SOLP

Base

130 mL

No

Yes

(EXCEL;NON-PVC)

Pre-Medications

atropine injection 0.25 mg

Dose: 0.25 mg
Start: S

Route: intravenous

once PRN

Supportive Care

☐ **LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg
Start: S

Route: intravenous

once PRN

☐ **LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg
Start: S

Route: oral

once PRN

Chemotherapy

bevacizumab (AVASTIN) 5 mg/kg in sodium chloride 0.9 % 100 mL IVPB

Dose: 5 mg/kg

Route: intravenous

once over 30 Minutes for 1 dose
Offset: 30 Minutes

Ingredients:

Name

Type

Dose

Selected
Main
Ingredient

Adds Vol.
Yes

BEVACIZUMAB 25
MG/ML

Medications

5 mg/kg

INTRAVENOUS
SOLUTION

SODIUM
CHLORIDE 0.9 %

QS Base

100 mL

Yes

Yes

INTRAVENOUS
SOLUTION

leucovorin 400 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 400 mg/m2

Route: intravenous

once over 90 Minutes for 1 dose
Offset: 1 Hours

Ingredients:

Name

Type

Dose

Selected
Main

Adds Vol.
Yes

LEUCOVORIN

Medications

400

CALCIUM 350 MG SOLUTION FOR INJECTION		mg/m2	Ingredient	
DEXTROSE 5 % IN WATER (D5W)	QS Base	250 mL	No	Yes
INTRAVENOUS SOLUTION				
SODIUM CHLORIDE 0.9 %	QS Base	250 mL	Yes	Yes
INTRAVENOUS SOLUTION				

irinotecan (CAMPTOSAR) 180 mg/m2 in dextrose 5% 500 mL chemo IVPB

Dose: 180 mg/m2 Route: intravenous once over 90 Minutes for 1 dose
Offset: 1 Hours

Instructions:
Protect from light

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
IRINOTECAN 100 MG/5 ML	Medications	180 mg/m2	Main Ingredient	Yes
INTRAVENOUS SOLUTION				
DEXTROSE 5 % IN WATER (D5W)	QS Base	500 mL	Yes	Yes
INTRAVENOUS SOLUTION				
SODIUM CHLORIDE 0.9 %	QS Base	500 mL	No	Yes
INTRAVENOUS SOLUTION				

fluorouracil (ADRUCIL) 400 mg/m2 in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 400 mg/m2 Route: intravenous once over 15 Minutes for 1 dose
Offset: 2.5 Hours

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
FLUOROURACIL 500 MG/10 ML	Medications	400 mg/m2	Main Ingredient	Yes
INTRAVENOUS SOLUTION				
DEXTROSE 5 % IN WATER (D5W)	Base	50 mL	No	Yes
INTRAVENOUS SOLUTION				
SODIUM CHLORIDE 0.9 %	Base	50 mL	Yes	Yes
INTRAVENOUS SOLUTION				

fluorouracil (ADRUCIL) 2,400 mg/m2 in sodium chloride 0.9 % 100 mL chemo infusion - AMBULATORY PUMP

Dose: 2,400 mg/m2 Route: intravenous once over 48 Hours for 1 dose
Offset: 2.75 Hours

Instructions:
Nurses to chart as GIVEN on the MAR.
Administer via CADD pump.

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
FLUOROURACIL 5 GRAM/100 ML	Medications	2,400 mg/m2	Main Ingredient	Yes
INTRAVENOUS SOLUTION				
DEXTROSE 5 % IN	QS Base	100 mL	No	Yes

WATER (D5W) INTRAVENOUS SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes
---	---------	--------	-----	-----

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.

4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epinephrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Day 3

Perform every 1 day x1

Appointment Requests

ONC PUMP DISCONNECT APPOINTMENT REQUEST

Interval: -- Occurrences: --

Discharge Nursing Orders

DISCONNECT CONTINUOUS INFUSION PUMP

Interval: -- Occurrences: --
Comments: Disconnect patient from continuous infusion pump.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

☒ **HEparin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Post-Medications

○ **pegfilgrastim (NEULASTA) on-body injection kit 6 mg**

Dose: 6 mg

Route: subcutaneous once for 1 dose

Start: S

End: S

Instructions:

Apply to intact, nonirritated skin on the back of the arm or abdomen (only use the back of the arm if caregiver is available to monitor On-body injection status).