

## OP FOLFIRI / PANITUMUMAB

*Types:* ONCOLOGY TREATMENT

*Synonyms:* FOLFIRI, IRINOTECAN, LEUCOVORIN, 5FU, 5-FLUOROURACIL, FLUOROURACIL, LEUCO, IRIN, CAMPTOSAR, CAMP, PANT, PANITUMUMAB, VEX, VECTIBIX, COLORECTAL

|                      |  |                                                                               |                       |
|----------------------|--|-------------------------------------------------------------------------------|-----------------------|
| <b>Cycles 1 to 6</b> |  | Repeat 6 times                                                                | Cycle length: 14 days |
| <b>Day 1</b>         |  | Perform every 1 day x1                                                        |                       |
| Appointment Requests |  |                                                                               |                       |
|                      |  | <b>INFUSION APPOINTMENT REQUEST</b>                                           |                       |
|                      |  | Interval: --                                                                  | Occurrences: --       |
| Labs                 |  |                                                                               |                       |
|                      |  | <input checked="" type="checkbox"/> <b>COMPREHENSIVE METABOLIC PANEL</b>      |                       |
|                      |  | Interval: --                                                                  | Occurrences: --       |
|                      |  | <input checked="" type="checkbox"/> <b>CBC WITH PLATELET AND DIFFERENTIAL</b> |                       |
|                      |  | Interval: --                                                                  | Occurrences: --       |
|                      |  | <input checked="" type="checkbox"/> <b>MAGNESIUM LEVEL</b>                    |                       |
|                      |  | Interval: --                                                                  | Occurrences: --       |

## Outpatient Electrolyte Replacement Protocol

### TREATMENT CONDITIONS 39

Interval: -- Occurrences: --  
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)  
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP  
o Protocol applies only to same day lab value.  
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP  
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO  
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO  
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement  
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"  
o Sign electrolyte replacement order as Per protocol: cosign required

### TREATMENT CONDITIONS 40

Interval: -- Occurrences: --  
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)  
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP  
o Protocol applies only to same day lab value.  
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP  
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV  
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV  
o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement  
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"  
o Sign electrolyte replacement order as Per protocol: cosign required

## Nursing Orders

### TREATMENT CONDITIONS 4

Interval: -- Occurrences: --  
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000; Total Bilirubin GREATER than 1.5; ALT/AST GREATER than 3 times upper normal limit; or Serum Creatinine GREATER than 1.2

## Provider Communication

### ONC PROVIDER COMMUNICATION 2

Interval: -- Occurrences: --  
Comments: Tumor KRAS gene status should be determined prior to initiation of therapy. KRAS type: Please Push F2:115540219.

## Line Flush

### sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN  
Start: S

## Nursing Orders

### sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
Start: S  
Instructions:  
To keep vein open.

## Pre-Medications

- ☒ **ondansetron (ZOFTRAN) 16 mg, dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: --

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:30 AM

**Ingredients:****Name****Type****Dose****Selected****Adds Vol.**

ONDANSETRON

Medications

16 mg

Yes

No

HCL (PF) 4 MG/2

ML INJECTION

SOLUTION

DEXAMETHASONE Medications

12 mg

Yes

No

4 MG/ML

INJECTION

SOLUTION

SODIUM

Base

50 mL

Always

Yes

CHLORIDE 0.9 %

INTRAVENOUS

SOLUTION

DEXTROSE 5 % IN

Base

No

Yes

WATER (D5W)

INTRAVENOUS

SOLUTION

- ☐ **ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg

Route: oral

once for 1 dose

Start: S

End: S 11:30 AM

- ☐ **dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg

Route: oral

once for 1 dose

Start: S

- ☐ **palonosetron (ALOXI) injection 0.25 mg**

Dose: 250 mcg

Route: intravenous

once for 1 dose

Start: S

End: S 3:00 PM

Instructions:

For OUTPATIENT use only.

- ☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg

Route: intravenous

once over 30 Minutes for 1 dose

Start: S

End: S

**Ingredients:****Name****Type****Dose****Selected****Adds Vol.**

APREPITANT 7.2

Medications

130 mg

Main

Yes

MG/ML

Ingredient

INTRAVENOUS

EMULSION

DEXTROSE 5 % IN

Base

130 mL

Yes

Yes

WATER (D5W) IV

SOLP (EXCEL;

NON-PVC)

SODIUM

Base

130 mL

No

Yes

CHLORIDE 0.9 % IV

SOLP

(EXCEL;NON-PVC)

## Pre-Medications

**atropine injection 0.25 mg**

Dose: 0.25 mg

Route: intravenous

once PRN

Start: S

## Supportive Care

☐ **LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg  
Start: S

Route: intravenous

once PRN

☐ **LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg  
Start: S

Route: oral

once PRN

**Chemotherapy**

**panitumumab (VECTIBIX) 6 mg/kg in sodium chloride 0.9 % 100 mL chemo IVPB**

Dose: 6 mg/kg

Route: intravenous

once over 60 Minutes for 1 dose  
Offset: 30 Minutes

Instructions:

USE 0.2 OR 0.22 MICRON INLINE FILTER.

**Ingredients:**

**Name**

PANITUMUMAB  
100 MG/5 ML (20  
MG/ML)  
INTRAVENOUS  
SOLUTION  
SODIUM  
CHLORIDE 0.9 %  
INTRAVENOUS  
SOLUTION

**Type**

Medications

**Dose**

6 mg/kg

**Selected**

Main  
Ingredient

**Adds Vol.**

Yes

**leucovorin 400 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB**

Dose: 400 mg/m2

Route: intravenous

once over 90 Minutes for 1 dose  
Offset: 1.5 Hours

**Ingredients:**

**Name**

LEUCOVORIN  
CALCIUM 350 MG  
SOLUTION FOR  
INJECTION  
DEXTROSE 5 % IN  
WATER (D5W)  
INTRAVENOUS  
SOLUTION  
SODIUM  
CHLORIDE 0.9 %  
INTRAVENOUS  
SOLUTION

**Type**

Medications

**Dose**

400  
mg/m2

**Selected**

Main  
Ingredient

**Adds Vol.**

Yes

**irinotecan (CAMPTOSAR) 180 mg/m2 in dextrose 5% 500 mL chemo IVPB**

Dose: 180 mg/m2

Route: intravenous

once over 90 Minutes for 1 dose  
Offset: 1.5 Hours

Instructions:

Protect from light

**Ingredients:**

**Name**

IRINOTECAN 100  
MG/5 ML  
INTRAVENOUS  
SOLUTION  
DEXTROSE 5 % IN  
WATER (D5W)  
INTRAVENOUS  
SOLUTION  
SODIUM  
CHLORIDE 0.9 %  
INTRAVENOUS

**Type**

Medications

**Dose**

180  
mg/m2

**Selected**

Main  
Ingredient

**Adds Vol.**

Yes

## SOLUTION

### fluorouracil (ADRUCIL) 400 mg/m<sup>2</sup> in sodium chloride 0.9 % 50 mL chemo IVPB

Dose: 400 mg/m<sup>2</sup>

Route: intravenous

once over 15 Minutes for 1 dose

Offset: 3 Hours

#### Ingredients:

#### Name

#### Type

#### Dose

#### Selected

#### Adds Vol.

FLUOROURACIL

Medications

400

Main

Yes

500 MG/10 ML

mg/m<sup>2</sup>

Ingredient

INTRAVENOUS

SOLUTION

DEXTROSE 5 % IN

Base

50 mL

No

Yes

WATER (D5W)

INTRAVENOUS

SOLUTION

SODIUM

Base

50 mL

Yes

Yes

CHLORIDE 0.9 %

INTRAVENOUS

SOLUTION

### fluorouracil (ADRUCIL) 2,400 mg/m<sup>2</sup> in sodium chloride 0.9 % 100 mL chemo infusion - AMBULATORY PUMP

Dose: 2,400 mg/m<sup>2</sup>

Route: intravenous

once over 48 Hours for 1 dose

Offset: 3.25 Hours

#### Instructions:

Nurses to chart as GIVEN on the MAR.

Administer via CADD pump.

#### Ingredients:

#### Name

#### Type

#### Dose

#### Selected

#### Adds Vol.

FLUOROURACIL 5

Medications

2,400

Main

Yes

GRAM/100 ML

mg/m<sup>2</sup>

Ingredient

INTRAVENOUS

SOLUTION

DEXTROSE 5 % IN

QS Base

100 mL

No

Yes

WATER (D5W)

INTRAVENOUS

SOLUTION

SODIUM

QS Base

100 mL

Yes

Yes

CHLORIDE 0.9 %

INTRAVENOUS

SOLUTION

## Hematology & Oncology Hypersensitivity Reaction Standing Order

### ONC NURSING COMMUNICATION 82

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.

2. Place the patient on continuous monitoring.

3. Obtain vital signs.

4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.

5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.

6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.

7. Notify the treating physician.

8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).

9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**ONC NURSING COMMUNICATION 4**

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**ONC NURSING COMMUNICATION 83**

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**diphenhydramine (BENADRYL) injection 25****mg**

Dose: 25 mg

Route: intravenous

PRN

Start: S

**fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg

Route: oral

PRN

Start: S

**famotidine (PEPCID) 20 mg/2 mL injection 20****mg**

Dose: 20 mg

Route: intravenous

PRN

Start: S

**hydrocortisone sodium succinate  
(Solu-CORTEF) injection 100 mg**

Dose: 100 mg      Route: intravenous      PRN

**dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg      Route: intravenous      PRN  
Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT  
injection syringe 0.3 mg**

Dose: 0.3 mg      Route: subcutaneous      PRN  
Start: S

**Nursing Orders**

**ONC NURSING COMMUNICATION 14**

Interval: --      Occurrences: --  
Comments:      Contact Provider if drug-induced acneiform rash develops and covers more than 25 per cent of the body.

**Discharge Nursing Orders**

**ONC NURSING COMMUNICATION 76**

Interval: --      Occurrences: --  
Comments:      Discontinue IV.

**Discharge Nursing Orders**

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL      Route: intravenous      PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units      Route: intra-catheter      once PRN  
Start: S  
Instructions:  
Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.

**Day 3**

Perform every 1 day x1

**Appointment Requests**

**ONC PUMP DISCONNECT APPOINTMENT  
REQUEST**

Interval: --      Occurrences: --

**Discharge Nursing Orders**

**DISCONNECT CONTINUOUS INFUSION PUMP**

Interval: --      Occurrences: --  
Comments:      Disconnect patient from continuous infusion pump.

**Discharge Nursing Orders**

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL      Route: intravenous      PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units      Route: intra-catheter      once PRN  
Start: S  
Instructions:  
Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.

**Post-Medications**

○ **pegfilgrastim (NEULASTA) on-body injection  
kit 6 mg**

Dose: 6 mg

Route: subcutaneous    once for 1 dose

Start: S

End: S

Instructions:

Apply to intact, nonirritated skin on the back of  
the arm or abdomen (only use the back of the  
arm if caregiver is available to monitor  
On-body injection status).