

OP FOLFOXIRI / CETUXIMAB (EVERY 14 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: FOLFOXIRI, FLUOROURACIL , LEUCOVORIN, OXALIPLATIN , IIRINOTECAN , AGE, CAMPTOSAR, ADRUCIL, ELOXATIN, FLOW, IRENE, CETUX, ERBIT

Cycles 1 to 6		Repeat 6 times	Cycle length: 14 days
Day 1		Perform every 1 day x1	
Appointment Requests			
		INFUSION APPOINTMENT REQUEST	
		Interval: --	Occurrences: --
Labs			
		☒ COMPREHENSIVE METABOLIC PANEL	
		Interval: --	Occurrences: --
		☒ CBC WITH PLATELET AND DIFFERENTIAL	
		Interval: --	Occurrences: --
		☒ MAGNESIUM LEVEL	
		Interval: --	Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39

Interval: -- Occurrences: --
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
o Protocol applies only to same day lab value.
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
o Protocol applies only to same day lab value.
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 4

Interval: -- Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000; Total Bilirubin GREATER than 1.5; ALT/AST GREATER than 3 times upper normal limit; or Serum Creatinine GREATER than 1.2

Provider Communication

ONC PROVIDER COMMUNICATION 2

Interval: -- Occurrences: --
Comments: Tumor KRAS gene status should be determined prior to initiation of therapy. KRAS type: Please Push F2:115540219.

Line Flush

dextrose 5% flush syringe 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S
Instructions:
Administer ONLY for Oxaliplatin.

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
Start: S
Instructions:

Do NOT administer with Oxaliplatin.

Nursing Orders

dextrose 5% infusion 250 mL

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open for Oxaliplatin.

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL

Route: intravenous

once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open. Do NOT administer with Oxaliplatin.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg, dexamethasone

- ☒ **(DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: --

Route: intravenous

once over 15 Minutes for 1 dose

Start: S

End: S 11:30 AM

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

ONDANSETRON
HCL (PF) 4 MG/2
ML INJECTION
SOLUTION

Medications

16 mg

Yes

No

DEXAMETHASONE
4 MG/ML
INJECTION
SOLUTION

Medications

12 mg

Yes

No

SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Base

50 mL

Always

Yes

DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION

Base

No

Yes

- ☐ **ondansetron (ZOFTRAN) tablet 16 mg**

Dose: 16 mg

Route: oral

once for 1 dose

Start: S

End: S 11:30 AM

- ☐ **dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg

Route: oral

once for 1 dose

Start: S

- ☐ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg

Route: intravenous

once over 30 Minutes for 1 dose

Start: S

End: S

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

APREPITANT 7.2
MG/ML
INTRAVENOUS
EMULSION

Medications

130 mg

Main
Ingredient

Yes

DEXTROSE 5 % IN
WATER (D5W) IV
SOLP (EXCEL;
NON-PVC)

Base

130 mL

Yes

Yes

SODIUM
CHLORIDE 0.9 % IV
SOLP

Base

130 mL

No

Yes

(EXCEL;NON-PVC)

Pre-Medications

atropine injection 0.25 mg

Dose: 0.25 mg

Route: intravenous

once PRN

Start: S

Pre-Medications

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg

Route: intravenous

once for 1 dose

Start: S

Instructions:

Give 30 minutes prior to cetuximab.

Supportive Care

☐ **LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg

Route: intravenous

once PRN

Start: S

☐ **LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg

Route: oral

once PRN

Start: S

Chemotherapy

cetuximab (ERBITUX) 400 mg/m2 in 0 mL

Dose: 400 mg/m2

Route: intravenous

once over 120 Minutes for 1 dose

Offset: 30 Minutes

Instructions:

Administer with low protein binding 0.22 micron filter. Do not shake. Do not mix with other medications. Flush IV line with NS at the end of infusion.

1st infusion: Infuse first 10 mL over 10 minutes and observe patient for 30 minutes for allergic reactions if infusion tolerated, infuse loading dose over 120 minutes.

Rate of infusion not to exceed 10 mg/minute (5 mL/minute)

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

CETUXIMAB 100
MG/50 ML
INTRAVENOUS
SOLUTION

Medications

400
mg/m2

Main
Ingredient

Yes

**leucovorin 200 mg/m2 in dextrose 5% 250 mL
chemo IVPB**

Dose: 200 mg/m2

Route: intravenous

once over 90 Minutes for 1 dose

Offset: 2.5 Hours

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

LEUCOVORIN
CALCIUM 350 MG
SOLUTION FOR
INJECTION
DEXTROSE 5 % IN
WATER (D5W)
INTRAVENOUS
SOLUTION
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS

Medications

200
mg/m2

Main
Ingredient

Yes

QS Base

250 mL

Yes

Yes

QS Base

250 mL

No

Yes

SOLUTION

irinotecan (CAMPTOSAR) 165 mg/m2 in dextrose 5% 500 mL chemo IVPB

Dose: 165 mg/m2 Route: intravenous once over 90 Minutes for 1 dose
Offset: 2.5 Hours

Instructions:
Protect from light

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	IRINOTECAN 100 MG/5 ML INTRAVENOUS SOLUTION	Medications	165 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	No	Yes

OXALiplatIn (ELOXATIN) 85 mg/m2 in dextrose 5% 500 mL chemo IVPB

Dose: 85 mg/m2 Route: intravenous once over 2 Hours for 1 dose
Offset: 4 Hours

Instructions:
Irritant - avoid extravasation. Flush line with D5W before and after oxaliplatin infusion.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	OXALIPLATIN 100 MG/20 ML INTRAVENOUS SOLUTION	Medications	85 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes

fluorouracil (ADRUCIL) 3,200 mg/m2 in sodium chloride 0.9 % 100 mL chemo infusion - AMBULATORY PUMP

Dose: 3,200 mg/m2 Route: intravenous once over 48 Hours for 1 dose
Offset: 6 Hours

Instructions:
Nurses to chart as GIVEN on the MAR.
Administer via CADD pump.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	FLUOROURACIL 5 GRAM/100 ML INTRAVENOUS SOLUTION	Medications	3,200 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	100 mL	No	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Comments:

Occurrences: --

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Comments:

Occurrences: --

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.

9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epinephrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Nursing Orders

ONC NURSING COMMUNICATION 11

Interval: -- Occurrences: --
Comments: Exposure to cold may exacerbate oxaliplatin-induced neuropathy (including pharyngolaryngeal dysesthesia). Encourage patient to keep blanket on their chest and/or throat during oxaliplatin infusion. Educate patient to avoid cold drinks/foods, ice chips or exposure to cold water or air for 7 days after oxaliplatin infusion.

ONC NURSING COMMUNICATION 12

Interval: -- Occurrences: --
Comments: Assess and notify provider for persistent neuropathy (Grade 2).

Nursing Orders

ONC NURSING COMMUNICATION 14

Interval: -- Occurrences: --
Comments: Contact Provider if drug-induced acneiform rash develops and covers more than 25 per cent of the body.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

☒ **HEparin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN
Start: S

Instructions:
Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Day 3

Perform every 1 day x1

Appointment Requests

**ONC PUMP DISCONNECT APPOINTMENT
REQUEST**

Interval: -- Occurrences: --

Discharge Nursing Orders

DISCONNECT CONTINUOUS INFUSION PUMP

Interval: -- Occurrences: --
Comments: Disconnect patient from continuous infusion pump.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Post-Medications

☐ **pegfilgrastim (NEULASTA) on-body injection
kit 6 mg**

Dose: 6 mg Route: subcutaneous once for 1 dose

Start: S End: S

Instructions:

Apply to intact, nonirritated skin on the back of
the arm or abdomen (only use the back of the
arm if caregiver is available to monitor
On-body injection status).