Protocol Index

OP GEMCITABINE / CISPLATIN (SPLIT DOSE)

Types: ONCOLOGY TREATMENT
Synonyms: GEMCITABINE , CISPLATIN , GEM, PLATINOL, GEMZAR, CIS, BLADDER

Cycles 1 to 4 Repeat 4 times Cycle length: 21 days

Day 1 Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST
Interval: -- Occurrences: --

Labs

☑ **COMPREHENSIVE METABOLIC PANEL**
Interval: -- Occurrences: --

☑ **CBC WITH PLATELET AND DIFFERENTIAL**
Interval: -- Occurrences: --

☑ **MAGNESIUM LEVEL**
Interval: -- Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39
Interval: -- Occurrences: --
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)
- Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- Protocol applies only to same day lab value.
- Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40
Interval: -- Occurrences: --
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)
- Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- Protocol applies only to same day lab value.
- Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: --
Occurrences: --
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL | Route: intravenous | PRN
Start: S

Pre-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL | Route: intravenous | once @ 500 mL/hr for 1 dose
Start: S

Pre-Medications

☑ **palonosetron (ALOXI) injection 0.25 mg**

Dose: 0.25 mg | Route: intravenous | once for 1 dose
Start: S | End: S 1:45 PM

☑ **dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% IVPB**

Dose: 12 mg | Route: intravenous | once over 15 Minutes for 1 dose
Start: S

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

☑ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg | Route: intravenous | once over 30 Minutes for 1 dose
Start: S | End: S

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

☐ **netupitant-palonosetron (AKYNZEO) 300-0.5 mg per capsule 1 capsule**

Dose: 1 capsule | Route: oral | once for 1 dose
Start: S | End: S 5:30 PM
Instructions:
Administer approximately 1 hour prior to chemotherapy.

☐ **ondansetron (ZOFRAN), dexamethasone (DECADRON) in sodium chloride 0.9% 50 mL IVPB**

Dose: -- Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:42 AM

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ONDANSETRON HCL 2 MG/ML INTRAVENOUS SOLUTION	Medications		Yes	No
DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications		Yes	No
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

Chemotherapy

gemcitabine (GEMZAR) 1,000 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 1,000 mg/m2 Route: intravenous once over 30 Minutes for 1 dose
Offset: 1.5 Hours

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
GEMCITABINE 200 MG/5.26 ML (38 MG/ML) INTRAVENOUS SOLUTION	Medications	1,000 mg/m2	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

CISplatin (PLATINOL) 35 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 35 mg/m2 Route: intravenous once over 1 Hours for 1 dose
Offset: 2 Hours

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
CISPLATIN 1 MG/ML INTRAVENOUS SOLUTION	Medications	35 mg/m2	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

Post-Hydration

○ **sodium chloride 0.9 % infusion 1,000 mL**

Dose: 1,000 mL | Route: intravenous | once @ 500 mL/hr for 1 dose
Offset: 3 Hours

Instructions:
Following chemotherapy.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76
Interval: -- | Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

☑ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL | Route: intravenous | PRN

☑ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units | Route: intra-catheter | once PRN
Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

Day 8 Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST
Interval: -- | Occurrences: --

Labs

☐ **COMPREHENSIVE METABOLIC PANEL**

Interval: -- | Occurrences: --

☑ **CBC WITH PLATELET AND DIFFERENTIAL**

Interval: -- | Occurrences: --

☐ **MAGNESIUM LEVEL**

Interval: -- | Occurrences: --

☐ **URINALYSIS, AUTOMATED WITH MICROSCOPY**

Interval: -- | Occurrences: --

Outpatient Electrolyte Replacement Protocol

TREATMENT CONDITIONS 39
Interval: -- | Occurrences: --
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP
- o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO
- o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO
- o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

Interval: -- Occurrences: --

Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)

- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
- o Protocol applies only to same day lab value.
- o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
- o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
- o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
- o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 8

Interval: -- Occurrences: --

Comments: On Day 8 of Gemcitabine: HOLD and notify provider if Platelets LESS than 75,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

Start: S

Pre-Hydration

sodium chloride 0.9 % infusion 1,000 mL

Dose: 1,000 mL Route: intravenous once @ 500 mL/hr for 1 dose

Start: S

Pre-Medications

☑ **palonosetron (ALOXI) injection 0.25 mg**

Dose: 0.25 mg Route: intravenous once for 1 dose

Start: S End: S 1:45 PM

☑ **dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% IVPB**

Dose: 12 mg Route: intravenous once over 15 Minutes for 1 dose

Start: S

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes

☑ **aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose

Start: S End: S

Ingredients: **Name** **Type** **Dose** **Selected** **Adds Vol.**

APREPITANT 7.2 MG/ML INTRAVENOUS EMULSION	Medications	130 mg	Main Ingredient	Yes
DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

☐ **netupitant-palonosetron (AKYNZEO) 300-0.5 mg per capsule 1 capsule**
Dose: 1 capsule Route: oral once for 1 dose
Start: S End: S 5:30 PM
Instructions:
Administer approximately 1 hour prior to chemotherapy.

☐ **ondansetron (ZOFRAN), dexamethasone (DECADRON) in sodium chloride 0.9% 50 mL IVPB**
Dose: -- Route: intravenous once over 15 Minutes for 1 dose
Start: S End: S 11:42 AM

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
ONDANSETRON HCL 2 MG/ML INTRAVENOUS SOLUTION	Medications		Yes	No
DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications		Yes	No
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

Chemotherapy

gemcitabine (GEMZAR) 1,000 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB
Dose: 1,000 mg/m2 Route: intravenous once over 30 Minutes for 1 dose
Offset: 30 Minutes

Ingredients:

Name	Type	Dose	Selected	Adds Vol.
GEMCITABINE 200 MG INTRAVENOUS SOLUTION	Medications	1,000 mg/m2	Main Ingredient	Yes
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes

CISplatin (PLATINOL) 35 mg/m2 in sodium chloride 0.9 % 250 mL chemo IVPB

Dose: 35 mg/m2 Route: intravenous once over 1 Hours for 1 dose
Offset: 1 Hours

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	CISPLATIN 1 MG/ML INTRAVENOUS SOLUTION	Medications	35 mg/m2	Main Ingredient	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base		Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base		No	Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76
Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

☑ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

☑ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN
Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.