

OP INOTUZUMAB OZOGAMICIN INDUCTION

Types: ONCOLOGY TREATMENT

Synonyms: BESPONSA, INDUCTION, BESPONSA INDUCTION

Cycle 1	Repeat 1 time	Cycle length: 21 days
Day 1 Perform every 1 day x1		
Appointment Requests		
INFUSION APPOINTMENT REQUEST Interval: -- Occurrences: --		
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> MAGNESIUM LEVEL Interval: -- Occurrences: --		
Line Flush		
sodium chloride 0.9 % flush 20 mL Dose: 20 mL Route: intravenous PRN Start: S		
Nursing Orders		
sodium chloride 0.9 % infusion 250 mL Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose Start: S Instructions: To keep vein open.		
Hydration		
sodium chloride 0.9 % infusion 500 mL Dose: 500 mL Route: intravenous once @ 500 mL/hr for 1 dose Start: S		
Pre-Medications		
acetaminophen (TYLENOL) tablet 650 mg Dose: 650 mg Route: oral once for 1 dose Start: S Instructions: Administer 30 minutes prior to chemotherapy.		
Pre-Medications		
diphenhydramine (BENADRYL) tablet 25 mg Dose: 25 mg Route: oral once for 1 dose Start: S Instructions: Administer 30 minutes prior to chemotherapy.		
Pre-Medications		
hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg Dose: 100 mg Route: intravenous once for 1 dose Instructions: Administer via slow IV push 30 minutes prior to chemotherapy.		
Chemotherapy		

inotuzumab ozogamicin (BESPO NSA) 0.8 mg/m² in sodium chloride 0.9% 50 mL IVPB (RESTRICTED)

Dose: 0.8 mg/m²

Route: intravenous

once over 1 Hours for 1 dose

Offset: 30 Minutes

Instructions:

Do not mix with or administer as an infusion with other drugs. If refrigerated, allow admixtures to warm to room temperature for approximately 1 hour prior to administration.

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

INOTUZUMAB
OZOGAMICIN 0.9
MG(0.25 MG/ML
INITIAL
CONCENTRATION)
IV SOLN
SODIUM
CHLORIDE 0.9 %
INTRAVENOUS
SOLUTION

Medications

0.8
mg/m²

Main

Ingredient

Yes

QS Base

50 mL

Yes

Yes

Nursing Orders

ONC NURSING COMMUNICATION 2

Interval: --

Occurrences: --

Comments:

Observe for symptoms of infusion reaction during and for at least 1 hour after the end of the infusion.

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxvaen at 2 L per minute via nasal cannula. Titrate to

- maintain O2 saturation of greater than or equal to 92%.
- 6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
- 7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
- 8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
- 9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

- Interval: -- Occurrences: --
- Comments: Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)
1. Stop the infusion.
 2. Notify the CERT team and treating physician immediately.
 3. Place the patient on continuous monitoring.
 4. Obtain vital signs.
 5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
 6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
 7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
 8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
 9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
 10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous PRN
Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg Route: oral PRN
Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20 mg

Dose: 20 mg Route: intravenous PRN
Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --

Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Day 8

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

MAGNESIUM LEVEL

Interval: -- Occurrences: --

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Start: S

Instructions:

To keep vein open.

Hydration

sodium chloride 0.9 % infusion 500 mL

Dose: 500 mL Route: intravenous once @ 500 mL/hr for 1 dose

Start: S

Pre-Medications

acetaminophen (TYLENOL) tablet 650 mg

Dose: 650 mg Route: oral once for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

Pre-Medications

diphenhydrAMINE (BENADRYL) tablet 25 mg

Dose: 25 mg Route: oral once for 1 dose

Start: S

Instructions:

Administer 30 minutes prior to chemotherapy.

Pre-Medications

**hydrocortisone sodium succinate
(Solu-CORTEF) injection 100 mg**

Dose: 100 mg Route: intravenous once for 1 dose

Instructions:
Administer via slow IV push 30 minutes prior to chemotherapy.

Chemotherapy

inotuzumab ozogamicin (BESPONSA) 0.5 mg/m² in sodium chloride 0.9% 50 mL IVPB (RESTRICTED)

Dose: 0.5 mg/m² Route: intravenous once over 1 Hours for 1 dose
Offset: 30 Minutes

Instructions:
Do not mix with or administer as an infusion with other drugs. If refrigerated, allow admixtures to warm to room temperature for approximately 1 hour prior to administration.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	INOTUZUMAB	Medications	0.5	Main	Yes
	OZOGAMICIN 0.9 MG(0.25 MG/ML INITIAL CONCENTRATION) IV SOLN		mg/m ²	Ingredient	
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	50 mL	Yes	Yes

Nursing Orders

ONC NURSING COMMUNICATION 2

Interval: -- Occurrences: --
Comments: Observe for symptoms of infusion reaction during and for at least 1 hour after the end of the infusion.

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --
Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: -- Occurrences: --
Comments: Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or

back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25

mg

Dose: 25 mg

Route: intravenous

PRN

Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg

Route: oral

PRN

Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20

mg

Dose: 20 mg

Route: intravenous

PRN

Start: S

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg

Route: intravenous

PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg

Route: intravenous

PRN

Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT

injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
 Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
 Comments: Discontinue IV.

Discharge Nursing Orders

 sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

 HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
 Start: S
 Instructions:
 Concentration: 100 units/mL. Heparin flush for
 Implanted Vascular Access Device
 maintenance.

Day 15

Perform every 1 day x1

Appointment Requests

INFUSION APPOINTMENT REQUEST

Interval: -- Occurrences: --

Labs

 COMPREHENSIVE METABOLIC PANEL

Interval: -- Occurrences: --

 CBC WITH PLATELET AND DIFFERENTIAL

Interval: -- Occurrences: --

 MAGNESIUM LEVEL

Interval: -- Occurrences: --

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions:
 To keep vein open.

Hydration

sodium chloride 0.9 % infusion 500 mL

Dose: 500 mL Route: intravenous once @ 500 mL/hr for 1 dose
 Start: S

Pre-Medications

acetaminophen (TYLENOL) tablet 650 mg

Dose: 650 mg Route: oral once for 1 dose
 Start: S
 Instructions:
 Administer 30 minutes prior to chemotherapy.

Pre-Medications

diphenhydrAMINE (BENADRYL) tablet 25 mg

Dose: 25 mg Route: oral once for 1 dose

Start: S
Instructions:
Administer 30 minutes prior to chemotherapy.

Pre-Medications

hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous once for 1 dose

Instructions:
Administer via slow IV push 30 minutes prior to chemotherapy.

Chemotherapy

inotuzumab ozogamicin (BESPONSA) 0.5 mg/m2 in sodium chloride 0.9% 50 mL IVPB (RESTRICTED)

Dose: 0.5 mg/m2 Route: intravenous once over 1 Hours for 1 dose
Offset: 30 Minutes

Instructions:
Do not mix with or administer as an infusion with other drugs. If refrigerated, allow admixtures to warm to room temperature for approximately 1 hour prior to administration.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	INOTUZUMAB	Medications	0.5	Main	Yes
	OZOGAMICIN 0.9 MG(0.25 MG/ML INITIAL CONCENTRATION) IV SOLN		mg/m2	Ingredient	
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	50 mL	Yes	Yes

Nursing Orders

ONC NURSING COMMUNICATION 2

Interval: -- Occurrences: --
Comments: Observe for symptoms of infusion reaction during and for at least 1 hour after the end of the infusion.

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: -- Occurrences: --
Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)
1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydramine (BENADRYL) injection 25

mg

Dose: 25 mg

Route: intravenous

PRN

Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg

Route: oral

PRN

Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20

mg

Dose: 20 mg

Route: intravenous

PRN

Start: S

hydrocortisone sodium succinate

(Solu-CORTEF) injection 100 mg

Dose: 100 mg Route: intravenous PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg Route: intravenous PRN
Start: S

epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.