

OP IRINOTECAN / CETUXIMAB / VEMURAFENIB (EVERY 14 DAYS)

Types: ONCOLOGY TREATMENT

Synonyms: BRAF, V600E, IRINOTECAN, IRIN, CAMPTOSAR, CAMP, CET, CETUXIMAB, VEMURAFENIB, ZELBORAF, COLORECTAL, VIC

Take-Home Medications	Repeat 1 time	Cycle length: 1 day
Day 1		Perform every 1 day x1
Provider Communication		
ONC PROVIDER COMMUNICATION		
Interval: --	Occurrences: --	
Comments:	Presence of V600 mutation of BRAF gene must be documented by a CLIA-approved laboratory prior to initiation of Vemurafenib therapy.	
Take-Home Medications Prior to Treatment		
vemurafenib (ZELBORAF) 240 mg tablet tablet		
Dose: 960 mg	Route: oral	2 times daily
Dispense: 224 tablet	Refills: 2	
Start: S	End: S+14	
Instructions:	Take 4 tablets (total 960 mg) by mouth twice daily on days 1 to 14.	
Cycles 1 to 6	Repeat 6 times	Cycle length: 14 days
Day 1		Perform every 1 day x1
Appointment Requests		
INFUSION APPOINTMENT REQUEST		
Interval: --	Occurrences: --	
Labs		
<input checked="" type="checkbox"/> COMPREHENSIVE METABOLIC PANEL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> CBC WITH PLATELET AND DIFFERENTIAL		
Interval: --	Occurrences: --	
<input checked="" type="checkbox"/> MAGNESIUM LEVEL		
Interval: --	Occurrences: --	
Nursing Orders		
ONC VERIFY PATIENT TAKING HOME MEDS		
Interval: --	Occurrences: --	
Comments:	Verify that patient is taking home meds as directed.	
Outpatient Electrolyte Replacement Protocol		
TREATMENT CONDITIONS 39		
Interval: --	Occurrences: --	
Comments:	Potassium (Normal range 3.5 to 5.0mEq/L) <ul style="list-style-type: none">o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NPo Protocol applies only to same day lab value.o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NPo Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or POo Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or POo Serum potassium 3.5 mEq/L or greater, do not give potassium replacemento If patient meets criteria, order SmartSet called "Outpatient	

Electrolyte Replacement"

- o Sign electrolyte replacement order as Per protocol: cosign required

TREATMENT CONDITIONS 40

- Interval: -- Occurrences: --
 Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)
- o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP
 - o Protocol applies only to same day lab value.
 - o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP
 - o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV
 - o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV
 - o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement
 - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
 - o Sign electrolyte replacement order as Per protocol: cosign required

Provider Communication

ONC PROVIDER COMMUNICATION 2

- Interval: -- Occurrences: --
 Comments: Tumor KRAS gene status should be determined prior to initiation of therapy. KRAS type: Please Push F2:115540219.

Nursing Orders

TREATMENT CONDITIONS 4

- Interval: -- Occurrences: --
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000; Total Bilirubin GREATER than 1.5; ALT/AST GREATER than 3 times upper normal limit; or Serum Creatinine GREATER than 1.2

Line Flush

sodium chloride 0.9 % flush 20 mL
 Dose: 20 mL Route: intravenous PRN
 Start: S

Nursing Orders

sodium chloride 0.9 % infusion 250 mL
 Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose
 Start: S
 Instructions:
 To keep vein open.

Pre-Medications

ondansetron (ZOFTRAN) 16 mg, dexamethasone (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB
 Dose: -- Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:30 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
	DEXAMETHASONE 4 MG/ML	Medications	12 mg	Yes	No

INJECTION SOLUTION SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

ondansetron (ZOFRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
Start: S End: S 11:30 AM

dexamethasone (DECADRON) tablet 12 mg

Dose: 12 mg Route: oral once for 1 dose
Start: S

palonosetron (ALOXI) injection 0.25 mg

Dose: 250 mcg Route: intravenous once for 1 dose
Start: S End: S 3:00 PM
Instructions:
For OUTPATIENT use only.

aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB

Dose: 130 mg Route: intravenous once over 30 Minutes for 1 dose
Start: S End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Pre-Medications

atropine injection 0.25 mg

Dose: 0.25 mg Route: intravenous once PRN
Start: S

Pre-Medications

diphenhydrAMINE (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous once for 1 dose
Start: S
Instructions:
Give 30 minutes prior to cetuximab.

Chemotherapy

cetuximab (ERBITUX) 500 mg/m2 in 0 mL

Dose: 500 mg/m2 Route: intravenous once over 120 Minutes for 1 dose
Offset: 30 Minutes

Instructions:
Administer with low protein binding 0.22 micron filter. Do not shake. Do not mix with other medications. Flush IV line with NS at the end

of infusion.

1st infusion: Infuse first 10 mL over 10 minutes and observe patient for 30 minutes for allergic reactions if infusion tolerated, infuse loading dose over 120 minutes.

Rate of infusion not to exceed 10 mg/minute (5 mL/minute)

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	CETUXIMAB 100 MG/50 ML INTRAVENOUS SOLUTION	Medications	500 mg/m2	Main Ingredient	Yes

Chemotherapy

irinotecan (CAMPTOSAR) 180 mg/m2 in dextrose 5% 500 mL chemo IVPB

Dose: 180 mg/m2 Route: intravenous once over 90 Minutes for 1 dose
Offset: 2.5 Hours

Instructions:
Protect from light

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	IRINOTECAN 100 MG/5 ML INTRAVENOUS SOLUTION	Medications	180 mg/m2	Main Ingredient	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	No	Yes

Hematology & Oncology Hypersensitivity Reaction Standing Order

ONC NURSING COMMUNICATION 82

Interval: --

Occurrences: --

Comments:

Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)

1. Stop the infusion.
2. Place the patient on continuous monitoring.
3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 4

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea.

vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

ONC NURSING COMMUNICATION 83

Interval: --

Comments:

Occurrences: --

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O₂ saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O₂ saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

diphenhydrAMINE (BENADRYL) injection 25

mg

Dose: 25 mg

Route: intravenous

PRN

Start: S

fexofenadine (ALLEGRA) tablet 180 mg

Dose: 180 mg

Route: oral

PRN

Start: S

famotidine (PEPCID) 20 mg/2 mL injection 20

mg

Dose: 20 mg

Route: intravenous

PRN

Start: S

hydrocortisone sodium succinate

(Solu-CORTEF) injection 100 mg

Dose: 100 mg

Route: intravenous

PRN

dexamethasone (DECADRON) injection 4 mg

Dose: 4 mg

Route: intravenous

PRN

Start: S

**epINEPHrine (ADRENALIN) 1 mg/10 mL ADULT
injection syringe 0.3 mg**

Dose: 0.3 mg Route: subcutaneous PRN
Start: S

Nursing Orders

ONC NURSING COMMUNICATION 14

Interval: -- Occurrences: --
Comments: Contact Provider if drug-induced acneiform rash develops and covers
more than 25 per cent of the body.

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: -- Occurrences: --
Comments: Discontinue IV.

Discharge Nursing Orders

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN

HEParin, porcine (PF) injection 500 Units

Dose: 500 Units Route: intra-catheter once PRN
Start: S
Instructions:
Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.