

OP IXABEPILONE

Types: ONCOLOGY TREATMENT

Synonyms: IXABEPILONE , IXEMPRA, ICS, ISA, ISE, X, BREAST

Cycles 1 to 4		Repeat 4 times	Cycle length: 21 days
Day 1		Perform every 1 day x1	
Appointment Requests			
		INFUSION APPOINTMENT REQUEST	
		Interval: --	Occurrences: --
Labs			
		☒ COMPREHENSIVE METABOLIC PANEL	
		Interval: --	Occurrences: --
		☒ CBC WITH PLATELET AND DIFFERENTIAL	
		Interval: --	Occurrences: --
		☒ MAGNESIUM LEVEL	
		Interval: --	Occurrences: --
		☐ URINALYSIS, AUTOMATED WITH MICROSCOPY	
		Interval: --	Occurrences: --
		☐ CANCER ANTIGEN 27-29 (CA BR)	
		Interval: --	Occurrences: --
Outpatient Electrolyte Replacement Protocol			
		TREATMENT CONDITIONS 39	
		Interval: --	Occurrences: --
		Comments:	Potassium (Normal range 3.5 to 5.0mEq/L) - o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP - o Protocol applies only to same day lab value. - o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP - o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO - o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO - o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement - o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement" - o Sign electrolyte replacement order as Per protocol: cosign required
		TREATMENT CONDITIONS 40	
		Interval: --	Occurrences: --
		Comments:	Magnesium (Normal range 1.6 to 2.6mEq/L) - o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP - o Protocol applies only to same day lab value. - o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP - o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV - o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV - o Serum Magnesium 1.6 mEq/L or areater, do not aive

magnesium replacement

- o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"
- o Sign electrolyte replacement order as Per protocol: cosign required

Nursing Orders

TREATMENT CONDITIONS 7

Interval: -- Occurrences: --
 Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000.

Line Flush

sodium chloride 0.9 % flush 20 mL

Dose: 20 mL Route: intravenous PRN
 Start: S

Nursing Orders

lactated ringer's infusion 250 mL

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose

Instructions:
 To keep vein open

Pre-Medications

☒ ondansetron (ZOFRAN) injection 8 mg

Dose: 8 mg Route: intravenous once for 1 dose
 Start: S End: S 11:15 AM

☐ ondansetron (ZOFRAN) tablet 16 mg

Dose: 16 mg Route: oral once for 1 dose
 Start: S

☐ ondansetron (ZOFRAN) 16 mg in dextrose 5% 50 mL IVPB

Dose: 16 mg Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:00 AM

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Main Ingredient	No
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes

Pre-Medications

☒ diphenhydramine (BENADRYL) injection 25 mg

Dose: 25 mg Route: intravenous once for 1 dose
 Start: S
 Instructions:
 Administer via slow IV push 30 minutes prior to chemotherapy.

☐ diphenhydramine (BENADRYL) 50 mg in sodium chloride 0.9% 50 mL IVPB

Dose: 50 mg Route: intravenous once over 15 Minutes for 1 dose
 Start: S End: S 11:45 AM
 Instructions:
 Administer 30 minutes prior to chemotherapy.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	DIPHENHYDRAMIN E 50 MG/ML INJECTION SOLUTION	Medications	50 mg	Main Ingredient	No
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Yes	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base	50 mL	No	Yes
☐ diphenhydrAMINE (BENADRYL) tablet 25 mg					
Dose: 25 mg		Route: oral	once for 1 dose Offset: 0 Hours		
Instructions: Administer 30 minutes prior to chemotherapy.					
☐ diphenhydrAMINE (BENADRYL) tablet 50 mg					
Dose: 50 mg		Route: oral	once for 1 dose Offset: 0 Hours		
Instructions: Administer 30 minutes prior to chemotherapy.					
☒ famotidine (PEPCID) injection 20 mg					
Dose: 20 mg		Route: intravenous	once for 1 dose Offset: 0 Hours		
Instructions: Administer 30 minutes prior to chemotherapy.					
☐ famotidine (PEPCID) tablet 20 mg					
Dose: 20 mg		Route: oral	once for 1 dose Offset: 0 Hours		
Instructions: Administer 30 minutes prior to chemotherapy.					
☐ acetaminophen (TYLENOL) tablet 650 mg					
Dose: 650 mg		Route: oral	once for 1 dose Offset: 0 Hours		
Instructions: Administer 30 minutes prior to chemotherapy.					
Supportive Care					
☐ LORAZepam (ATIVAN) injection 1 mg					
Dose: 1 mg Start: S		Route: intravenous	once PRN		
☐ LORAZepam (ATIVAN) tablet 1 mg					
Dose: 1 mg Start: S		Route: oral	once PRN		
Antiemetics					
☐ promethazine (PHENERGAN) injection 12.5 mg					
Dose: 12.5 mg Start: S		Route: injection	once PRN		
Chemotherapy					
ixabepilone (IXEMPRA) 40 mg/m2 in lactated ringer's 250 mL chemo IVPB					

Dose: 40 mg/m2 Route: intravenous once over 3 Hours for 1 dose
Offset: 30 Minutes

Instructions:

Maximum dose = 88 mg. Dose must be
infused through a 0.2 micron inline filter. Use
DEHP-free bag & tubing.

Rule-Based Template: RULE ONCBCN
IXABEPILONE 40MG/M2

Conditions:

BSA < 2.2 m2

BSA > = 2.2 m2

Modifications:

Set dose to 40 mg/m2

Set dose to 88 mg

Ingredients:

Name

Type

Dose

Selected

Adds Vol.

IXABEPILONE 45
MG INTRAVENOUS
SOLUTION
LACTATED
RINGERS
INTRAVENOUS
SOLUTION

Medications

40 mg/m2

Main

Yes

Ingredient

QS Base

Yes

Yes

Discharge Nursing Orders

ONC NURSING COMMUNICATION 76

Interval: --

Occurrences: --

Comments:

Discontinue IV.

Discharge Nursing Orders

☒ **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL

Route: intravenous

PRN

☒ **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units

Route: intra-catheter

once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for
Implanted Vascular Access Device
maintenance.

Post-Medications

☐ **pegfilgrastim (NEULASTA) on-body injection
kit 6 mg**

Dose: 6 mg

Route: subcutaneous

once for 1 dose

Start: S

End: S

Instructions:

Apply to intact, nonirritated skin on the back of
the arm or abdomen (only use the back of the
arm if caregiver is available to monitor
On-body injection status).