

# OP MFOLFIRINOX (EVERY 2 WEEKS) / ADJUVANT FOR PANCREATIC CANCER

Types: ONCOLOGY TREATMENT

Synonyms: MFOLFIRINOX , FLUOROURACIL , LEUCOVORIN, OXALIPLATIN , IIRINOTECAN , AGE, CAMPTOSAR, ADRUCIL, ELOXATIN, FLOW, IRENE, PANCREATIC, PANCREAS, ADJUVANT FOR PANC CANCER

<b>Cycles 1 to 12</b>	Repeat 12 times	Cycle length: 14 days
<b>Day 1</b>		Perform every 1 day x1
Appointment Requests		
<b>INFUSION APPOINTMENT REQUEST</b>		
Interval: -- Occurrences: --		
Labs		
<input checked="" type="checkbox"/> <b>COMPREHENSIVE METABOLIC PANEL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>CBC WITH PLATELET AND DIFFERENTIAL</b>		
Interval: -- Occurrences: --		
<input checked="" type="checkbox"/> <b>MAGNESIUM LEVEL</b>		
Interval: -- Occurrences: --		
Outpatient Electrolyte Replacement Protocol		
<b>TREATMENT CONDITIONS 39</b>		
Interval: -- Occurrences: --		
Comments: Potassium (Normal range 3.5 to 5.0mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum potassium less than 3.0mEq/L, give 40mEq KCL IV or PO and contact MD/NP		
o Serum potassium 3.0 to 3.2mEq/L, give 40mEq KCL IV or PO		
o Serum potassium 3.3 to 3.4mEq/L, give 20mEq KCL IV or PO		
o Serum potassium 3.5 mEq/L or greater, do not give potassium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign required		
<b>TREATMENT CONDITIONS 40</b>		
Interval: -- Occurrences: --		
Comments: Magnesium (Normal range 1.6 to 2.6mEq/L)		
o Protocol applies for SCr less than 1.5. Otherwise, contact MD/NP		
o Protocol applies only to same day lab value.		
o Serum Magnesium less than 1.0mEq/L, give 2 gram magnesium sulfate IV and contact MD/NP		
o Serum Magnesium 1.0 to 1.2mEq/L, give 2 gram magnesium sulfate IV		
o Serum Magnesium 1.3 to 1.5mEq/L, give 1 gram magnesium sulfate IV		
o Serum Magnesium 1.6 mEq/L or greater, do not give magnesium replacement		
o If patient meets criteria, order SmartSet called "Outpatient Electrolyte Replacement"		
o Sign electrolyte replacement order as Per protocol: cosign		

required

Nursing Orders

**TREATMENT CONDITIONS 4**

Interval: -- Occurrences: --  
Comments: HOLD and notify provider if ANC LESS than 1000; Platelets LESS than 100,000; Total Bilirubin GREATER than 1.5; ALT/AST GREATER than 3 times upper normal limit; or Serum Creatinine GREATER than 1.2

Line Flush

**dextrose 5% flush syringe 20 mL**

Dose: 20 mL Route: intravenous PRN  
Start: S  
Instructions:  
Administer ONLY for Oxaliplatin.

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN  
Start: S  
Instructions:  
Do NOT administer with Oxaliplatin.

Nursing Orders

**dextrose 5% infusion 250 mL**

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
Start: S  
Instructions:  
To keep vein open for Oxaliplatin.

**sodium chloride 0.9 % infusion 250 mL**

Dose: 250 mL Route: intravenous once @ 30 mL/hr for 1 dose  
Start: S  
Instructions:  
To keep vein open. Do NOT administer with Oxaliplatin.

Pre-Medications

**ondansetron (ZOFRAN) 16 mg, dexamethasone**

- (DECADRON) 12 mg in sodium chloride 0.9% 50 mL IVPB**

Dose: -- Route: intravenous once over 15 Minutes for 1 dose  
Start: S End: S 11:30 AM

**Ingredients:**

Name	Type	Dose	Selected	Adds Vol.
ONDANSETRON HCL (PF) 4 MG/2 ML INJECTION SOLUTION	Medications	16 mg	Yes	No
DEXAMETHASONE 4 MG/ML INJECTION SOLUTION	Medications	12 mg	Yes	No
SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	Base	50 mL	Always	Yes
DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	Base		No	Yes

- ondansetron (ZOFRAN) tablet 16 mg**

Dose: 16 mg Route: oral once for 1 dose  
Start: S End: S 11:30 AM

**dexamethasone (DECADRON) tablet 12 mg**

Dose: 12 mg                      Route: oral                      once for 1 dose  
Start: S

**palonosetron (ALOXI) injection 0.25 mg**

Dose: 250 mcg                      Route: intravenous                      once for 1 dose  
Start: S                      End: S 3:00 PM  
Instructions:  
For OUTPATIENT use only.

**aprepitant (CINVANTI) 130 mg in dextrose (NON-PVC) 5% 130 mL IVPB**

Dose: 130 mg                      Route: intravenous                      once over 30 Minutes for 1 dose  
Start: S                      End: S

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	APREPITANT 7.2 MG/ML	Medications	130 mg	Main Ingredient	Yes
	INTRAVENOUS EMULSION				
	DEXTROSE 5 % IN WATER (D5W) IV SOLP (EXCEL; NON-PVC)	Base	130 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % IV SOLP (EXCEL;NON-PVC)	Base	130 mL	No	Yes

Pre-Medications

**atropine injection 0.25 mg**

Dose: 0.25 mg                      Route: intravenous                      once PRN  
Start: S

Supportive Care

**LORAZepam (ATIVAN) injection 1 mg**

Dose: 1 mg                      Route: intravenous                      once PRN  
Start: S

**LORAZepam (ATIVAN) tablet 1 mg**

Dose: 1 mg                      Route: oral                      once PRN  
Start: S

Chemotherapy

**OXALIplatin (ELOXATIN) 85 mg/m2 in dextrose 5% 500 mL chemo IVPB**

Dose: 85 mg/m2                      Route: intravenous                      once over 2 Hours for 1 dose  
Offset: 30 Minutes

Instructions:  
Irritant - avoid extravasation. Flush line with D5W before and after oxaliplatin infusion.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	OXALIPLATIN 100 MG/20 ML	Medications	85 mg/m2	Main Ingredient	Yes
	INTRAVENOUS SOLUTION				
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes

**leucovorin 400 mg/m2 in dextrose 5% 250 mL IVPB**

Dose: 400 ma/m2                      Route: intravenous                      once over 2 Hours for 1 dose

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	LEUCOVORIN	Medications	400	Main	Yes
	CALCIUM 350 MG SOLUTION FOR INJECTION		mg/m2	Ingredient	
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	250 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	250 mL	No	Yes

Offset: 30 Minutes

**irinotecan (CAMPTOSAR) 150 mg/m2 in dextrose 5% 500 mL chemo IVPB**

Dose: 150 mg/m2      Route: intravenous      once over 90 Minutes for 1 dose  
Offset: 2.5 Hours

Instructions:  
Protect from light

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	IRINOTECAN 100 MG/5 ML INTRAVENOUS SOLUTION	Medications	150	Main	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	500 mL	Yes	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	500 mL	No	Yes

**fluorouracil (ADRUCIL) 2,400 mg/m2 in sodium chloride 0.9% 100 mL chemo infusion - AMBULATORY PUMP**

Dose: 2,400 mg/m2      Route: intravenous      once over 46 Hours for 1 dose  
Offset: 4 Hours

Instructions:  
Nurses to chart as GIVEN on the MAR.  
Administer via CADD pump.

Ingredients:	Name	Type	Dose	Selected	Adds Vol.
	FLUOROURACIL 5 GRAM/100 ML INTRAVENOUS SOLUTION	Medications	2,400	Main	Yes
	DEXTROSE 5 % IN WATER (D5W) INTRAVENOUS SOLUTION	QS Base	100 mL	No	Yes
	SODIUM CHLORIDE 0.9 % INTRAVENOUS SOLUTION	QS Base	100 mL	Yes	Yes

**Hematology & Oncology Hypersensitivity Reaction Standing Order**

**ONC NURSING COMMUNICATION 82**

Interval: --      Occurrences: --  
Comments: Grade 1 - MILD Symptoms (cutaneous and subcutaneous symptoms only – itching, flushing, periorbital edema, rash, or runny nose)  
1. Stop the infusion.  
2. Place the patient on continuous monitoring.

3. Obtain vital signs.
4. Administer Normal Saline at 50 mL per hour using a new bag and new intravenous tubing.
5. If greater than or equal to 30 minutes since the last dose of Diphenhydramine, administer Diphenhydramine 25 mg intravenous once.
6. If less than 30 minutes since the last dose of Diphenhydramine, administer Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
7. Notify the treating physician.
8. If no improvement after 15 minutes, advance level of care to Grade 2 (Moderate) or Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 4**

Interval: --

Occurrences: --

Comments:

Grade 2 – MODERATE Symptoms (cardiovascular, respiratory, or gastrointestinal symptoms – shortness of breath, wheezing, nausea, vomiting, dizziness, diaphoresis, throat or chest tightness, abdominal or back pain)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
6. Administer Normal Saline at 150 mL per hour using a new bag and new intravenous tubing.
7. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous), Fexofenadine 180 mg orally and Famotidine 20 mg intravenous once.
8. If no improvement after 15 minutes, advance level of care to Grade 3 (Severe).
9. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

#### **ONC NURSING COMMUNICATION 83**

Interval: --

Occurrences: --

Comments:

Grade 3 – SEVERE Symptoms (hypoxia, hypotension, or neurologic compromise – cyanosis or O2 saturation less than 92%, hypotension with systolic blood pressure less than 90 mmHg, confusion, collapse, loss of consciousness, or incontinence)

1. Stop the infusion.
2. Notify the CERT team and treating physician immediately.
3. Place the patient on continuous monitoring.
4. Obtain vital signs.
5. If heart rate is less than 50 or greater than 120, or blood pressure is less than 90/50 mmHg, place patient in reclined or flattened position.
6. Administer Oxygen at 2 L per minute via nasal cannula. Titrate to maintain O2 saturation of greater than or equal to 92%.
7. Administer Normal Saline at 1000 mL intravenous bolus using a new bag and new intravenous tubing.
8. Administer Hydrocortisone 100 mg intravenous (if patient has allergy to Hydrocortisone, please administer Dexamethasone 4 mg intravenous) and Famotidine 20 mg intravenous once.
9. Administer Epinephrine (1:1000) 0.3 mg subcutaneous.
10. Assess vital signs every 15 minutes until resolution of symptoms or otherwise ordered by covering physician.

**diphenhydrAMINE (BENADRYL) injection 25 mg**

Dose: 25 mg                      Route: intravenous                      PRN  
Start: S

**fexofenadine (ALLEGRA) tablet 180 mg**

Dose: 180 mg                      Route: oral                      PRN  
Start: S

**famotidine (PEPCID) 20 mg/2 mL injection 20 mg**

Dose: 20 mg                      Route: intravenous                      PRN  
Start: S

**hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg**

Dose: 100 mg                      Route: intravenous                      PRN

**dexamethasone (DECADRON) injection 4 mg**

Dose: 4 mg                      Route: intravenous                      PRN  
Start: S

**epiNEPHrine (ADRENALIN) 1 mg/10 mL ADULT injection syringe 0.3 mg**

Dose: 0.3 mg                      Route: subcutaneous                      PRN  
Start: S

**Nursing Orders**

**ONC NURSING COMMUNICATION 11**

Interval: --                      Occurrences: --  
Comments:                      Exposure to cold may exacerbate oxaliplatin-induced neuropathy (including pharyngolaryngeal dysesthesia). Encourage patient to keep blanket on their chest and/or throat during oxaliplatin infusion. Educate patient to avoid cold drinks/foods, ice chips or exposure to cold water or air for 7 days after oxaliplatin infusion.

**ONC NURSING COMMUNICATION 12**

Interval: --                      Occurrences: --  
Comments:                      Assess and notify provider for persistent neuropathy (Grade 2).

**Discharge Nursing Orders**

**ONC NURSING COMMUNICATION 76**

Interval: --                      Occurrences: --  
Comments:                      Discontinue IV.

**Discharge Nursing Orders**

**sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL                      Route: intravenous                      PRN

**HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units                      Route: intra-catheter                      once PRN  
Start: S

Instructions:  
Concentration: 100 units/mL. Heparin flush for Implanted Vascular Access Device maintenance.

**Day 3**

Perform every 1 day x1

**Appointment Requests**

**ONC PUMP DISCONNECT APPOINTMENT REQUEST**

Interval: --                      Occurrences: --

**Discharge Nursing Orders**

**DISCONNECT CONTINUOUS INFUSION PUMP**

Interval: -- Occurrences: --  
Comments: Disconnect patient from continuous infusion pump.

## Discharge Nursing Orders

 **sodium chloride 0.9 % flush 20 mL**

Dose: 20 mL Route: intravenous PRN

 **HEParin, porcine (PF) injection 500 Units**

Dose: 500 Units Route: intra-catheter once PRN

Start: S

Instructions:

Concentration: 100 units/mL. Heparin flush for  
Implanted Vascular Access Device  
maintenance.

## Post-Medications

 **pegfilgrastim (NEULASTA) on-body injection  
kit 6 mg**

Dose: 6 mg Route: subcutaneous once for 1 dose

Start: S

End: S

Instructions:

Apply to intact, nonirritated skin on the back of  
the arm or abdomen (only use the back of the  
arm if caregiver is available to monitor  
On-body injection status).